Rances Blanco

List of Publications by Citations

Source: https://exaly.com/author-pdf/5145368/rances-blanco-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

31 333 9 17 g-index

36 395 ext. papers ext. citations 3.8 avg, IF L-index

#	Paper	IF	Citations
31	Induction of immunogenic apoptosis by blockade of epidermal growth factor receptor activation with a specific antibody. <i>Journal of Immunology</i> , 2011 , 187, 4954-66	5.3	70
30	Biological activity in vitro of anti-epidermal growth factor receptor monoclonal antibodies with different affinities. <i>Hybridoma</i> , 2007 , 26, 423-31		42
29	Immunoreactivity of the 14F7 Mab Raised against N-Glycolyl GM3 Ganglioside in Epithelial Malignant Tumors from Digestive System. <i>ISRN Gastroenterology</i> , 2011 , 2011, 645641		26
28	Human antibodies reactive to NeuGcGM3 ganglioside have cytotoxic antitumor properties. <i>European Journal of Immunology</i> , 2013 , 43, 826-37	6.1	25
27	Immunoreactivity of the 14F7 Mab (Raised against N-Glycolyl GM3 Ganglioside) as a Positive Prognostic Factor in Non-Small-Cell Lung Cancer. <i>Pathology Research International</i> , 2012 , 2012, 235418		22
26	Immunohistochemical Reactivity of the 14F7 Monoclonal Antibody Raised against N-Glycolyl GM3 Ganglioside in Some Benign and Malignant Skin Neoplasms. <i>ISRN Dermatology</i> , 2011 , 2011, 848909		21
25	Prognostic Significance of N-Glycolyl GM3 Ganglioside Expression in Non-Small Cell Lung Carcinoma Patients: New Evidences. <i>Pathology Research International</i> , 2015 , 2015, 132326		19
24	Tissue Reactivity of the 14F7 Mab Raised against N-Glycolyl GM3 Ganglioside in Tumors of Neuroectodermal, Mesodermal, and Epithelial Origin. <i>Journal of Biomarkers</i> , 2013 , 2013, 602417	О	13
23	Sticholysin II-mediated cytotoxicity involves the activation of regulated intracellular responses that anticipates cell death. <i>Biochimie</i> , 2018 , 148, 18-35	4.6	10
22	Immunorecognition of the 14F7 Mab Raised against N-Glycolyl GM3 Ganglioside in Some Normal and Malignant Tissues from Genitourinary System. <i>ISRN Pathology</i> , 2011 , 2011, 1-10		9
21	High-Risk Human Papillomavirus and Tobacco Smoke Interactions in Epithelial Carcinogenesis. <i>Cancers</i> , 2020 , 12,	6.6	8
20	Immunoreactivity of the 14F7 Mab Raised against N-Glycolyl GM3 Ganglioside in Primary Lymphoid Tumors and Lymph Node Metastasis. <i>Pathology Research International</i> , 2013 , 2013, 920972		7
19	Synergistic potentiation of the anti-metastatic effect of anti EGFR mAb by its combination with immunotherapies targeting the ganglioside NGcGM3. <i>Oncotarget</i> , 2018 , 9, 24069-24080	3.3	7
18	Human Papillomavirus 16 E7 Promotes EGFR/PI3K/AKT1/NRF2 Signaling Pathway Contributing to PIR/NF- B Activation in Oral Cancer Cells. <i>Cancers</i> , 2020 , 12,	6.6	7
17	Flow Cytometric Measurement of Aneuploid DNA Content Correlates with High S-Phase Fraction and Poor Prognosis in Patients with Non-Small-Cell Lung Cancer 2013 , 2013, 1-8		5
16	Role of BamHI-A Rightward Frame 1 in Epstein-Barr Virus-Associated Epithelial Malignancies. <i>Biology</i> , 2020 , 9,	4.9	5
15	Functional expression of human-epidermal-growth-factor receptor in a melanoma cell line. <i>Biotechnology and Applied Biochemistry</i> , 2007 , 48, 21-7	2.8	4

LIST OF PUBLICATIONS

14	Preclinical Efficacy of Nimotuzumab, an Anti-Egfr Monoclonal Antibody as a Single Agent Therapy in Human GBM u87mg Xenografts. <i>Journal of Cancer Therapy</i> , 2012 , 03, 245-255	0.2	4	
13	Role of Epstein-Barr Virus and Human Papillomavirus Coinfection in Cervical Cancer: Epidemiology, Mechanisms and Perspectives. <i>Pathogens</i> , 2020 , 9,	4.5	4	
12	Interplay between Epstein-Barr virus infection and environmental xenobiotic exposure in cancer. <i>Infectious Agents and Cancer</i> , 2021 , 16, 50	3.5	4	
11	Aberrant expression of N-glycolyl GM3 ganglioside is associated with the aggressive biological behavior of human sarcomas. <i>BMC Cancer</i> , 2019 , 19, 556	4.8	3	
10	Immunohistochemical Characterization of Three Monoclonal Antibodies Raised against the Epidermal Growth Factor and Its Receptor in Non-Small-Cell Lung Cancer: Their Potential Use in the Selection of Patients for Immunotherapy. <i>Journal of Biomarkers</i> , 2013 , 2013, 627845	0	3	
9	Human Papillomavirus in Breast Carcinogenesis: A Passenger, a Cofactor, or a Causal Agent?. <i>Biology</i> , 2021 , 10,	4.9	3	
8	Merkel cell polyomavirus detected in head and neck carcinomas from Chile. <i>Infectious Agents and Cancer</i> , 2020 , 15, 4	3.5	2	
7	Epidermal growth factor receptor modulates the tumorigenic potential of melanoma. <i>Frontiers in Bioscience - Landmark</i> , 2009 , 14, 159-66	2.8	2	
6	High-Risk Human Papillomavirus and Epstein-Barr Virus Coinfection: A Potential Role in Head and Neck Carcinogenesis <i>Biology</i> , 2021 , 10,	4.9	2	
5	Tissue Expression of Low and High Molecular Weight Cytokeratins in Lung Carcinoma Sections: Its Correlation with Some Clinic-Pathological Features. <i>Journal of Histology</i> , 2014 , 2014, 1-9		1	
4	Tumor Expression of the Carcinoembryonic Antigen Correlates with High Mitotic Activity and Cell Pleomorphism Index in Lung Carcinoma. <i>Journal of Histology</i> , 2013 , 2013, 1-8		1	
3	Phenotypic heterogeneity in the NCI-H125 cell line affects biological activity using the epidermal growth factor receptor as target. <i>Acta Pharmaceutica</i> , 2012 , 62, 581-91	3.2	1	
2	Epstein B arr Virus Association with Breast Cancer: Evidence and Perspectives. <i>Biology</i> , 2022 , 11, 799	4.9	О	
1	Characterization of High-Risk HPV/EBV Co-Presence in Pre-Malignant Cervical Lesions and	4.9		