Lluis Catasus

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/514643/publications.pdf Version: 2024-02-01



Ι τιμο Ολτλομο

#	Article	IF	CITATIONS
1	hMLH1 Promoter Hypermethylation Is an Early Event in Human Endometrial Tumorigenesis. American Journal of Pathology, 1999, 155, 1767-1772.	3.8	280
2	Molecular pathology of endometrial hyperplasia and carcinoma. Human Pathology, 2001, 32, 569-577.	2.0	278
3	Endometrial carcinoma: pathology and genetics. Pathology, 2007, 39, 72-87.	0.6	172
4	Molecular genetic alterations in endometrioid carcinomas of the ovary: Similar frequency of beta-catenin abnormalities but lower rate of microsatellite instability and PTEN alterations than in uterine endometrioid carcinomas. Human Pathology, 2004, 35, 1360-1368.	2.0	167
5	Microsatellite instability in endometrial carcinomas: clinicopathologic correlations in a series of 42 cases. Human Pathology, 1998, 29, 1160-1164.	2.0	144
6	CTNNB1 mutations and β-catenin expression in endometrial carcinomas. Human Pathology, 2002, 33, 206-212.	2.0	130
7	Synchronous endometrioid carcinomas of the uterine corpus and ovary: alterations in the β-catenin (CTNNB1) pathway are associated with independent primary tumors and favorable prognosis. Human Pathology, 2005, 36, 605-619.	2.0	121
8	PIK3CA mutations in the kinase domain (exon 20) of uterine endometrial adenocarcinomas are associated with adverse prognostic parameters. Modern Pathology, 2008, 21, 131-139.	5.5	118
9	K-ras mutations in endometrial carcinomas with microsatellite instability. Journal of Pathology, 2001, 193, 193-199.	4.5	115
10	Concomitant PI3K–AKT and p53 alterations in endometrial carcinomas are associated with poor prognosis. Modern Pathology, 2009, 22, 522-529.	5.5	110
11	Microsatellite instability, MLH-1 promoter hypermethylation, and frameshift mutations at coding mononucleotide repeat microsatellites in ovarian tumors. Cancer, 2001, 92, 2829-2836.	4.1	103
12	Prognostic significance of FOXL2 mutation and mRNA expression in adult and juvenile granulosa cell tumors of the ovary. Modern Pathology, 2011, 24, 1360-1367.	5.5	89
13	Repression of E-cadherin by SNAIL, ZEB1, and TWIST in invasive ductal carcinomas of the breast: a cooperative effort?. Human Pathology, 2011, 42, 103-110.	2.0	76
14	Microsatellite instability and immunostaining for MSHâ€2 and MLHâ€1 in cutaneous and internal tumors from patients with the Muir–Torre syndrome. Journal of Cutaneous Pathology, 2002, 29, 415-420.	1.3	72
15	Promoter hypermethylation and reduced expression of RASSF1A are frequent molecular alterations of endometrial carcinoma. Modern Pathology, 2008, 21, 691-699.	5.5	71
16	Myometrial Invasion and Lymph Node Metastasis in Endometrioid Carcinomas: Tumor-associated Macrophages, Microvessel Density, and HIF1A Have a Crucial Role. American Journal of Surgical Pathology, 2010, 34, 1708-1714.	3.7	67
17	β- And γ-catenin expression in endometrial carcinoma. Relationship with clinicopathological features and microsatellite instability. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2001, 438, 464-469	2.8	52
18	Expression profiling of 22 genes involved in the PI3K–AKT pathway identifies two subgroups of high-grade endometrial carcinomas with different molecular alterations. Modern Pathology, 2010, 23, 694-702.	5.5	47

LLUIS CATASUS

#	Article	IF	CITATIONS
19	Frameshift mutations at coding mononucleotide repeat microsatellites in endometrial carcinoma with microsatellite instability. Cancer, 2000, 88, 2290-2297.	4.1	46
20	Clonality Analysis in Synchronous or Metachronous Tumors of the Female Genital Tract. International Journal of Gynecological Pathology, 2002, 21, 205-211.	1.4	43
21	Malignant Mullerian Mixed Tumor Arising From Ovarian Serous Carcinoma: A Clinicopathologic and Molecular Study of Two Cases. International Journal of Gynecological Pathology, 2002, 21, 268-272.	1.4	42
22	The STK11/LKB1 Peutz-Jegher gene is not involved in the pathogenesis of sporadic sex cord-stromal tumors, although loss of heterozygosity at 19p13.3 indicates other gene alteration in these tumors. Human Pathology, 2004, 35, 1101-1104.	2.0	41
23	Is loss of heterozygosity at 9q22.3 (PTCH gene) and 19p13.3 (STK11 gene) involved in the pathogenesis of ovarian stromal tumors?. Human Pathology, 2005, 36, 792-796.	2.0	38
24	Molecular pathology of atypical polypoid adenomyoma of the uterus. Human Pathology, 2003, 34, 784-788.	2.0	35
25	Exon 20 PIK3CA mutations decreases survival in aggressive (HER-2 positive) breast carcinomas. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2008, 453, 133-139.	2.8	34
26	The Sequence and Conformation of Human Pancreatic Procarboxypeptidase A2. Journal of Biological Chemistry, 1995, 270, 6651-6657.	3.4	31
27	Gene expression analysis identifies two groups of ovarian high-grade serous carcinomas with different prognosis. Modern Pathology, 2011, 24, 846-854.	5.5	26
28	Expression profile of heptahelical putative membrane progesterone receptors in epithelial ovarian tumors. Human Pathology, 2008, 39, 1026-1033.	2.0	25
29	Molecular genetics of endometrial carcinoma. Diagnostic Histopathology, 2009, 15, 554-563.	0.4	24
30	Stromal signatures in endometrioid endometrial carcinomas. Modern Pathology, 2014, 27, 631-639.	5.5	23
31	Cardiac chimerism in recipients of peripheral-blood and bone marrow stem cells. European Journal of Heart Failure, 2004, 6, 399-402.	7.1	22
32	The FOXO1-miR27 tandem regulates myometrial invasion in endometrioid endometrial adenocarcinoma. Human Pathology, 2014, 45, 942-951.	2.0	21
33	Low-density lipoprotein receptor–related protein 1 is associated with proliferation and invasiveness in Her-2/neu and triple-negative breast carcinomas. Human Pathology, 2011, 42, 1581-1588.	2.0	20
34	Characterisation and preliminary X-ray diffraction analysis of human pancreatic procarboxypeptidase A2. FEBS Letters, 1997, 420, 7-10.	2.8	19
35	Low-density lipoprotein receptor-related protein 1 (LRP-1) is associated with high-grade, advanced stage and p53 and p16 alterations in endometrial carcinomas. Histopathology, 2011, 59, 567-571.	2.9	17
36	Promoter hypermethylation contributes to <scp><scp>TIMP</scp></scp> <i>3</i> downâ€regulation in high stage endometrioid endometrial carcinomas. Histopathology, 2013, 62, 632-641.	2.9	17

LLUIS CATASUS

#	Article	IF	CITATIONS
37	Prognostic significance of the Fas-receptor/Fas-ligand system in cervical squamous cell carcinoma. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2008, 452, 65-74.	2.8	15
38	Comparative Analysis of the Sequences and Three-Dimensional Models of Human Procarboxypeptidases A1, A2 and B. Biological Chemistry, 1998, 379, 149-156.	2.5	13
39	Mutation Analysis of the SDHD Gene in Four Kindreds with Familial Paraganglioma. Diagnostic Molecular Pathology, 2005, 14, 109-114.	2.1	13
40	A novel germline mutation in exon 5 of the multiple endocrine neoplasia type 1 gene. Journal of Molecular Medicine, 1998, 76, 837-839.	3.9	11
41	Synchronous Bilateral Breast Carcinoma in a Patient with Cowden Syndrome: A Case Report with Morphologic, Immunohistochemical and Genetic Analysis. Breast Journal, 2010, 16, 77-81.	1.0	11
42	Molecular Pathology of Multiple Endocrine Neoplasia Type I. Diagnostic Molecular Pathology, 1999, 8, 195-204.	2.1	10
43	Genetic, Clinical, And Biochemical Analysis Of Unrelated Spanish Families With Multiple Endocrine Neoplasia Type I. Endocrine Practice, 2000, 6, 13-19.	2.1	4
44	Frameshift mutations at coding mononucleotide repeat microsatellites in endometrial carcinoma with microsatellite instability. Cancer, 2000, 88, 2290-2297.	4.1	2
45	K-ras mutations in endometrial carcinomas with microsatellite instability. , 2001, 193, 193.		1