

Akira Fujiyoshi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5167537/publications.pdf>

Version: 2024-02-01

143
papers

3,119
citations

172386

29
h-index

197736

49
g-index

147
all docs

147
docs citations

147
times ranked

4879
citing authors

#	ARTICLE	IF	CITATIONS
1	Caloric Restriction, the Traditional Okinawan Diet, and Healthy Aging. <i>Annals of the New York Academy of Sciences</i> , 2007, 1114, 434-455.	1.8	291
2	Patient-Important Outcomes in Registered Diabetes Trials. <i>JAMA - Journal of the American Medical Association</i> , 2008, 299, 2543.	3.8	194
3	Blood pressure categories and long-term risk of cardiovascular disease according to age group in Japanese men and women. <i>Hypertension Research</i> , 2012, 35, 947-953.	1.5	134
4	Factors Associated with Mortality of Myxedema Coma: Report of Eight Cases and Literature Survey. <i>Thyroid</i> , 1999, 9, 1167-1174.	2.4	125
5	Dietary sodium-to-potassium ratio as a risk factor for stroke, cardiovascular disease and all-cause mortality in Japan: the NIPPON DATA80 cohort study. <i>BMJ Open</i> , 2016, 6, e011632.	0.8	104
6	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population. <i>Journal of Hypertension</i> , 2017, 35, 401-408.	0.3	74
7	Long-term risk of BP values above normal for cardiovascular mortality. <i>Journal of Hypertension</i> , 2012, 30, 2299-2306.	0.3	70
8	Impact of Metabolic Syndrome on the Risk of Cardiovascular Disease Mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , 2014, 113, 84-89.	0.7	69
9	Aortic stiffness and calcification in men in a population-based international study. <i>Atherosclerosis</i> , 2012, 222, 473-477.	0.4	63
10	Influence of Smoking Combined with Another Risk Factor on the Risk of Mortality from Coronary Heart Disease and Stroke: Pooled Analysis of 10 Japanese Cohort Studies. <i>Cerebrovascular Diseases</i> , 2012, 33, 480-491.	0.8	62
11	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , 2014, 112, 916-924.	1.2	59
12	Epidemiology of Cardiovascular Risk Factors in Asian Countries. <i>Circulation Journal</i> , 2013, 77, 2851-2859.	0.7	56
13	Long-chain n-3 polyunsaturated fatty acids intake and cardiovascular disease mortality risk in Japanese: A 24-year follow-up of NIPPON DATA80. <i>Atherosclerosis</i> , 2014, 232, 384-389.	0.4	51
14	Increased Aortic Calcification Is Associated With Arterial Stiffness Progression in Multiethnic Middle-Aged Men. <i>Hypertension</i> , 2017, 69, 102-108.	1.3	51
15	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016, 23, 800-809.	0.9	48
16	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016, 246, 141-147.	0.4	48
17	Long chain n-3 polyunsaturated fatty acids and incidence rate of coronary artery calcification in Japanese men in Japan and white men in the USA: population based prospective cohort study. <i>Heart</i> , 2014, 100, 569-573.	1.2	47
18	Relationship of Insulin Resistance to Prevalence and Progression of Coronary Artery Calcification Beyond Metabolic Syndrome Components. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016, 36, 1703-1708.	1.1	44

#	ARTICLE	IF	CITATIONS
19	Coronary Artery Calcium and Risk of Dementia in MESA (Multi-Ethnic Study of Atherosclerosis). <i>Circulation: Cardiovascular Imaging</i> , 2017, 10, .	1.3	44
20	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2013, 20, 755-766.	0.9	43
21	Validity of Death Certificate and Hospital Discharge ICD Codes for Dementia Diagnosis. <i>Alzheimer Disease and Associated Disorders</i> , 2017, 31, 168-172.	0.6	40
22	Comparison of HOMA-IR, HOMA- β % and disposition index between US white men and Japanese men in Japan: the ERA JUMP study. <i>Diabetologia</i> , 2015, 58, 265-271.	2.9	39
23	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980â€“2010. <i>Hypertension Research</i> , 2015, 38, 790-795.	1.5	39
24	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , 2016, 5, .	1.6	39
25	Coronary Artery Calcium and Carotid Artery Intima Media Thickness and Plaque: Clinical Use in Need of Clarification. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 227-239.	0.9	38
26	Cross-Sectional Comparison of Coronary Artery Calcium Scores Between Caucasian Men in the United States and Japanese Men in Japan: The Multi-Ethnic Study of Atherosclerosis and the Shiga Epidemiological Study of Subclinical Atherosclerosis. <i>American Journal of Epidemiology</i> , 2014, 180, 590-598.	1.6	36
27	Association Between J-Point Elevation and Death From Coronary Artery Disease. <i>Circulation Journal</i> , 2013, 77, 1260-1266.	0.7	35
28	Serum levels of marine-derived n-3 fatty acids in Icelanders, Japanese, Koreans, and Americansâ€”A descriptive epidemiologic study. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2012, 87, 11-16.	1.0	31
29	Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. <i>British Journal of Nutrition</i> , 2017, 117, 260-266.	1.2	31
30	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Preventive Medicine Reports</i> , 2016, 4, 225-232.	0.8	30
31	Relationship of serum irisin levels to prevalence and progression of coronary artery calcification: A prospective, population-based study. <i>International Journal of Cardiology</i> , 2018, 267, 177-182.	0.8	30
32	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , 2016, 220, 262-267.	0.8	29
33	Home blood pressure variability and subclinical atherosclerosis in multiple vascular beds. <i>Journal of Hypertension</i> , 2018, 36, 2193-2203.	0.3	28
34	Mendelian randomization analysis in three Japanese populations supports a causal role of alcohol consumption in lowering low-density lipid cholesterol levels and particle numbers. <i>Atherosclerosis</i> , 2016, 254, 242-248.	0.4	27
35	Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. <i>Atherosclerosis</i> , 2018, 273, 145-152.	0.4	27
36	Coronary Artery Calcification by Computed Tomography in Epidemiologic Research and Cardiovascular Disease Prevention. <i>Journal of Epidemiology</i> , 2012, 22, 188-198.	1.1	26

#	ARTICLE	IF	CITATIONS
37	Influence of cigarette smoking on coronary artery and aortic calcium among random samples from populations of middle-aged Japanese and Korean men. <i>Journal of Epidemiology and Community Health</i> , 2013, 67, 119-124.	2.0	24
38	Brachial-ankle pulse wave velocity is associated with coronary calcification among 1131 healthy middle-aged men. <i>International Journal of Cardiology</i> , 2015, 189, 67-72.	0.8	24
39	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , 2014, 236, 237-243.	0.4	22
40	Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. <i>European Journal of Clinical Nutrition</i> , 2019, 73, 783-792.	1.3	22
41	LOX-1 ligands containing apolipoprotein B and carotid intima-media thickness in middle-aged community-dwelling US Caucasian and Japanese men. <i>Atherosclerosis</i> , 2013, 229, 240-245.	0.4	21
42	A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and US Whites. <i>European Heart Journal Cardiovascular Imaging</i> , 2013, 14, 921-927.	0.5	21
43	Association between Pulse Wave Velocity and Coronary Artery Calcification in Japanese men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015, 22, 1266-1277.	0.9	21
44	Association of blood pressure with estimates of 24-h urinary sodium and potassium excretion from repeated single-spot urine samples. <i>Hypertension Research</i> , 2019, 42, 411-418.	1.5	21
45	Fatty Acids Intakes and Serum Lipid Profiles: NIPPON DATA90 and the National Nutrition Monitoring. <i>Journal of Epidemiology</i> , 2010, 20, S544-S548.	1.1	20
46	Associations between Inflammatory Markers and Subclinical Atherosclerosis in Middle-aged White, Japanese-American and Japanese Men: The ERA-JUMP Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015, 22, 590-598.	0.9	20
47	Relationship Between Step Counts and Cerebral Small Vessel Disease in Japanese Men. <i>Stroke</i> , 2020, 51, 3584-3591.	1.0	19
48	The prevalence of aortic calcification in Japanese compared to white and Japanese-American middle-aged men is confounded by the amount of cigarette smoking. <i>International Journal of Cardiology</i> , 2013, 167, 134-139.	0.8	18
49	A Cross-Sectional Association Between Bone Mineral Density and Parathyroid Hormone and Other Biomarkers in Community-Dwelling Young Adults: The CARDIA Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 4038-4046.	1.8	18
50	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015, 239, 444-450.	0.4	18
51	Intracranial Artery Stenosis and Its Association With Conventional Risk Factors in a General Population of Japanese Men. <i>Stroke</i> , 2019, 50, 2967-2969.	1.0	18
52	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 606-620.	0.9	17
53	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 198-206.	0.9	17
54	Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. <i>European Journal of Clinical Nutrition</i> , 2020, 74, 67-76.	1.3	17

#	ARTICLE	IF	CITATIONS
55	Association of Total Energy Intake with 29-Year Mortality in the Japanese: NIPPON DATA80. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016, 23, 339-354.	0.9	16
56	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. <i>Environmental Pollution</i> , 2016, 213, 460-467.	3.7	16
57	Population Attributable Fraction of Smoking and Metabolic Syndrome on Cardiovascular Disease Mortality in Japan: a 15-Year Follow Up of NIPPON DATA90. <i>BMC Public Health</i> , 2010, 10, 306.	1.2	15
58	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S29-S34.	1.1	15
59	Differences Between Coronary Artery Calcification and Aortic Artery Calcification in Relation to Cardiovascular Disease Risk Factors in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 452-464.	0.9	13
60	Long-term outcomes associated with prolonged PR interval in the general Japanese population. <i>International Journal of Cardiology</i> , 2015, 184, 291-293.	0.8	12
61	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1195-1202.e1.	0.6	12
62	The relationship between serum levels of LOX-1 ligand containing ApoAI as a novel marker of dysfunctional HDL and coronary artery calcification in middle-aged Japanese men. <i>Atherosclerosis</i> , 2020, 313, 20-25.	0.4	12
63	The Association Between Coronary Artery Calcification and Subclinical Cerebrovascular Diseases in Men: An Observational Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 995-1009.	0.9	12
64	Relationship of moderate metabolic risk factor clustering to cardiovascular disease mortality in non-lean Japanese: A 15-year follow-up of NIPPON DATA90. <i>Atherosclerosis</i> , 2011, 215, 209-213.	0.4	11
65	Metabolic syndrome and its components are underdiagnosed in cardiology clinics. <i>Journal of Evaluation in Clinical Practice</i> , 2011, 17, 78-83.	0.9	11
66	Association between Alanine Aminotransferase and Intracerebral Hemorrhage in East Asian Populations. <i>Neuroepidemiology</i> , 2013, 41, 131-138.	1.1	11
67	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: A 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , 2014, 64, 218-224.	0.8	11
68	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia. <i>Journal of Clinical Lipidology</i> , 2016, 10, 172-180.e1.	0.6	11
69	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. <i>Circulation Journal</i> , 2018, 82, 2542-2548.	0.7	11
70	Cardiovascular Risk Assessment Chart by Dietary Factors in Japan. <i>NIPPON DATA80</i> . <i>Circulation Journal</i> , 2019, 83, 1254-1260.	0.7	11
71	Association of Total Marine Fatty Acids, Eicosapentaenoic and Docosahexaenoic Acids, With Aortic Stiffness in Koreans, Whites, and Japanese Americans. <i>American Journal of Hypertension</i> , 2013, 26, 1321-1327.	1.0	10
72	Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , 2018, 37, 182-188.	2.3	10

#	ARTICLE	IF	CITATIONS
73	Reduced Lung Function and Cerebral Small Vessel Disease in Japanese Men: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 1009-1021.	0.9	10
74	Coronary Artery Calcium Progression Among the US and Japanese Men. <i>Circulation: Cardiovascular Imaging</i> , 2019, 12, e008104.	1.3	10
75	Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2020, 30, 244-252.	1.1	10
76	Lipoprotein-Associated Phospholipase A2 Regulates Macrophage Apoptosis via the Akt and Caspase-7 Pathways. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014, 21, 839-853.	0.9	9
77	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. <i>Atherosclerosis</i> , 2016, 252, 116-121.	0.4	9
78	The role of initial and longitudinal change in blood pressure on progression of arterial stiffness among multiethnic middle-aged men. <i>Journal of Hypertension</i> , 2017, 35, 111-117.	0.3	9
79	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. <i>Metabolic Syndrome and Related Disorders</i> , 2018, 16, 166-173.	0.5	9
80	Comparison of carotid plaque burden among healthy middle-aged men living in the US, Japan, and South Korea. <i>International Journal of Cardiology</i> , 2018, 266, 245-249.	0.8	9
81	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019, 29, 133-138.	1.1	9
82	Elevated Fasting Blood Glucose Levels Are Associated With Lower Cognitive Function, With a Threshold in Non-Diabetic Individuals: A Population-Based Study. <i>Journal of Epidemiology</i> , 2020, 30, 121-127.	1.1	9
83	Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Epidemiology</i> , 2019, 29, 205-212.	1.1	9
84	Relationship between Kidney Function and Subclinical Atherosclerosis Progression Evaluated by Coronary Artery Calcification. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 1359-1371.	0.9	9
85	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. <i>Hypertension Research</i> , 2016, 39, 670-679.	1.5	8
86	Smoking habits and parathyroid hormone concentrations in young adults: The CARDIA study. <i>Bone Reports</i> , 2016, 5, 104-109.	0.2	8
87	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. <i>European Journal of Preventive Cardiology</i> , 2017, 24, 505-513.	0.8	8
88	Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , 2018, 18, 799-805.	0.7	8
89	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 417-423.	1.2	8
90	The effects of renin-angiotensin system inhibitors on mortality, cardiovascular events, and renal events in hypertensive patients with diabetes: a systematic review and meta-analysis of randomized controlled trials. <i>Hypertension Research</i> , 2019, 42, 669-680.	1.5	8

#	ARTICLE	IF	CITATIONS
91	Comparison of the National Nutritional Survey in Japan Estimated Individual-Based Nutritional Data and NIPPON DATA80 Food Frequency Questionnaires. <i>Journal of Epidemiology</i> , 2010, 20, S582-S586.	1.1	7
92	Comparability in coronary artery calcium scores on CT scan between two community-based cohort studies. <i>International Journal of Cardiology</i> , 2011, 149, 244-245.	0.8	7
93	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , 2013, 99, 1024-1029.	1.2	7
94	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men—A 20-Year Follow-up of NIPPON DATA90. <i>Circulation Journal</i> , 2019, 83, 1506-1513.	0.7	7
95	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , 2019, 37, 1676-1681.	0.3	7
96	Alcohol consumption and cognitive function in elderly Japanese men. <i>Alcohol</i> , 2020, 85, 145-152.	0.8	7
97	Smoking habits and progression of coronary and aortic artery calcification: A 5-year follow-up of community-dwelling Japanese men. <i>International Journal of Cardiology</i> , 2020, 314, 89-94.	0.8	7
98	Relationship of Higher-level Functional Capacity With Long-term Mortality in Japanese Older People: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2023, 33, 136-141.	1.1	7
99	A Comparison of Segment-Specific and Composite Measures of Carotid Intima-Media Thickness and their Relationships with Coronary Calcium. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 282-295.	0.9	7
100	Significant inverse associations of serum n-6 fatty acids with plasma plasminogen activator inhibitor-1. <i>British Journal of Nutrition</i> , 2012, 107, 567-572.	1.2	6
101	Association of alcohol consumption and aortic calcification in healthy men aged 40–49 years for the ERA JUMP Study. <i>Atherosclerosis</i> , 2018, 268, 84-91.	0.4	6
102	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. <i>Hypertension Research</i> , 2021, 44, 80-87.	1.5	6
103	Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. <i>Alcohol</i> , 2021, 90, 57-65.	0.8	6
104	̢-Glutamyltransferase and mortality risk from heart disease and stroke in Japanese men and women: NIPPON DATA90. <i>CVD Prevention and Control</i> , 2010, 5, 27.	0.7	5
105	Self-reported Sleep Duration and Subclinical Atherosclerosis in a General Population of Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 186-198.	0.9	5
106	Hyperglycemia, duration of diabetes, and intracranial atherosclerotic stenosis by magnetic resonance angiography: The ARIC-NCS study. <i>Journal of Diabetes and Its Complications</i> , 2020, 34, 107605.	1.2	5
107	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 252, 253-262.	0.5	5
108	Risk factors for heart failure and coronary heart disease mortality over 24-year follow-up period in Japan: NIPPON DATA80. <i>CVD Prevention and Control</i> , 2010, 5, 97.	0.7	4

#	ARTICLE	IF	CITATIONS
109	Association of Coronary Artery Calcification with Estimated Coronary Heart Disease Risk from Prediction Models in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 477-489.	0.9	4
110	Passive Smoking at Home by Socioeconomic Factors in a Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S40-S45.	1.1	4
111	Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical) <i>Tj ETQq1 1 0.784314 rgBTi/Overlock 10 Tf 5016</i>		
112	Coronary Artery Calcium Assessed Years Before Was Positively Associated With Subtle White Matter Injury of the Brain in Asymptomatic Middle-Aged Men: The Framingham Heart Study. <i>Circulation: Cardiovascular Imaging</i> , 2021, 14, e011753.	1.3	4
113	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. <i>PLoS ONE</i> , 2016, 11, e0157563.	1.1	4
114	Serum long-chain n-3 polyunsaturated fatty acids and aortic calcification in middle-aged men: The population-based cross-sectional ERA-JUMP study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019, 29, 837-846.	1.1	3
115	Anthropometric Obesity Indices were Stronger than CT-Based Indices in Associations with Carotid Intima-Media Thickness in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 1102-1114.	0.9	3
116	Cross-sectional association of bone mineral density with coronary artery calcification in an international multi-ethnic population-based cohort of men aged 40-49: ERA JUMP study. <i>IJC Heart and Vasculature</i> , 2020, 30, 100618.	0.6	3
117	Seven-Day Pedometer-Assessed Step Counts and Brain Volume: A Population-Based Observational Study. <i>Journal of Physical Activity and Health</i> , 2021, 18, 157-164.	1.0	3
118	Association of self-measured home, ambulatory, and strictly measured office blood pressure and their variability with intracranial arterial stenosis. <i>Journal of Hypertension</i> , 2021, 39, 2030-2039.	0.3	3
119	Ventricular Premature Complexes and Their Associated Factors in a General Population of Japanese Men. <i>American Journal of Cardiology</i> , 2022, 169, 51-56.	0.7	3
120	Association of equol producing status with aortic calcification in middle-aged Japanese men: The ERA JUMP study. <i>International Journal of Cardiology</i> , 2022, 352, 158-164.	0.8	3
121	Progression of coronary artery calcium in Japanese American men and white men in the ERA JUMP study. <i>International Journal of Cardiology</i> , 2017, 228, 672-676.	0.8	2
122	Data on alcohol consumption and coronary artery calcification among asymptomatic middle-aged men for the ERA-JUMP study. <i>Data in Brief</i> , 2018, 17, 1091-1098.	0.5	2
123	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S35-S39.	1.1	2
124	Eczema and Risk of Cardiovascular Disease: Heightened Awareness Needed. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 760-761.	0.9	2
125	Relationship of Four Blood Pressure Indexes to Subclinical Cerebrovascular Diseases Assessed by Brain MRI in General Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 174-187.	0.9	2
126	Lipoprotein Particle Profiles Compared With Standard Lipids in the Association With Subclinical Aortic Valve Calcification in Apparently Healthy Japanese Men. <i>Circulation Journal</i> , 2021, 85, 1076-1082.	0.7	2

#	ARTICLE	IF	CITATIONS
127	Predictors of lower limb fractures in general Japanese: NIPPON DATA90. PLoS ONE, 2022, 17, e0261716.	1.1	2
128	Premature Atrial Contractions and Their Determinants in a General Population of Japanese Men. Circulation Journal, 2022, 86, 1298-1306.	0.7	2
129	Association of ambulatory blood pressure with aortic valve and coronary artery calcification. Journal of Hypertension, 2022, 40, 1344-1351.	0.3	2
130	Cardiorespiratory fitness, a promising modality for treatment and risk prediction. European Journal of Preventive Cardiology, 2018, 25, 306-308.	0.8	1
131	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). Journal of Atherosclerosis and Thrombosis. 2019, 26, 746-746.	0.9	1
132	Lipoprotein particles and coronary artery calcium in middle-aged US-White and Japanese men. Open Heart, 2019, 6, e001119.	0.9	1
133	Factors Associated with Lower Cognitive Performance Scores Among Older Japanese Men in Hawaii and Japan. Journal of Alzheimer's Disease, 2021, 81, 403-412.	1.2	1
134	Use of heated tobacco products and pulmonary function in a real-world: more studies needed to answer many important questions. Journal of Epidemiology, 2021, , .	1.1	1
135	Two Cases of Myxedema Coma Successfully Treated with Low-Dose Thyroid Hormone Replacement.. Nihon Kyukyu Igakukai Zasshi, 2000, 11, 22-26.	0.0	1
136	Relation of Progression of Coronary Artery Calcium to Dementia (from the Multi-Ethnic Study of) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50 3	0.7	1
137	Corrigendum to "Gamma-Glutamyltransferase and mortality risk from heart disease and stroke in Japanese men and women: NIPPON DATA 90". CVD Prevention and Control, 2011, 6, 63.	0.7	0
138	J-005%AN INCREASED SERUM ALANINE AMINOTRANSFERASE IS ASSOCIATED WITH AN INCREASED MORTALITY FROM INTRACEREBRAL HEMORRHAGE IN THE EAST ASIAN POPULATION. Journal of Hypertension, 2011, 29, e25.	0.3	0
139	ICâ€Pâ€112: OBJECTIVELY MEASURED PHYSICAL ACTIVITY AND BRAIN VOLUME IN JAPANESE ADULT MEN: FINDINGS FROM THE SHIGA EPIDEMIOLOGICAL STUDY OF SUBCLINICAL ATHEROSCLEROSIS. Alzheimer's and Dementia, 2019, 15, P96.	0.4	0
140	Is Measuring Risk Marker Progression Useful for Cardiovascular Disease Prediction?. Cerebrovascular Diseases, 2021, 50, 1-4.	0.8	0
141	How Is Socioeconomic Status Associated With the Incidence of Heart Failure?. Circulation Journal, 2021, 85, 1553-1554.	0.7	0
142	913Factors of premature atrial contractions among general Japanese men. International Journal of Epidemiology, 2021, 50, .	0.9	0
143	Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men. Cerebrovascular Diseases, 2022, 51, 774-780.	0.8	0