

Akira Fujiyoshi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5167537/akira-fujiyoshi-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

130
papers

2,175
citations

26
h-index

41
g-index

147
ext. papers

2,681
ext. citations

3.6
avg, IF

4.24
L-index

#	Paper	IF	Citations
130	Predictors of lower limb fractures in general Japanese: NIPPON DATA90.. <i>PLoS ONE</i> , 2022 , 17, e0261716	3.7	0
129	Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men.. <i>Cerebrovascular Diseases</i> , 2022 , 1-7	3.2	
128	Factors Associated with Lower Cognitive Performance Scores Among Older Japanese Men in Hawaii and Japan. <i>Journal of Alzheimer's Disease</i> , 2021 , 81, 403-412	4.3	
127	Lipoprotein Particle Profiles Compared With Standard Lipids in the Association With Subclinical Aortic Valve Calcification in Apparently Healthy Japanese Men. <i>Circulation Journal</i> , 2021 , 85, 1076-1082	2.9	1
126	Association of self-measured home, ambulatory, and strictly measured office blood pressure and their variability with intracranial arterial stenosis. <i>Journal of Hypertension</i> , 2021 , 39, 2030-2039	1.9	0
125	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. <i>Hypertension Research</i> , 2021 , 44, 80-87	4.7	2
124	Relationship of higher-level functional capacity with long-term mortality in Japanese older people: NIPPON DATA90. <i>Journal of Epidemiology</i> , 2021 ,	3.4	2
123	Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. <i>Alcohol</i> , 2021 , 90, 57-65	2.7	2
122	Seven-Day Pedometer-Assessed Step Counts and Brain Volume: A Population-Based Observational Study. <i>Journal of Physical Activity and Health</i> , 2021 , 18, 157-164	2.5	0
121	Coronary Artery Calcium Assessed Years Before Was Positively Associated With Subtle White Matter Injury of the Brain in Asymptomatic Middle-Aged Men: The Framingham Heart Study. <i>Circulation: Cardiovascular Imaging</i> , 2021 , 14, e011753	3.9	0
120	Is Measuring Risk Marker Progression Useful for Cardiovascular Disease Prediction?. <i>Cerebrovascular Diseases</i> , 2021 , 50, 752-755	3.2	
119	How Is Socioeconomic Status Associated With the Incidence of Heart Failure?. <i>Circulation Journal</i> , 2021 , 85, 1553-1554	2.9	
118	Relationship Between Step Counts and Cerebral Small Vessel Disease in Japanese Men. <i>Stroke</i> , 2020 , 51, 3584-3591	6.7	5
117	Smoking habits and progression of coronary and aortic artery calcification: A 5-year follow-up of community-dwelling Japanese men. <i>International Journal of Cardiology</i> , 2020 , 314, 89-94	3.2	3
116	The Association Between Coronary Artery Calcification and Subclinical Cerebrovascular Diseases in Men: An Observational Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 995-1009	4	5
115	Hyperglycemia, duration of diabetes, and intracranial atherosclerotic stenosis by magnetic resonance angiography: The ARIC-NCS study. <i>Journal of Diabetes and Its Complications</i> , 2020 , 34, 107605	3.2	2
114	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 252, 253-262	2.4	0

113	Alcohol consumption and cognitive function in elderly Japanese men. <i>Alcohol</i> , 2020 , 85, 145-152	2.7	3
112	Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical Atherosclerosis). <i>Diabetes Research and Clinical Practice</i> , 2020 , 160, 108002	7.4	2
111	The relationship between serum levels of LOX-1 ligand containing ApoAI as a novel marker of dysfunctional HDL and coronary artery calcification in middle-aged Japanese men. <i>Atherosclerosis</i> , 2020 , 313, 20-25	3.1	1
110	Cross-sectional association of bone mineral density with coronary artery calcification in an international multi-ethnic population-based cohort of men aged 40-49: ERA JUMP study. <i>IJC Heart and Vasculature</i> , 2020 , 30, 100618	2.4	0
109	Proteinuria and Reduced Estimated Glomerular Filtration Rate are Independently Associated With Lower Cognitive Abilities in Apparently Healthy Community-Dwelling Elderly Men in Japan: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2020 , 30, 244-252	3.4	3
108	Elevated Fasting Blood Glucose Levels Are Associated With Lower Cognitive Function, With a Threshold in Non-Diabetic Individuals: A Population-Based Study. <i>Journal of Epidemiology</i> , 2020 , 30, 121-127	3.4	1
107	Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. <i>European Journal of Clinical Nutrition</i> , 2020 , 74, 67-76	5.2	4
106	Cardiovascular Risk Assessment Chart by Dietary Factors in Japan - NIPPON DATA80. <i>Circulation Journal</i> , 2019 , 83, 1254-1260	2.9	5
105	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men - A 20-Year Follow-up of NIPPON DATA90. <i>Circulation Journal</i> , 2019 , 83, 1506-1513	2.9	4
104	Anthropometric Obesity Indices were Stronger than CT-Based Indices in Associations with Carotid Intima-Media Thickness in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 1102-1114 ⁴		2
103	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , 2019 , 33, 417-423	3.2	6
102	Differences Between Coronary Artery Calcification and Aortic Artery Calcification in Relation to Cardiovascular Disease Risk Factors in Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 452-464	4	3
101	The effects of renin-angiotensin system inhibitors on mortality, cardiovascular events, and renal events in hypertensive patients with diabetes: a systematic review and meta-analysis of randomized controlled trials. <i>Hypertension Research</i> , 2019 , 42, 669-680	4.7	4
100	Coronary Artery Calcium Progression Among the US and Japanese Men. <i>Circulation: Cardiovascular Imaging</i> , 2019 , 12, e008104	3.9	3
99	Association of blood pressure with estimates of 24-h urinary sodium and potassium excretion from repeated single-spot urine samples. <i>Hypertension Research</i> , 2019 , 42, 411-418	4.7	15
98	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019 , 29, 133-138	3.4	3
97	Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. <i>European Journal of Clinical Nutrition</i> , 2019 , 73, 783-792	5.2	13
96	Intracranial Artery Stenosis and Its Association With Conventional Risk Factors in a General Population of Japanese Men. <i>Stroke</i> , 2019 , 50, 2967-2969	6.7	13

95	Serum long-chain n-3 polyunsaturated fatty acids and aortic calcification in middle-aged men: The population-based cross-sectional ERA-JUMP study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019 , 29, 837-846	4.5	3
94	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA).	4	
93	Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Epidemiology</i> , 2019 , 29, 205-212	3.4	6
92	Lipoprotein particles and coronary artery calcium in middle-aged US-White and Japanese men. <i>Open Heart</i> , 2019 , 6, e001119	3	
91	IC-P-112: OBJECTIVELY MEASURED PHYSICAL ACTIVITY AND BRAIN VOLUME IN JAPANESE ADULT MEN: FINDINGS FROM THE SHIGA EPIDEMIOLOGICAL STUDY OF SUBCLINICAL ATHEROSCLEROSIS 2019 , 15, P96-P96		
90	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , 2019 , 37, 1676-1681	1.9	4
89	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 198-206	4	12
88	Data on alcohol consumption and coronary artery calcification among asymptomatic middle-aged men for the ERA-JUMP study. <i>Data in Brief</i> , 2018 , 17, 1091-1098	1.2	2
87	Association of Coronary Artery Calcification with Estimated Coronary Heart Disease Risk from Prediction Models in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 477-489	4	3
86	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. <i>Metabolic Syndrome and Related Disorders</i> , 2018 , 16, 166-173	2.6	5
85	Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 799-805	2.9	5
84	Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. <i>Atherosclerosis</i> , 2018 , 273, 145-152	3.1	12
83	Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , 2018 , 37, 182-188	5.9	7
82	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. <i>Circulation Journal</i> , 2018 , 82, 2542-2548	2.9	4
81	Comparison of carotid plaque burden among healthy middle-aged men living in the US, Japan, and South Korea. <i>International Journal of Cardiology</i> , 2018 , 266, 245-249	3.2	5
80	Relationship of serum irisin levels to prevalence and progression of coronary artery calcification: A prospective, population-based study. <i>International Journal of Cardiology</i> , 2018 , 267, 177-182	3.2	15
79	Reduced Lung Function and Cerebral Small Vessel Disease in Japanese Men: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 1009-1021	4	6
78	Association of alcohol consumption and aortic calcification in healthy men aged 40-49 years for the ERA JUMP Study. <i>Atherosclerosis</i> , 2018 , 268, 84-91	3.1	4

77	Home blood pressure variability and subclinical atherosclerosis in multiple vascular beds: a population-based study. <i>Journal of Hypertension</i> , 2018 , 36, 2193-2203	1.9	13
76	Self-reported Sleep Duration and Subclinical Atherosclerosis in a General Population of Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 186-198	4	5
75	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S29-S34	3.4	11
74	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S35-S39	3.4	2
73	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , 2018 , 25, 606-620	4	10
72	Passive Smoking at Home by Socioeconomic Factors in a Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018 , 28 Suppl 3, S40-S45	3.4	3
71	Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. <i>British Journal of Nutrition</i> , 2017 , 117, 260-266	3.6	25
70	Coronary Artery Calcium and Risk of Dementia in MESA (Multi-Ethnic Study of Atherosclerosis). <i>Circulation: Cardiovascular Imaging</i> , 2017 , 10,	3.9	25
69	The role of initial and longitudinal change in blood pressure on progression of arterial stiffness among multiethnic middle-aged men. <i>Journal of Hypertension</i> , 2017 , 35, 111-117	1.9	8
68	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population: NIPPON DATA2010. <i>Journal of Hypertension</i> , 2017 , 35, 401-408	1.9	59
67	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. <i>European Journal of Preventive Cardiology</i> , 2017 , 24, 505-513	3.9	5
66	Coronary Artery Calcium and Carotid Artery Intima Media Thickness and Plaque: Clinical Use in Need of Clarification. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017 , 24, 227-239	4	26
65	Validity of Death Certificate and Hospital Discharge ICD Codes for Dementia Diagnosis: The Multi-Ethnic Study of Atherosclerosis. <i>Alzheimer Disease and Associated Disorders</i> , 2017 , 31, 168-172	2.5	22
64	Increased Aortic Calcification Is Associated With Arterial Stiffness Progression in Multiethnic Middle-Aged Men. <i>Hypertension</i> , 2017 , 69, 102-108	8.5	37
63	Progression of coronary artery calcium in Japanese American men and white men in the ERA JUMP study. <i>International Journal of Cardiology</i> , 2017 , 228, 672-676	3.2	1
62	Dietary sodium-to-potassium ratio as a risk factor for stroke, cardiovascular disease and all-cause mortality in Japan: the NIPPON DATA80 cohort study. <i>BMJ Open</i> , 2016 , 6, e011632	3	53
61	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , 2016 , 220, 262-7	3.2	16
60	Smoking habits and parathyroid hormone concentrations in young adults: The CARDIA study. <i>Bone Reports</i> , 2016 , 5, 104-109	2.6	7

59	Relationship of Insulin Resistance to Prevalence and Progression of Coronary Artery Calcification Beyond Metabolic Syndrome Components: Shiga Epidemiological Study of Subclinical Atherosclerosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016 , 36, 1703-8	9.4	26
58	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016 , 246, 141-7	3.1	36
57	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia LOX-1 ligand and IMT in Japanese. <i>Journal of Clinical Lipidology</i> , 2016 , 10, 172-80.e1	4.9	10
56	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016 , 10, 1195-1202.e1	4.9	9
55	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. <i>PLoS ONE</i> , 2016 , 11, e0157563 ³	3.7	3
54	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016 , 23, 800-9	4	33
53	Association of Total Energy Intake with 29-Year Mortality in the Japanese: NIPPON DATA80. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016 , 23, 339-54	4	12
52	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. <i>Environmental Pollution</i> , 2016 , 213, 460-467	9.3	13
51	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. <i>Hypertension Research</i> , 2016 , 39, 670-9	4.7	8
50	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , 2016 , 5,	6	26
49	Mendelian randomization analysis in three Japanese populations supports a causal role of alcohol consumption in lowering low-density lipid cholesterol levels and particle numbers. <i>Atherosclerosis</i> , 2016 , 254, 242-248	3.1	18
48	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. <i>Atherosclerosis</i> , 2016 , 252, 116-121	3.1	6
47	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Preventive Medicine Reports</i> , 2016 , 4, 225-32	2.6	26
46	Comparison of HOMA-IR, HOMA-% and disposition index between US white men and Japanese men in Japan: the ERA JUMP study. <i>Diabetologia</i> , 2015 , 58, 265-71	10.3	28
45	Long-term outcomes associated with prolonged PR interval in the general Japanese population. <i>International Journal of Cardiology</i> , 2015 , 184, 291-293	3.2	9
44	Brachial-ankle pulse wave velocity is associated with coronary calcification among 1131 healthy middle-aged men. <i>International Journal of Cardiology</i> , 2015 , 189, 67-72	3.2	20
43	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980-2010. <i>Hypertension Research</i> , 2015 , 38, 790-5	4.7	32
42	Associations between Inflammatory Markers and Subclinical Atherosclerosis in Middle-aged White, Japanese-American and Japanese Men: The ERA-JUMP Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 590-8	4	17

41	Association between Pulse Wave Velocity and Coronary Artery Calcification in Japanese men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 1266-77	4	17
40	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015 , 239, 444-50	3.1	16
39	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , 2014 , 236, 237-43	3.1	17
38	Long-chain n-3 polyunsaturated fatty acids intake and cardiovascular disease mortality risk in Japanese: a 24-year follow-up of NIPPON DATA80. <i>Atherosclerosis</i> , 2014 , 232, 384-9	3.1	42
37	Lipoprotein-associated phospholipase A2 regulates macrophage apoptosis via the Akt and caspase-7 pathways. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014 , 21, 839-53	4	3
36	Cross-sectional comparison of coronary artery calcium scores between Caucasian men in the United States and Japanese men in Japan: the multi-ethnic study of atherosclerosis and the Shiga epidemiological study of subclinical atherosclerosis. <i>American Journal of Epidemiology</i> , 2014 , 180, 590-8	3.8	29
35	Long chain n-3 polyunsaturated fatty acids and incidence rate of coronary artery calcification in Japanese men in Japan and white men in the USA: population based prospective cohort study. <i>Heart</i> , 2014 , 100, 569-73	5.1	39
34	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , 2014 , 112, 916-24	3.6	35
33	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: a 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , 2014 , 64, 218-24	3	9
32	Impact of metabolic syndrome on the risk of cardiovascular disease mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , 2014 , 113, 84-9	3	60
31	LOX-1 ligands containing apolipoprotein B and carotid intima-media thickness in middle-aged community-dwelling US Caucasian and Japanese men. <i>Atherosclerosis</i> , 2013 , 229, 240-5	3.1	17
30	The prevalence of aortic calcification in Japanese compared to white and Japanese-American middle-aged men is confounded by the amount of cigarette smoking. <i>International Journal of Cardiology</i> , 2013 , 167, 134-9	3.2	13
29	A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and U.S. whites. <i>European Heart Journal Cardiovascular Imaging</i> , 2013 , 14, 921-7	4.1	16
28	A cross-sectional association between bone mineral density and parathyroid hormone and other biomarkers in community-dwelling young adults: the CARDIA study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 4038-46	5.6	13
27	Influence of cigarette smoking on coronary artery and aortic calcium among random samples from populations of middle-aged Japanese and Korean men. <i>Journal of Epidemiology and Community Health</i> , 2013 , 67, 119-24	5.1	18
26	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , 2013 , 99, 1024-9	5.1	6
25	Association of total marine fatty acids, eicosapentaenoic and docosahexaenoic acids, with aortic stiffness in Koreans, whites, and Japanese Americans. <i>American Journal of Hypertension</i> , 2013 , 26, 1321-7	7.3	8
24	Association between J-point elevation and death from coronary artery disease--15-year follow up of the NIPPON DATA90. <i>Circulation Journal</i> , 2013 , 77, 1260-6	2.9	28

23	Epidemiology of cardiovascular risk factors in Asian countries. <i>Circulation Journal</i> , 2013 , 77, 2851-9	2.9	44
22	Association between alanine aminotransferase and intracerebral hemorrhage in East Asian populations. <i>Neuroepidemiology</i> , 2013 , 41, 131-8	5.4	8
21	Carotid intima-media thickness and plaque in apparently healthy Japanese individuals with an estimated 10-year absolute risk of CAD death according to the Japan Atherosclerosis Society (JAS) guidelines 2012: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 755-66	4	35
20	Coronary artery calcification by computed tomography in epidemiologic research and cardiovascular disease prevention. <i>Journal of Epidemiology</i> , 2012 , 22, 188-98	3.4	23
19	Aortic stiffness and calcification in men in a population-based international study. <i>Atherosclerosis</i> , 2012 , 222, 473-7	3.1	52
18	Serum levels of marine-derived n-3 fatty acids in Icelanders, Japanese, Koreans, and Americans—a descriptive epidemiologic study. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2012 , 87, 11-6	2.8	26
17	Significant inverse associations of serum n-6 fatty acids with plasma plasminogen activator inhibitor-1. <i>British Journal of Nutrition</i> , 2012 , 107, 567-72	3.6	6
16	Influence of smoking combined with another risk factor on the risk of mortality from coronary heart disease and stroke: pooled analysis of 10 Japanese cohort studies. <i>Cerebrovascular Diseases</i> , 2012 , 33, 480-91	3.2	50
15	Blood pressure categories and long-term risk of cardiovascular disease according to age group in Japanese men and women. <i>Hypertension Research</i> , 2012 , 35, 947-53	4.7	87
14	Long-term risk of BP values above normal for cardiovascular mortality: a 24-year observation of Japanese aged 30 to 92 years. <i>Journal of Hypertension</i> , 2012 , 30, 2299-306	1.9	59
13	Comparability in coronary artery calcium scores on CT scan between two community-based cohort studies. <i>International Journal of Cardiology</i> , 2011 , 149, 244-245	3.2	7
12	Relationship of moderate metabolic risk factor clustering to cardiovascular disease mortality in non-lean Japanese: a 15-year follow-up of NIPPON DATA90. <i>Atherosclerosis</i> , 2011 , 215, 209-13	3.1	8
11	J-005 AN INCREASED SERUM ALANINE AMINOTRANSFERASE IS ASSOCIATED WITH AN INCREASED MORTALITY FROM INTRACEREBRAL HEMORRHAGE IN THE EAST ASIAN POPULATION. <i>Journal of Hypertension</i> , 2011 , 29, e25	1.9	
10	Metabolic syndrome and its components are underdiagnosed in cardiology clinics. <i>Journal of Evaluation in Clinical Practice</i> , 2011 , 17, 78-83	2.5	4
9	Glutamyltransferase and mortality risk from heart disease and stroke in Japanese men and women: NIPPON DATA90. <i>CVD Prevention and Control</i> , 2010 , 5, 27-34		5
8	Risk factors for heart failure and coronary heart disease mortality over 24-year follow-up period in Japan: NIPPON DATA80. <i>CVD Prevention and Control</i> , 2010 , 5, 97-103		4
7	Comparison of the National Nutritional Survey in Japan estimated individual-based nutritional data and NIPPON DATA80 food frequency questionnaires. <i>Journal of Epidemiology</i> , 2010 , 20 Suppl 3, S582-6	3.4	7
6	Fatty acids intakes and serum lipid profiles: NIPPON DATA90 and the national nutrition monitoring. <i>Journal of Epidemiology</i> , 2010 , 20 Suppl 3, S544-8	3.4	13

5	Population attributable fraction of smoking and metabolic syndrome on cardiovascular disease mortality in Japan: a 15-year follow up of NIPPON DATA90. <i>BMC Public Health</i> , 2010 , 10, 306	4.1	13
4	Patient-important outcomes in registered diabetes trials. <i>JAMA - Journal of the American Medical Association</i> , 2008 , 299, 2543-9	27.4	159
3	Caloric restriction, the traditional Okinawan diet, and healthy aging: the diet of the world's longest-lived people and its potential impact on morbidity and life span. <i>Annals of the New York Academy of Sciences</i> , 2007 , 1114, 434-55	6.5	216
2	Two Cases of Myxedema Coma Successfully Treated with Low-Dose Thyroid Hormone Replacement.. <i>Nihon Kyukyu Igakukai Zasshi</i> , 2000 , 11, 22-26	0	
1	Factors associated with mortality of myxedema coma: report of eight cases and literature survey. <i>Thyroid</i> , 1999 , 9, 1167-74	6.2	99