Fabiola Carrara

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5171953/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Glomerular Hyperfiltration and Renal Disease Progression in Type 2 Diabetes. Diabetes Care, 2012, 35, 2061-2068. | 8.6 | 259 |
| 2 | Effect of longacting somatostatin analogue on kidney and cyst growth in autosomal dominant polycystic kidney disease (ALADIN): a randomised, placebo-controlled, multicentre trial. Lancet, The, 2013, 382, 1485-1495. | 13.7 | 218 |
| 3 | Estimated GFR: time for a critical appraisal. Nature Reviews Nephrology, 2019, 15, 177-190. | 9.6 | 187 |
| 4 | Performance of Different Prediction Equations for Estimating Renal Function in Kidney Transplantation. American Journal of Transplantation, 2004, 4, 1826-1835. | 4.7 | 184 |
| 5 | The GFR and GFR decline cannot be accurately estimated in type 2 diabetics. Kidney International, 2013, 84, 164-173. | 5.2 | 131 |
| 6 | Unlike each drug alone, lisinopril if combined with avosentan promotes regression of renal lesions in experimental diabetes. American Journal of Physiology - Renal Physiology, 2009, 297, F1448-F1456. | 2.7 | 114 |
| 7 | Analogs of bardoxolone methyl worsen diabetic nephropathy in rats with additional adverse effects. American Journal of Physiology - Renal Physiology, 2013, 304, F808-F819. | 2.7 | 90 |
| 8 | Renal and Systemic Effects of Calorie Restriction in Patients With Type 2 Diabetes With Abdominal Obesity: A Randomized Controlled Trial. Diabetes, 2017, 66, 75-86. | 0.6 | 66 |
| 9 | Therapeutic Drug Monitoring of Sirolimus: Effect of Concomitant Immunosuppressive Therapy and Optimization of Drug Dosing. American Journal of Transplantation, 2004, 4, 1345-1351. | 4.7 | 57 |
| 10 | Measuring and Estimating GFR and Treatment Effect in ADPKD Patients: Results and Implications of a Longitudinal Cohort Study. PLoS ONE, 2012, 7, e32533. | 2.5 | 46 |
| 11 | C5 Convertase Blockade in Membranoproliferative Glomerulonephritis: A Single-Arm Clinical Trial. American Journal of Kidney Diseases, 2019, 74, 224-238. | 1.9 | 45 |
| 12 | Octreotide-LAR in later-stage autosomal dominant polycystic kidney disease (ALADIN 2): A randomized, double-blind, placebo-controlled, multicenter trial. PLoS Medicine, 2019, 16, e1002777. | 8.4 | 42 |
| 13 | Effect of Sirolimus on Disease Progression in Patients with Autosomal Dominant Polycystic Kidney Disease and CKD Stages 3b-4. Clinical Journal of the American Society of Nephrology: CJASN, 2016, 11, 785-794. | 4.5 | 35 |
| 14 | Estimated Glomerular Filtration Rate in Renal Transplantation. Transplantation, 2015, 99, 2625-2633. | 1.0 | 30 |
| 15 | lohexol plasma clearance simplified by dried blood spot testing. Nephrology Dialysis Transplantation, 2018, 33, 1597-1603. | 0.7 | 29 |
| 16 | Moderate salt restriction with or without paricalcitol in type 2 diabetes and losartan-resistant macroalbuminuria (PROCEED): a randomised, double-blind, placebo-controlled, crossover trial. Lancet Diabetes and Endocrinology,the, 2018, 6, 27-40. | 11.4 | 24 |
| 17 | Safety of Iohexol Administration to Measure Clomerular Filtration Rate in Different Patient Populations: A 25-Year Experience. Nephron, 2018, 140, 1-8. | 1.8 | 21 |
| 18 | Effects of Sevelamer Carbonate in Patients With CKD and Proteinuria: The ANSWER Randomized Trial. American Journal of Kidney Diseases, 2019, 74, 338-350. | 1.9 | 17 |

FABIOLA CARRARA

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Measurement of glomerular filtration rate: Internal and external validations of the iohexol plasma clearance technique by HPLC. Clinica Chimica Acta, 2014, 430, 84-85. | 1.1 | 16 |
| 20 | Effects of valsartan, benazepril and their combination in overt nephropathy of type 2 diabetes: A prospective, randomized, controlled trial. Diabetes, Obesity and Metabolism, 2019, 21, 1177-1190. | 4.4 | 14 |
| 21 | Effect of ACE inhibition on glomerular permselectivity and tubular albumin concentration in the renal ablation model. American Journal of Physiology - Renal Physiology, 2011, 300, F1291-F1300. | 2.7 | 13 |
| 22 | Iohexol plasma clearance, a simple and reliable method to measure renal function in conscious mice. Pflugers Archiv European Journal of Physiology, 2016, 468, 1587-1594. | 2.8 | 13 |
| 23 | Effects of Rosuvastatin on Glomerular Capillary Size-Selectivity Function in Rats with Renal Mass Ablation. American Journal of Nephrology, 2007, 27, 630-638. | 3.1 | 12 |
| 24 | Determination of iohexol in human serum by a semi-automated liquid chromatography tandem mass spectrometry method. Clinical Biochemistry, 2015, 48, 679-685. | 1.9 | 12 |
| 25 | GFR measured by iohexol: the best choice from a laboratory perspective. Journal of Laboratory and Precision Medicine, 0, 3, 77-77. | 1.1 | 12 |
| 26 | A Simple Method to Measure Renal Function in Swine by the Plasma Clearance of Iohexol. International Journal of Molecular Sciences, 2018, 19, 232. | 4.1 | 10 |
| 27 | Simplified Method to Measure Glomerular Filtration Rate by Iohexol Plasma Clearance in Conscious Rats. Nephron, 2016, 133, 62-70. | 1.8 | 9 |
| 28 | Preventing microalbuminuria with benazepril, valsartan, and benazepril–valsartan combination therapy in diabetic patients with high-normal albuminuria: A prospective, randomized, open-label, blinded endpoint (PROBE) study. PLoS Medicine, 2021, 18, e1003691. | 8.4 | 7 |
| 29 | Reply to â€~Strengths and limitations of estimated and measured GFR'. Nature Reviews Nephrology, 2019, 15, 785-786. | 9.6 | 5 |
| 30 | Conversion from Brand-Name Neoral to the Generic Ciqorin in Stable Renal Transplant Recipients. Nephron, 2017, 135, 173-180. | 1.8 | 4 |
| 31 | Preimplantation Histological Score Associates with 6-Month GFR in Recipients of Perfused, Older Kidney Grafts: Results from a Pilot Study. Nephron, 2021, 145, 137-149. | 1.8 | 3 |
| 32 | Long-Term Outcomes of Kidney Transplants from Older/Marginal Donors: A Cohort Study. Nephron, 2021, 145, 642-652. | 1.8 | 3 |
| 33 | SaO006INCREASED PRE-GLOMERULAR RESISTANCE AND KIDNEY HYPOPERFUSION MAY SUSTAIN ACCELERATED GFR DECLINE IN HYPERTENSIVE, TYPE 2 DIABETICS WITH NORMAL AND HIGH NORMAL ALBUMINURIA. Nephrology Dialysis Transplantation, 2018, 33, i317-i317. | 0.7 | 2 |
| 34 | Glomerular resistances predict long-term GFR decline in type 2 diabetic patients without overt nephropathy: a longitudinal subgroup analysis of the DEMAND trial. Acta Diabetologica, 2022, 59, 309-317. | 2.5 | 2 |
| 35 | Response to "lohexol-measured glomerular filtration rate in children and adolescents with chronic kidney disease: a pilot study comparing venous and finger stick methods― Pediatric Nephrology, 2019, 34, 1629-1630. | 1.7 | 0 |