

# Christopher Grefen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5181678/publications.pdf>

Version: 2024-02-01

45  
papers

5,642  
citations

136950

32  
h-index

233421

45  
g-index

48  
all docs

48  
docs citations

48  
times ranked

7510  
citing authors

#	ARTICLE	IF	CITATIONS
1	Looking for a safe haven: tail-anchored proteins and their membrane insertion pathways. <i>Plant Physiology</i> , 2021, 187, 1916-1928.	4.8	9
2	Endoplasmic reticulum membrane receptors of the GET pathway are conserved throughout eukaryotes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	7.1	13
3	Distinct RopGEFs Successively Drive Polarization and Outgrowth of Root Hairs. <i>Current Biology</i> , 2019, 29, 1854-1865.e5.	3.9	78
4	Constitutive signaling activity of a receptor-associated protein links fertilization with embryonic patterning in <i>Arabidopsis thaliana</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 5795-5804.	7.1	39
5	The GET pathway can increase the risk of mitochondrial outer membrane proteins to be mistargeted to the ER. <i>Journal of Cell Science</i> , 2018, 131, .	2.0	34
6	2in1 Vectors Improve In Planta BiFC and FRET Analyses. <i>Methods in Molecular Biology</i> , 2018, 1691, 139-158.	0.9	24
7	ER Membrane Protein Interactions Using the Split-Ubiquitin System (SUS). <i>Methods in Molecular Biology</i> , 2018, 1691, 191-203.	0.9	5
8	Detecting Interactions of Membrane Proteins: The Split-Ubiquitin System. <i>Methods in Molecular Biology</i> , 2018, 1794, 49-60.	0.9	4
9	Loss of GET pathway orthologs in <i>Arabidopsis thaliana</i> causes root hair growth defects and affects SNARE abundance. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E1544-E1553.	7.1	56
10	ATP sensing in living plant cells reveals tissue gradients and stress dynamics of energy physiology. <i>ELife</i> , 2017, 6, .	6.0	125
11	Techniques for the analysis of protein-protein interactions in vivo. <i>Plant Physiology</i> , 2016, 171, pp.00470.2016.	4.8	177
12	An RLP23-SOBIR1-BAK1 complex mediates NLP-triggered immunity. <i>Nature Plants</i> , 2015, 1, 15140.	9.3	373
13	A vesicle-trafficking protein commandeers Kv channel voltage sensors for voltage-dependent secretion. <i>Nature Plants</i> , 2015, 1, 15108.	9.3	53
14	Binding of SEC11 Indicates Its Role in SNARE Recycling after Vesicle Fusion and Identifies Two Pathways for Vesicular Traffic to the Plasma Membrane. <i>Plant Cell</i> , 2015, 27, 675-694.	6.6	55
15	The Arabidopsis R-SNARE VAMP721 Interacts with KAT1 and KC1 K+ Channels to Moderate K+ Current at the Plasma Membrane. <i>Plant Cell</i> , 2015, 27, 1697-1717.	6.6	84
16	Binary 2in1 Vectors Improve in Planta (Co)localization and Dynamic Protein Interaction Studies. <i>Plant Physiology</i> , 2015, 168, 776-787.	4.8	84
17	Voltage-Sensor Transitions of the Inward-Rectifying K+ Channel KAT1 Indicate a Latching Mechanism Biased by Hydration within the Voltage Sensor Å Å. <i>Plant Physiology</i> , 2014, 166, 960-975.	4.8	21
18	The Golgi S-acylation machinery comprises zDHHC enzymes with major differences in substrate affinity and S-acylation activity. <i>Molecular Biology of the Cell</i> , 2014, 25, 3870-3883.	2.1	62

#	ARTICLE	IF	CITATIONS
19	Applications of Fluorescent Marker Proteins in Plant Cell Biology. <i>Methods in Molecular Biology</i> , 2014, 1062, 487-507.	0.9	31
20	The Split-Ubiquitin System for the Analysis of Three-Component Interactions. <i>Methods in Molecular Biology</i> , 2014, 1062, 659-678.	0.9	17
21	<i>Arabidopsis</i> SNAREs SYP61 and SYP121 Coordinate the Trafficking of Plasma Membrane Aquaporin PIP2;7 to Modulate the Cell Membrane Water Permeability. <i>Plant Cell</i> , 2014, 26, 3132-3147.	6.6	192
22	Protein Delivery to Vacuole Requires SAND Protein-Dependent Rab GTPase Conversion for MVB-Vacuole Fusion. <i>Current Biology</i> , 2014, 24, 1383-1389.	3.9	144
23	SDM-Assist software to design site-directed mutagenesis primers introducing restriction sites. <i>BMC Bioinformatics</i> , 2013, 14, 105.	2.6	34
24	<i>Arabidopsis</i> Sec1/Munc18 Protein SEC11 Is a Competitive and Dynamic Modulator of SNARE Binding and SYP121-Dependent Vesicle Traffic. <i>Plant Cell</i> , 2013, 25, 1368-1382.	6.6	66
25	Do Calcineurin B-Like Proteins Interact Independently of the Serine Threonine Kinase CIPK23 with the K <sup>+</sup> Channel AKT1? Lessons Learned from a <i>Maize</i> . <i>Plant Physiology</i> , 2012, 159, 915-919.	4.8	46
26	Palmitoylation-induced Aggregation of Cysteine-string Protein Mutants That Cause Neuronal Ceroid Lipofuscinosis. <i>Journal of Biological Chemistry</i> , 2012, 287, 37330-37339.	3.4	57
27	Selective Regulation of Maize Plasma Membrane Aquaporin Trafficking and Activity by the SNARE SYP121. <i>Plant Cell</i> , 2012, 24, 3463-3481.	6.6	109
28	A 2in1 cloning system enables ratiometric bimolecular fluorescence complementation (rBiFC). <i>BioTechniques</i> , 2012, 53, 311-314.	1.8	178
29	The trafficking protein SYP121 of <i>Arabidopsis</i> connects programmed stomatal closure and K <sup>+</sup> channel activity with vegetative growth. <i>Plant Journal</i> , 2012, 69, 241-251.	5.7	115
30	A bicistronic, <i>Ubiquitin-10</i> promoter-based vector cassette for transient transformation and functional analysis of membrane transport demonstrates the utility of quantitative voltage clamp studies on intact <i>Arabidopsis</i> root epidermis. <i>Plant, Cell and Environment</i> , 2011, 34, 554-564.	5.7	12
31	Ion transport, membrane traffic and cellular volume control. <i>Current Opinion in Plant Biology</i> , 2011, 14, 332-339.	7.1	29
32	A molecular framework for coupling cellular volume and osmotic solute transport control. <i>Journal of Experimental Botany</i> , 2011, 62, 2363-2370.	4.8	35
33	Evidence for the localization of the <i>Arabidopsis</i> cytokinin receptors AHK3 and AHK4 in the endoplasmic reticulum. <i>Journal of Experimental Botany</i> , 2011, 62, 5571-5580.	4.8	155
34	A ubiquitin-10 promoter-based vector set for fluorescent protein tagging facilitates temporal stability and native protein distribution in transient and stable expression studies. <i>Plant Journal</i> , 2010, 64, 355-365.	5.7	499
35	A Novel Motif Essential for SNARE Interaction with the K <sup>+</sup> Channel KC1 and Channel Gating in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2010, 22, 3076-3092.	6.6	119
36	A Tripartite SNARE-K <sup>+</sup> Channel Complex Mediates in Channel-Dependent K <sup>+</sup> Nutrition in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2009, 21, 2859-2877.	6.6	156

#	ARTICLE	IF	CITATIONS
37	The Determination of Protein-protein Interactions by the Mating-based Split-ubiquitin system (mbSUS). <i>Methods in Molecular Biology</i> , 2009, 479, 217-233.	0.9	94
38	The Arabidopsis thaliana response regulator ARR22 is a putative AHP phospho-histidine phosphatase expressed in the chalaza of developing seeds. <i>BMC Plant Biology</i> , 2008, 8, 77.	3.6	88
39	SNAREsâ€”molecular governors in signalling and development. <i>Current Opinion in Plant Biology</i> , 2008, 11, 600-609.	7.1	49
40	Subcellular Localization and In Vivo Interactions of the Arabidopsis thaliana Ethylene Receptor Family Members. <i>Molecular Plant</i> , 2008, 1, 308-320.	8.3	207
41	The Histidine Kinase AHK5 Integrates Endogenous and Environmental Signals in Arabidopsis Guard Cells. <i>PLoS ONE</i> , 2008, 3, e2491.	2.5	138
42	Splitâ€”Ubiquitin System for Identifying Proteinâ€”Protein Interactions in Membrane and Fullâ€”Length Proteins. <i>Current Protocols in Neuroscience</i> , 2007, 41, Unit 5.27.	2.6	64
43	Functional cross-talk between two-component and phytochrome B signal transduction in Arabidopsis. <i>Journal of Experimental Botany</i> , 2007, 58, 2595-2607.	4.8	64
44	Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. <i>Plant Journal</i> , 2004, 40, 428-438.	5.7	1,514
45	Plant two-component systems: principles, functions, complexity and cross talk. <i>Planta</i> , 2004, 219, 733-42.	3.2	133