

Miwa Yamaguchi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/518722/publications.pdf>

Version: 2024-02-01

17
papers

386
citations

932766

10
h-index

887659

17
g-index

17
all docs

17
docs citations

17
times ranked

542
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence of Frailty Assessed by Fried and Kihon Checklist Indexes in a Prospective Cohort Study: Design and Demographics of the Kyoto-Kameoka Longitudinal Study. <i>Journal of the American Medical Directors Association</i> , 2017, 18, 733.e7-733.e15.	1.2	68
2	Eating healthy to impress: How conspicuous consumption, perceived self-control motivation, and descriptive normative influence determine functional food choices. <i>Appetite</i> , 2018, 131, 59-67.	1.8	59
3	Association of social jetlag with metabolic syndrome among Japanese working population: the Furukawa Nutrition and Health Study. <i>Sleep Medicine</i> , 2018, 51, 53-58.	0.8	47
4	Community Social Capital and Depressive Symptoms Among Older People in Japan: A Multilevel Longitudinal Study. <i>Journal of Epidemiology</i> , 2019, 29, 363-369.	1.1	41
5	Sex Difference in the Association Between Protein Intake and Frailty: Assessed Using the Kihon Checklist Indexes Among Older Adults. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 801-805.	1.2	26
6	Validation of Energy and Nutrition Intake in Japanese Elderly Individuals Estimated Based on a Short Food Frequency Questionnaire Compared against a 7-day Dietary Record: The Kyoto-Kameoka Study. <i>Nutrients</i> , 2019, 11, 688.	1.7	24
7	Estimation of Energy Intake by a Food Frequency Questionnaire: Calibration and Validation with the Doubly Labeled Water Method in Japanese Older People. <i>Nutrients</i> , 2019, 11, 1546.	1.7	22
8	Sociodemographic and physical predictors of non-participation in community based physical checkup among older neighbors: a case-control study from the Kyoto-Kameoka longitudinal study, Japan. <i>BMC Public Health</i> , 2018, 18, 568.	1.2	19
9	Association between the Frequency of Protein-Rich Food Intakes and Kihon-Checklist Frailty Indices in Older Japanese Adults: The Kyoto-Kameoka Study. <i>Nutrients</i> , 2018, 10, 84.	1.7	17
10	Comparison of Objective and Perceived Access to Food Stores Associated with Intake Frequencies of Vegetables/Fruits and Meat/Fish among Community-Dwelling Older Japanese. <i>International Journal of Environmental Research and Public Health</i> , 2019, 16, 772.	1.2	12
11	Healthy lifestyle index and its association with hypertension among community adults in Sri Lanka: A cross-sectional study. <i>PLoS ONE</i> , 2020, 15, e0226773.	1.1	12
12	BMI and Medically Certified Long-Term Sickness Absence Among Japanese Employees. <i>Obesity</i> , 2020, 28, 437-444.	1.5	10
13	The effect of youths as change agents on cardiovascular disease risk factors among adult neighbours: a cluster randomised controlled trial in Sri Lanka. <i>BMC Public Health</i> , 2019, 19, 893.	1.2	8
14	Measures of Perceived Neighborhood Food Environments and Dietary Habits: A Systematic Review of Methods and Associations. <i>Nutrients</i> , 2022, 14, 1788.	1.7	8
15	The association of work-related stressors and their changes over time with the development of metabolic syndrome: The Furukawa Nutrition and Health Study. <i>Journal of Occupational Health</i> , 2018, 60, 485-493.	1.0	6
16	An assessment of implementation gaps and priority recommendations on food environment policies: the Healthy Food Environment Policy Index (Food-EPI) in Japan. <i>Public Health Nutrition</i> , 2021, , 1-36.	1.1	6
17	The Role of the School Food Environment in Improving the Healthiness of School Canteens and Readiness to Reopen Post COVID-19 Pandemic: A Study Conducted in Indonesia. <i>Journal of Public Health Research</i> , 2022, 11, jphr.2021.2287.	0.5	1