

Philippe Conus

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5187414/philippe-conus-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

192
papers

6,153
citations

45
h-index

73
g-index

221
ext. papers

7,573
ext. citations

4.5
avg, IF

5.56
L-index

#	Paper	IF	Citations
192	N-acetyl cysteine as a glutathione precursor for schizophrenia--a double-blind, randomized, placebo-controlled trial. <i>Biological Psychiatry</i> , 2008 , 64, 361-8	7.9	415
191	Glutathione precursor, N-acetyl-cysteine, improves mismatch negativity in schizophrenia patients. <i>Neuropsychopharmacology</i> , 2008 , 33, 2187-99	8.7	284
190	Impaired glutathione synthesis in schizophrenia: convergent genetic and functional evidence. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 16621-6	11.5	247
189	The impact of cannabis use on cognitive functioning in patients with schizophrenia: a meta-analysis of existing findings and new data in a first-episode sample. <i>Schizophrenia Bulletin</i> , 2012 , 38, 316-30	1.3	190
188	A 600 kb deletion syndrome at 16p11.2 leads to energy imbalance and neuropsychiatric disorders. <i>Journal of Medical Genetics</i> , 2012 , 49, 660-8	5.8	182
187	Setting the stage: from prodrome to treatment resistance in bipolar disorder. <i>Bipolar Disorders</i> , 2007 , 9, 671-8	3.8	168
186	Pre-treatment, baseline, and outcome differences between early-onset and adult-onset psychosis in an epidemiological cohort of 636 first-episode patients. <i>Schizophrenia Research</i> , 2007 , 95, 1-8	3.6	142
185	Does stage of illness impact treatment response in bipolar disorder? Empirical treatment data and their implication for the staging model and early intervention. <i>Bipolar Disorders</i> , 2011 , 13, 87-98	3.8	135
184	Developing psychosis and its risk states through the lens of schizotypy. <i>Schizophrenia Bulletin</i> , 2015 , 41 Suppl 2, S396-407	1.3	120
183	Defining the Effect of the 16p11.2 Duplication on Cognition, Behavior, and Medical Comorbidities. <i>JAMA Psychiatry</i> , 2016 , 73, 20-30	14.5	120
182	Stage managing bipolar disorder. <i>Bipolar Disorders</i> , 2014 , 16, 471-7	3.8	112
181	Impact of duration of untreated psychosis on pre-treatment, baseline, and outcome characteristics in an epidemiological first-episode psychosis cohort. <i>Journal of Psychiatric Research</i> , 2008 , 42, 982-90	5.2	111
180	Diagnostic stability 18 months after treatment initiation for first-episode psychosis. <i>Journal of Clinical Psychiatry</i> , 2005 , 66, 1239-46	4.6	106
179	Pretreatment and outcome correlates of sexual and physical trauma in an epidemiological cohort of first-episode psychosis patients. <i>Schizophrenia Bulletin</i> , 2010 , 36, 1105-14	1.3	99
178	Is diagnosis enough to guide interventions in mental health? Using case formulation in clinical practice. <i>BMC Medicine</i> , 2012 , 10, 111	11.4	93
177	Rate and predictors of service disengagement in an epidemiological first-episode psychosis cohort. <i>Schizophrenia Research</i> , 2010 , 118, 256-63	3.6	87
176	N-acetylcysteine in a Double-Blind Randomized Placebo-Controlled Trial: Toward Biomarker-Guided Treatment in Early Psychosis. <i>Schizophrenia Bulletin</i> , 2018 , 44, 317-327	1.3	83

175	M3. Topological Analyses of Metabolomic Data to Identify Markers of Early Psychosis and Disease Biotypes. <i>Schizophrenia Bulletin</i> , 2017 , 43, S211-S212	1.3	78
174	29.3 TOPOLOGICAL ANALYSIS ENABLES PREDICTION OF FUNCTIONAL AND SOCIAL OUTCOMES IN EARLY PSYCHOSIS. <i>Schizophrenia Bulletin</i> , 2019 , 45, S137-S137	1.3	78
173	M16. PROFILING GLUTAMATE AND D-SERINE PATHWAYS IN TREATMENT RESISTANT EARLY PSYCHOSIS PATIENTS. <i>Schizophrenia Bulletin</i> , 2020 , 46, S139-S139	1.3	78
172	SA58. The First-Episode Psychosis Outcome Study: Preliminary Data on the Long-Term Follow-Up of First-Episode Psychosis Patients Who Were Treated at the Early Psychosis Prevention and Intervention Centre Between 1998 and 2000. <i>Schizophrenia Bulletin</i> , 2017 , 43, S134-S134	1.3	78
171	16.2 CHILDHOOD TRAUMA ENGAGES OXIDATIVE STRESS, HIPPOCAMPUS ALTERATIONS, AND POORER CLINICAL OUTCOME IN EARLY PSYCHOSIS PATIENTS. <i>Schizophrenia Bulletin</i> , 2018 , 44, S25-S26	1.3	78
170	S69. CLINICAL HIGH RISK STATE: STRATIFICATION BASED ON CLINICAL PROFILE AND REDOX STATUS. <i>Schizophrenia Bulletin</i> , 2020 , 46, S60-S60	1.3	78
169	S143. NEURAL MECHANISMS OF ROBOT-INDUCED HALLUCINATIONS IN HEALTHY PARTICIPANTS AND SYMPTOMATIC HALLUCINATIONS OF NEUROLOGICAL AND PSYCHIATRIC ORIGIN. <i>Schizophrenia Bulletin</i> , 2020 , 46, S90-S91	1.3	78
168	S164. AT-RISK MENTAL STATES PROGRAM IN LAUSANNE: INFLUENCE OF RECRUITMENT STRATEGIES ON THE RATE OF FALSE POSITIVES. <i>Schizophrenia Bulletin</i> , 2018 , 44, S389-S389	1.3	78
167	10.2 REDOX DYSREGULATION, OLIGODENDROCYTES AND WHITE MATTER ALTERATIONS IN SCHIZOPHRENIA. <i>Schizophrenia Bulletin</i> , 2018 , 44, S15-S16	1.3	78
166	The proximal prodrome to first episode mania--a new target for early intervention. <i>Bipolar Disorders</i> , 2008 , 10, 555-65	3.8	74
165	Predictors of service disengagement in first-admitted adolescents with psychosis. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2006 , 45, 990-999	7.2	70
164	From neuroprogression to neuroprotection: implications for clinical care. <i>Medical Journal of Australia</i> , 2010 , 193, S36-40	4	67
163	Pretreatment and outcome correlates of past sexual and physical trauma in 118 bipolar I disorder patients with a first episode of psychotic mania. <i>Bipolar Disorders</i> , 2010 , 12, 244-52	3.8	65
162	The First-Episode Psychosis Outcome Study: premorbid and baseline characteristics of an epidemiological cohort of 661 first-episode psychosis patients. <i>Microbial Biotechnology</i> , 2007 , 1, 191-200	3.3	64
161	Symptomatic and functional outcome 12 months after a first episode of psychotic mania: barriers to recovery in a catchment area sample. <i>Bipolar Disorders</i> , 2006 , 8, 221-31	3.8	62
160	Genetic Polymorphism Associated Prefrontal Glutathione and Its Coupling With Brain Glutamate and Peripheral Redox Status in Early Psychosis. <i>Schizophrenia Bulletin</i> , 2016 , 42, 1185-96	1.3	62
159	Transdiagnostic clinical staging in youth mental health: a first international consensus statement. <i>World Psychiatry</i> , 2020 , 19, 233-242	14.4	61
158	An evidence map of psychosocial interventions for the earliest stages of bipolar disorder. <i>Lancet Psychiatry</i> , 2015 , 2, 548-63	23.3	59

157	Treatment and early intervention in psychosis program (TIPP-Lausanne): Implementation of an early intervention programme for psychosis in Switzerland. <i>Microbial Biotechnology</i> , 2013 , 7, 322-8	3.3	59
156	Prevalence and predictors of suicide attempt in an incidence cohort of 661 young people with first-episode psychosis. <i>Australian and New Zealand Journal of Psychiatry</i> , 2009 , 43, 149-57	2.6	59
155	First-episode mania: a neglected priority for early intervention. <i>Australian and New Zealand Journal of Psychiatry</i> , 2002 , 36, 158-72	2.6	59
154	Randomized controlled trial of interventions for young people at ultra-high risk of psychosis: study design and baseline characteristics. <i>Australian and New Zealand Journal of Psychiatry</i> , 2009 , 43, 818-29	2.6	58
153	Early intervention in bipolar disorders: opportunities and pitfalls. <i>Medical Journal of Australia</i> , 2007 , 187, S11-4	4	58
152	The Number of Genomic Copies at the 16p11.2 Locus Modulates Language, Verbal Memory, and Inhibition. <i>Biological Psychiatry</i> , 2016 , 80, 129-139	7.9	57
151	Characterizing the connectome in schizophrenia with diffusion spectrum imaging. <i>Human Brain Mapping</i> , 2015 , 36, 354-66	5.9	55
150	Characterisation of the prodrome to a first episode of psychotic mania: results of a retrospective study. <i>Journal of Affective Disorders</i> , 2010 , 124, 341-5	6.6	53
149	Genetic dysregulation of glutathione synthesis predicts alteration of plasma thiol redox status in schizophrenia. <i>Antioxidants and Redox Signaling</i> , 2011 , 15, 2003-10	8.4	50
148	Prevalence, predictors, and consequences of long-term refusal of antipsychotic treatment in first-episode psychosis. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 565-72	1.7	46
147	Importance of early weight changes to predict long-term weight gain during psychotropic drug treatment. <i>Journal of Clinical Psychiatry</i> , 2015 , 76, e1417-23	4.6	43
146	Quetiapine lithium in the maintenance phase following a first episode of mania: randomised controlled trial. <i>British Journal of Psychiatry</i> , 2017 , 210, 413-421	5.4	41
145	Public health significance of bipolar disorder: implications for early intervention and prevention. <i>Bipolar Disorders</i> , 2014 , 16, 548-56	3.8	40
144	Cannabis use disorder and age at onset of psychosis--a study in first-episode patients. <i>Schizophrenia Research</i> , 2011 , 129, 52-6	3.6	40
143	Motivational intervention to reduce cannabis use in young people with psychosis: a randomized controlled trial. <i>Psychotherapy and Psychosomatics</i> , 2011 , 80, 287-97	9.4	40
142	MMP9/RAGE pathway overactivation mediates redox dysregulation and neuroinflammation, leading to inhibitory/excitatory imbalance: a reverse translation study in schizophrenia patients. <i>Molecular Psychiatry</i> , 2020 , 25, 2889-2904	15.1	38
141	Influence of CRT1 polymorphisms on body mass index and fat mass in psychiatric patients and the general adult population. <i>JAMA Psychiatry</i> , 2013 , 70, 1011-9	14.5	37
140	Genetics-Based Population Pharmacokinetics and Pharmacodynamics of Risperidone in a Psychiatric Cohort. <i>Clinical Pharmacokinetics</i> , 2015 , 54, 1259-72	6.2	35

139	Pharmacological treatment in the early phase of bipolar disorders: what stage are we at?. <i>Australian and New Zealand Journal of Psychiatry</i> , 2006 , 40, 199-207	2.6	35
138	Early Psychosis Prevention and Intervention Centre long-term follow-up study of first-episode psychosis: methodology and baseline characteristics. <i>Microbial Biotechnology</i> , 2007 , 1, 49-60	3.3	34
137	Pharmacotherapy of first-episode psychosis. <i>Expert Opinion on Pharmacotherapy</i> , 2003 , 4, 717-50	4	34
136	Quantifying the Effects of 16p11.2 Copy Number Variants on Brain Structure: A Multisite Genetic-First Study. <i>Biological Psychiatry</i> , 2018 , 84, 253-264	7.9	33
135	An International Society of Bipolar Disorders task force report: Precursors and prodromes of bipolar disorder. <i>Bipolar Disorders</i> , 2019 , 21, 720-740	3.8	32
134	Impaired metabolic reactivity to oxidative stress in early psychosis patients. <i>Schizophrenia Bulletin</i> , 2014 , 40, 973-83	1.3	31
133	N-acetylcysteine add-on treatment leads to an improvement of fornix white matter integrity in early psychosis: a double-blind randomized placebo-controlled trial. <i>Translational Psychiatry</i> , 2018 , 8, 220	8.6	31
132	Can a targeted psychological intervention be effective for young people following a first manic episode? Results from an 18-month pilot study. <i>Microbial Biotechnology</i> , 2012 , 6, 380-8	3.3	30
131	Unpacking 'the City': An experience-based approach to the role of urban living in psychosis. <i>Health and Place</i> , 2016 , 42, 104-110	4.6	29
130	Psychological needs of adolescents in the early phase of bipolar disorder: implications for early intervention. <i>Microbial Biotechnology</i> , 2011 , 5, 100-7	3.3	29
129	Readiness to participate in psychiatric research. <i>Canadian Journal of Psychiatry</i> , 2003 , 48, 480-4	4.8	29
128	Schneiderian first rank symptoms predict poor outcome within first episode manic psychosis. <i>Journal of Affective Disorders</i> , 2004 , 81, 259-68	6.6	29
127	Mild Depressive Symptoms Mediate the Impact of Childhood Trauma on Long-Term Functional Outcome in Early Psychosis Patients. <i>Schizophrenia Bulletin</i> , 2017 , 43, 1027-1035	1.3	26
126	Duration of untreated psychosis: a proposition regarding treatment definition. <i>Microbial Biotechnology</i> , 2011 , 5, 301-8	3.3	26
125	Difficult-to-engage patients: a specific target for time-limited assertive outreach in a Swiss setting. <i>Canadian Journal of Psychiatry</i> , 2005 , 50, 845-50	4.8	26
124	Duration of untreated psychosis: Impact of the definition of treatment onset on its predictive value over three years of treatment. <i>Journal of Psychiatric Research</i> , 2016 , 77, 15-21	5.2	24
123	Age at the time of exposure to trauma modulates the psychopathological profile in patients with early psychosis. <i>Journal of Clinical Psychiatry</i> , 2016 , 77, e612-8	4.6	23
122	Treatment in early psychosis with N-acetyl-cysteine for 6months improves low-level auditory processing: Pilot study. <i>Schizophrenia Research</i> , 2018 , 191, 80-86	3.6	22

121	Cannabis use in early psychosis is associated with reduced glutamate levels in the prefrontal cortex. <i>Psychopharmacology</i> , 2018 , 235, 13-22	4.7	22
120	Cannabis, a Significant Risk Factor for Violent Behavior in the Early Phase Psychosis. Two Patterns of Interaction of Factors Increase the Risk of Violent Behavior: Cannabis Use Disorder and Impulsivity; Cannabis Use Disorder, Lack of Insight and Treatment Adherence. <i>Frontiers in Psychiatry</i> , 2018 , 9, 234	5	22
119	Rates and predictors of 18-months remission in an epidemiological cohort of 661 patients with first-episode psychosis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017 , 52, 1089-1099	4.5	19
118	Pharmacogenetic study on risperidone long-acting injection: influence of cytochrome P450 2D6 and pregnane X receptor on risperidone exposure and drug-induced side-effects. <i>Journal of Clinical Psychopharmacology</i> , 2013 , 33, 289-98	1.7	19
117	Redox dysregulation as a link between childhood trauma and psychopathological and neurocognitive profile in patients with early psychosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 12495-12500	11.5	19
116	Impulsivity in early psychosis: A complex link with violent behaviour and a target for intervention. <i>European Psychiatry</i> , 2018 , 49, 30-36	6	18
115	N-Acetyl-Cysteine Supplementation Improves Functional Connectivity Within the Cingulate Cortex in Early Psychosis: A Pilot Study. <i>International Journal of Neuropsychopharmacology</i> , 2019 , 22, 478-487	5.8	18
114	High b-value diffusion-weighted imaging: a sensitive method to reveal white matter differences in schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , 2012 , 201, 144-51	2.9	18
113	Brain connectivity alterations in early psychosis: from clinical to neuroimaging staging. <i>Translational Psychiatry</i> , 2019 , 9, 62	8.6	17
112	Olanzapine or chlorpromazine plus lithium in first episode psychotic mania: An 8-week randomised controlled trial. <i>European Psychiatry</i> , 2015 , 30, 975-82	6	17
111	Pre-morbid and outcome correlates of first episode mania with psychosis: is a distinction between schizoaffective and bipolar I disorder valid in the early phase of psychotic disorders?. <i>Journal of Affective Disorders</i> , 2010 , 126, 88-95	6.6	17
110	History of offending behavior in first episode psychosis patients: a marker of specific clinical needs and a call for early detection strategies among young offenders. <i>Schizophrenia Research</i> , 2015 , 161, 163-8	3.6	16
109	Sensorimotor Induction of Auditory Misattribution in Early Psychosis. <i>Schizophrenia Bulletin</i> , 2020 , 46, 947-954	1.3	16
108	Emplacing recovery: how persons diagnosed with psychosis handle stress in cities. <i>Psychosis</i> , 2017 , 9, 322-329	1.2	16
107	Bipolar Disorder in Young People: A Psychological Intervention Manual 2009 ,		16
106	Association of CRTC1 polymorphisms with obesity markers in subjects from the general population with lifetime depression. <i>Journal of Affective Disorders</i> , 2016 , 198, 43-9	6.6	15
105	Predictors of favourable outcome in young people with a first episode psychosis without antipsychotic medication. <i>Schizophrenia Research</i> , 2017 , 185, 130-136	3.6	14
104	A psychoeducation tool for patients with first-episode psychosis. <i>Acta Neuropsychiatrica</i> , 2011 , 23, 75-77	3.9	14

103	Sudden death among young people with first-episode psychosis: An 8-10 year follow-up study. <i>Psychiatry Research</i> , 2010 , 177, 305-8	9.9	14
102	Assertive outreach for "difficult to engage" patients: A useful tool for a subgroup of patients in specialized early psychosis intervention programs. <i>Psychiatry Research</i> , 2016 , 239, 212-9	9.9	14
101	Early changes of blood lipid levels during psychotropic drug treatment as predictors of long-term lipid changes and of new onset dyslipidemia. <i>Journal of Clinical Lipidology</i> , 2018 , 12, 219-229	4.9	14
100	Gender differences in first episode psychotic mania. <i>BMC Psychiatry</i> , 2013 , 13, 82	4.2	13
99	Violent behaviour in early psychosis patients: Can we identify clinical risk profiles?. <i>Microbial Biotechnology</i> , 2019 , 13, 517-524	3.3	12
98	Understanding urbanicity: how interdisciplinary methods help to unravel the effects of the city on mental health. <i>Psychological Medicine</i> , 2021 , 51, 1099-1110	6.9	12
97	Second-Generation Antipsychotics in Adolescent Psychiatric Patients: Metabolic Effects and Impact of an Early Weight Change to Predict Longer Term Weight Gain. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2018 , 28, 258-265	2.9	11
96	Trajectory and predictors of quality of life in first episode psychotic mania. <i>Journal of Affective Disorders</i> , 2016 , 195, 148-55	6.6	11
95	The Swiss Early Psychosis Project SWEPP: a national network. <i>Microbial Biotechnology</i> , 2012 , 6, 106-11	3.3	11
94	Influence of MCHR2 and MCHR2-AS1 Genetic Polymorphisms on Body Mass Index in Psychiatric Patients and In Population-Based Subjects with Present or Past Atypical Depression. <i>PLoS ONE</i> , 2015 , 10, e0139155	3.7	11
93	Promotion of Wellbeing for Children of Parents With Mental Illness: A Model Protocol for Research and Intervention. <i>Frontiers in Psychiatry</i> , 2019 , 10, 606	5	10
92	Identifying patterns in psychiatric hospital stays with statistical methods: towards a typology of post-deinstitutionalization hospitalization trajectories. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2019 , 54, 1411-1417	4.5	10
91	The impact of past direct-personal traumatic events on 12-month outcome in first episode psychotic mania: trauma and early psychotic mania. <i>Australian and New Zealand Journal of Psychiatry</i> , 2014 , 48, 1017-24	2.6	10
90	Differentiating schizoaffective and bipolar I disorder in first-episode psychotic mania. <i>Schizophrenia Research</i> , 2012 , 140, 31-6	3.6	10
89	Impact of HSD11B1 polymorphisms on BMI and components of the metabolic syndrome in patients receiving psychotropic treatments. <i>Pharmacogenetics and Genomics</i> , 2015 , 25, 246-58	1.9	9
88	Intervention dans la phase pr̄coce des troubles psychotiques : objectifs et organisation du programme TIPP (Traitement et intervention dans la phase pr̄coce des troubles psychotiques) ~Lausanne. <i>Information Psychiatrique</i> , 2010 , 86, 145		9
87	Redox dysregulation and oxidative stress in schizophrenia: nutrigenetics as a challenge in psychiatric disease prevention. <i>World Review of Nutrition and Dietetics</i> , 2010 , 101, 131-153	0.2	9
86	Visuospatial working memory deficits and visual pursuit impairments are not directly related in schizophrenia. <i>Australian and New Zealand Journal of Psychiatry</i> , 2009 , 43, 766-74	2.6	9

85	Internal and Predictive Validity of the French Health of the Nation Outcome Scales: Need for Future Directions. <i>PLoS ONE</i> , 2016 , 11, e0160360	3.7	9
84	Early intervention for bipolar disorder - Do current treatment guidelines provide recommendations for the early stages of the disorder?. <i>Journal of Affective Disorders</i> , 2019 , 257, 669-677	6.6	8
83	Benefits of adjunctive N-acetylcysteine in a sub-group of clozapine-treated individuals diagnosed with schizophrenia. <i>Psychiatry Research</i> , 2015 , 230, 982-3	9.9	8
82	Urbanicity: The need for new avenues to explore the link between urban living and psychosis. <i>Microbial Biotechnology</i> , 2020 , 14, 398-409	3.3	8
81	Change in Emotional and Theory of Mind Processing in Borderline Personality Disorder: A Pilot Study. <i>Journal of Nervous and Mental Disease</i> , 2018 , 206, 935-943	1.8	8
80	Clozapine is more efficacious for tongue dystonia than olanzapine. <i>Psychopharmacology</i> , 2002 , 162, 89	4.7	7
79	Olanzapine in subjects with a first-episode psychosis non-responsive, intolerant or non-compliant to a first-line trial of risperidone. <i>International Journal of Psychiatry in Clinical Practice</i> , 2005 , 9, 244-50	2.4	7
78	Early onset of cannabis use and violent behavior in psychosis. <i>European Psychiatry</i> , 2020 , 63, e78	6	7
77	Prediction of early weight gain during psychotropic treatment using a combinatorial model with clinical and genetic markers. <i>Pharmacogenetics and Genomics</i> , 2016 , 26, 547-557	1.9	7
76	Community Treatment Order: Identifying the need for more evidence based justification of its use in first episode psychosis patients. <i>Schizophrenia Research</i> , 2017 , 185, 67-72	3.6	6
75	Social and academic premorbid adjustment domains predict different functional outcomes among youth with first episode mania. <i>Journal of Affective Disorders</i> , 2017 , 219, 133-140	6.6	6
74	Migration in patients with early psychosis: A 3-year prospective follow-up study. <i>Psychiatry Research</i> , 2019 , 275, 108-114	9.9	6
73	Familial psychopathology in the First Episode Psychosis Outcome Study. <i>Australian and New Zealand Journal of Psychiatry</i> , 2008 , 42, 617-26	2.6	6
72	Caught in vicious circles: a perspective on dynamic feed-forward loops driving oxidative stress in schizophrenia. <i>Molecular Psychiatry</i> , 2021 ,	15.1	6
71	Urban remediation: a new recovery-oriented strategy to manage urban stress after first-episode psychosis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2020 , 55, 273-283	4.5	6
70	Partial-volume modeling reveals reduced gray matter in specific thalamic nuclei early in the time course of psychosis and chronic schizophrenia. <i>Human Brain Mapping</i> , 2020 , 41, 4041-4061	5.9	6
69	The coupling of low-level auditory dysfunction and oxidative stress in psychosis patients. <i>Schizophrenia Research</i> , 2017 , 190, 52-59	3.6	5
68	Stable biomarker identification for predicting schizophrenia in the human connectome. <i>NeuroImage: Clinical</i> , 2020 , 27, 102316	5.3	5

67	Les premiers épisodes psychotiques: de l'importance du case management. <i>Evolution Psychiatrique</i> , 2013 , 78, 41-51	0.4	5
66	Association of PCK1 with Body Mass Index and Other Metabolic Features in Patients With Psychotropic Treatments. <i>Journal of Clinical Psychopharmacology</i> , 2015 , 35, 544-52	1.7	5
65	Predictors of substance use reduction in an epidemiological first-episode psychosis cohort. <i>Microbial Biotechnology</i> , 2014 , 8, 358-65	3.3	5
64	Pharmacological Treatment in the Early Phase of Bipolar Disorders: What Stage are we at?		5
63	Mitochondrial, exosomal miR137-COX6A2 and gamma synchrony as biomarkers of parvalbumin interneurons, psychopathology, and neurocognition in schizophrenia. <i>Molecular Psychiatry</i> , 2021 ,	15.1	5
62	The impact of insight in a first-episode mania with psychosis population on outcome at 18 months. <i>Journal of Affective Disorders</i> , 2014 , 167, 74-9	6.6	4
61	Influence of polygenic risk scores on lipid levels and dyslipidemia in a psychiatric population receiving weight gain-inducing psychotropic drugs. <i>Pharmacogenetics and Genomics</i> , 2017 , 27, 464-472	1.9	4
60	Amisulpride: Real-World Evidence of Dose Adaptation and Effect on Prolactin Concentrations and Body Weight Gain by Pharmacokinetic/Pharmacodynamic Analyses. <i>Clinical Pharmacokinetics</i> , 2020 , 59, 371-382	6.2	4
59	Beyond clinical staging of psychiatric disorders. <i>Lancet Psychiatry</i> , 2015 , 2, e27	23.3	3
58	'Is "early intervention" in bipolar disorder what it claims to be?' Malhi et al. <i>Bipolar Disorders</i> , 2018 , 20, 181-183	3.8	3
57	City Avoidance in the Early Phase of Psychosis: A Neglected Domain of Assessment and a Potential Target for Recovery Strategies. <i>Frontiers in Psychiatry</i> , 2019 , 10, 342	5	3
56	Association of variants in SH2B1 and RABEP1 with worsening of low-density lipoprotein and glucose parameters in patients treated with psychotropic drugs. <i>Gene</i> , 2017 , 628, 8-15	3.8	3
55	Trauma and psychosis: some aspects of a complex relationship. <i>Acta Neuropsychiatrica</i> , 2009 , 21, 148-150	9.9	3
54	Duration of untreated psychosis: What are we talking about?. <i>Acta Neuropsychiatrica</i> , 2009 , 21, 106-8	3.9	3
53	Psychological trauma occurring during adolescence is associated with an increased risk of greater waist circumference in Early Psychosis patients treated with psychotropic medication. <i>PLoS ONE</i> , 2020 , 15, e0242569	3.7	3
52	Fronto-Temporal Disconnection Within the Presence Hallucination Network in Psychotic Patients With Passivity Experiences. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1718-1728	1.3	3
51	Prevalence of ECG abnormalities and risk factors for QTc interval prolongation in hospitalized psychiatric patients. <i>Therapeutic Advances in Psychopharmacology</i> , 2019 , 9, 2045125319891386	4.9	3
50	Association Between Plasma Caffeine and Other Methylxanthines and Metabolic Parameters in a Psychiatric Population Treated With Psychotropic Drugs Inducing Metabolic Disturbances. <i>Frontiers in Psychiatry</i> , 2018 , 9, 573	5	3

49	Patients participating to neurobiological research in early psychosis: A selected subgroup?. <i>Schizophrenia Research</i> , 2018 , 201, 249-253	3.6	2
48	Termination of therapy: what can clinicians do to maximise gains?. <i>Acta Neuropsychiatrica</i> , 2010 , 22, 43-45	3.9	2
47	Redox dysregulation and oxidative stress in schizophrenia: nutrigenetics as a challenge in psychiatric disease prevention. <i>Journal of Nutrigenetics and Nutrigenomics</i> , 2010 , 3, 267-89		2
46	Encoding dysfunctions in a dynamic-static paradigm for visuospatial working memory in first-episode psychosis patients: a 2-year follow-up study. <i>Microbial Biotechnology</i> , 2009 , 3, 44-51	3.3	2
45	Evaluation of Cardiometabolic Risk in a Large Psychiatric Cohort and Comparison With a Population-Based Sample in Switzerland. <i>Journal of Clinical Psychiatry</i> , 2020 , 81,	4.6	2
44	Lipid Disturbances in Adolescents Treated With Second-Generation Antipsychotics: Clinical Determinants of Plasma Lipid Worsening and New-Onset Hypercholesterolemia. <i>Journal of Clinical Psychiatry</i> , 2019 , 80,	4.6	2
43	Moderating role of cannabis use between insight and depression in early psychosis. <i>Schizophrenia Research</i> , 2020 , 215, 61-65	3.6	2
42	Timely N-Acetyl-Cysteine and Environmental Enrichment Rescue Oxidative Stress-Induced Parvalbumin Interneuron Impairments via MMP9/RAGE Pathway: A Translational Approach for Early Intervention in Psychosis. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1782-1794	1.3	2
41	The Collaborative Outcomes study on Health and Functioning during Infection Times in Adults (COH-FIT-Adults): Design and methods of an international online survey targeting physical and mental health effects of the COVID-19 pandemic.. <i>Journal of Affective Disorders</i> , 2021 , 299, 393-393	6.6	2
40	Association of genetic risk scores with body mass index in Swiss psychiatric cohorts. <i>Pharmacogenetics and Genomics</i> , 2016 , 26, 208-17	1.9	2
39	Psychotropic drug-induced genetic-epigenetic modulation of CRT1 gene is associated with early weight gain in a prospective study of psychiatric patients. <i>Clinical Epigenetics</i> , 2019 , 11, 198	7.7	2
38	Topology predicts long-term functional outcome in early psychosis. <i>Molecular Psychiatry</i> , 2021 , 26, 5335-5346	5.4	1
37	Frontal cortical thickness correlates positively with impulsivity in early psychosis male patients. <i>Microbial Biotechnology</i> , 2019 , 13, 848-852	3.3	1
36	Effet des groupes motivationnels pour les personnes atteintes de psychose qui consomment du cannabis. <i>Evolution Psychiatrique</i> , 2013 , 78, 97-106	0.4	1
35	Preventive strategies in bipolar disorders	223-240	1
34	Health-related quality of life and functioning in bipolar disorder: the impact of pharmacotherapy. <i>Expert Review of Pharmacoeconomics and Outcomes Research</i> , 2006 , 6, 509-23	2.2	1
33	Evolution of impulsivity levels in relation to early cannabis use in violent patients in the early phase of psychosis.. <i>Psychological Medicine</i> , 2022 , 1-10	6.9	1
32	Evidence of mediation of severity of anxiety and depressive symptoms between abuse and positive symptoms of psychosis. <i>Journal of Psychiatric Research</i> , 2021 ,	5.2	1

31	L'hôpital psychiatrique : un lieu pour la formation à la psychothérapie des troubles psychotiques. <i>Information Psychiatrique</i> , 2012 , 88, 527		1
30	Quel rôle pour les neurosciences dans l'amélioration du traitement de la phase précoce des troubles psychotiques?. <i>Information Psychiatrique</i> , 2010 , 86, 484		1
29	Trauma et psychose : quelles pistes neurobiologiques?. <i>Information Psychiatrique</i> , 2010 , 86, 505		1
28	Syndrome métabolique et antipsychotiques atypiques : recommandations et prise en charge clinique. <i>Information Psychiatrique</i> , 2011 , 87, 715		1
27	Rate and predictors of disengagement in an early psychosis program with time limited intensification of treatment. <i>Journal of Psychiatric Research</i> , 2020 , 131, 33-38	5.2	1
26	Valproate is associated with early decrease of high-density lipoprotein cholesterol levels in the psychiatric population. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2021 , 129, 26-35	3.1	1
25	Socio-economic position as a moderator of cardiometabolic outcomes in patients receiving psychotropic treatment associated with weight gain: results from a prospective 12-month inception cohort study and a large population-based cohort. <i>Translational Psychiatry</i> , 2021 , 11, 360	8.6	1
24	T52. N-ACETYL-CYSTEINE ADD-ON TREATMENT LEADS TO AN IMPROVEMENT OF FORNIX WHITE MATTER INTEGRITY IN EARLY PSYCHOSIS. <i>Schizophrenia Bulletin</i> , 2018 , 44, S133-S134	1.3	1
23	Metabolomic alteration induced by psychotropic drugs: Short-term metabolite profile as a predictor of weight gain evolution. <i>Clinical and Translational Science</i> , 2021 , 14, 2544-2555	4.9	1
22	Associations Between High Plasma Methylxanthine Levels, Sleep Disorders and Polygenic Risk Scores of Caffeine Consumption or Sleep Duration in a Swiss Psychiatric Cohort.. <i>Frontiers in Psychiatry</i> , 2021 , 12, 756403	5	1
21	Détection et intervention précoce pour la psychose : pourquoi et comment?. <i>Santé Mentale Au Québec</i> , 2021 , 46, 45	0.2	1
20	Caught in vicious circles: a perspective on dynamic feed-forward loops driving oxidative stress in schizophrenia; Response to "Adaptive changes to oxidative stress in schizophrenia by Lena Palaniyappan" .. <i>Molecular Psychiatry</i> , 2022 ,	15.1	1
19	Six months functional response to early psychosis intervention program best predicts outcome after three years. <i>Schizophrenia Research</i> , 2021 , 238, 62-69	3.6	0
18	Mechanisms of change in brief treatments for borderline personality disorder: a protocol of a randomized controlled trial. <i>Trials</i> , 2020 , 21, 335	2.8	0
17	Effect of Quetiapine, from Low to High Dose, on Weight and Metabolic Traits: Results from a Prospective Cohort Study. <i>Pharmacopsychiatry</i> , 2021 , 54, 279-286	2	0
16	A narrative review of intervention in first-episode affective psychoses. <i>Journal of Psychiatric Research</i> , 2021 , 143, 123-137	5.2	0
15	Patients', carers' and clinicians' attitudes towards alternative terms to describe the at-risk for psychosis state. <i>Schizophrenia Research</i> , 2021 , 237, 69-75	3.6	0
14	Intervention précoce pour les premiers épisodes psychotiques d'ici à demain : comment relever les défis liés à son déploiement pour en maximiser les bénéfices?. <i>Santé Mentale Au Québec</i> , 2021 , 46, 391	0.2	0

- 13 Insomnia disorders are associated with increased cardiometabolic disturbances and death risks from cardiovascular diseases in psychiatric patients treated with weight-gain-inducing psychotropic drugs: results from a Swiss cohort.. *BMC Psychiatry*, **2022**, 22, 342 4.2 0
- 12 Qui a besoin de suivi intensif dans le milieu dans un programme spcialis pour les psychoses mergentes?. *Annales Medico-Psychologiques*, **2018**, 176, 74-79 0.2
- 11 Nonpharmalogical Substances for Early Intervention. *Key Issues in Mental Health*, 159-167
- 10 Early Intervention in Bipolar Disorder **2014**, 267-287
- 9 La pharmacoghtique/pharmacoghomique en psychiatrie, perspectives thdriques et perspectives pour l'avenir. *Information Psychiatrique*, **2011**, 87, 697
- 8 Intervention prcoce dans les troubles bipolaires: un dveloppement justifi. *Psn*, **2005**, 3, S69-S78
- 7 Traitement pharmacologique du premier pisode psychotique. *Psn*, **2005**, 3, S80-S91
- 6 Recovering the Capacity to Live outside of a Psychiatric Hospital: Impact of a Specialized Inpatient Program. *Psychiatric Quarterly*, **2021**, 92, 751-759 4.1
- 5 Dveloppement de l'insight dans les troubles psychotiques : objectif ou stratgie thfapeutique ? **2018**, 300-309
- 4 La Rencontre de Pierre Magistretti. *Information Psychiatrique*, **2010**, 86, 529
- 3 Video ethnography: introducing a new tool from research to psychiatric practice?. *Psychosis*, **2021**, 13, 24-34 1.2
- 2 White Matter Alterations Between Brain Network Hubs Underlie Processing Speed Impairment in Patients With Schizophrenia.. *Schizophrenia Bulletin Open*, **2021**, 2, sgab033 2.2
- 1 L'union fait la force : initier un mouvement francophone national et international pour l'implantation de l'intervention prcoce. *Sante Mentale Au Quebec*, **2021**, 46, 365 0.2