

Tatjana Cvrkovic

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/51881/publications.pdf>

Version: 2024-02-01

51
papers

831
citations

516710

16
h-index

526287

27
g-index

51
all docs

51
docs citations

51
times ranked

479
citing authors

#	ARTICLE	IF	CITATIONS
1	Genetic Diversity of Flavescence DorÃ©e Phytoplasmas in Vineyards of Serbia: From the Widespread Occurrence of Autochthonous Map-M51 to the Emergence of Endemic Map-FD2 (Vectotype II) and New Map-FD3 (Vectotype III) Epidemic Genotypes. <i>Agronomy</i> , 2022, 12, 448.	3.0	6
2	Integrative Taxonomy and Synonymization of <i>Aculus mosoniensis</i> (Acari: Eriophyidae), a Potential Biological Control Agent for Tree of Heaven (<i>Ailanthus altissima</i>). <i>Insects</i> , 2022, 13, 489.	2.2	4
3	Molecular phylogeny of the phytoparasitic mite family Phytoptidae (Acariformes: Eriophyoidea) identified the female genitalic anatomy as a major macroevolutionary factor and revealed multiple origins of gall induction. <i>Experimental and Applied Acarology</i> , 2021, 83, 31-68.	1.6	16
4	Symptomatology, (Co)occurrence and Differential Diagnostic PCR Identification of â€Ca. <i>Phytoplasma solani</i> â€™ and â€Ca. <i>Phytoplasma convolvuli</i> â€™ in Field Bindweed. <i>Pathogens</i> , 2021, 10, 160.	2.8	4
5	Field Assessment of the Host Range of <i>Aculus mosoniensis</i> (Acari: Eriophyidae), a Biological Control Agent of the Tree of Heaven (<i>Ailanthus altissima</i>). <i>Insects</i> , 2021, 12, 637.	2.2	7
6	The â€code redâ€™ for Balkan vineyards: occurrence of <i>Orienteus ishidae</i> (Matsumura, 1902) (Hemiptera: Tj ETQq1 1 0.784314 rgBT /Overlock		
7	Diversity of phytoplasmas identified in the polyphagous leafhopper <i>Euscelis incisus</i> (Cicadellidae,) Tj ETQq1 1 0.784314 rgBT /Overlock <i>European Journal of Plant Pathology</i> , 2020, 156, 201-221.	1.7	14
8	Molecular phylogenetic analyses reveal a deep dichotomy in the conifer-inhabiting genus <i>Trisetacus</i> (Eriophyoidea: Nalepellidae), with the two lineages differing in their female genital morphology and host associations. <i>Experimental and Applied Acarology</i> , 2020, 81, 287-316.	1.6	10
9	Role of plant-specialized <i>Hyalesthes obsoletus</i> associated with <i>Convolvulus arvensis</i> and <i>Crepis foetida</i> in the transmission of â€Candidatus <i>Phytoplasma solani</i> â€™-inflicted bois noir disease of grapevine in Serbia. <i>European Journal of Plant Pathology</i> , 2019, 153, 183-195.	1.7	31
10	<p class="Body">Supplementary description of five species from the genus Cecidophyopsis (Eriophyoidea: Eriophyidae: Cecidophyinae) </p>. <i>Systematic and Applied Acarology</i> , 2019, 24, 1555-1578.	0.5	1
11	<p class="Body">The description and molecular phylogenetic position of a new conifer-associated mite, <i>Setoptus tsugivagus</i> n. sp. (Eriophyoidea, Phytoptidae, Nalepellinae)<p align="center">
. <i>Systematic and Applied Acarology</i> , 2019, 24, 683.	0.5	8
12	<i>Clematis vitalba</i>-sourced â€â€™flavescence dorÃ©eâ€™ phytoplasmas and <i>Wolbachia</i> in naturally infected populations of <i>Dictyophara europaea</i>. <i>Phytopathogenic Mollicutes</i> , 2019, 9, 113.	0.1	0
13	Experimental and molecular evidence of <i>Neoliturus fenestratus</i> role in the transmission of â€estolburâ€™-phytoplasma to lettuce and carrot plants. <i>Phytopathogenic Mollicutes</i> , 2019, 9, 109.	0.1	3
14	Can polyphagous insect vectors contribute to phytoplasma inventory in diverse ecosystems?. <i>Phytopathogenic Mollicutes</i> , 2019, 9, 103.	0.1	0
15	High genetic diversity of â€Candidatus</i> <i>Phytoplasma solani</i> â€™ infecting pepper in Serbia. <i>Phytopathogenic Mollicutes</i> , 2019, 9, 37.	0.1	0
16	<i>Wolbachia</i> infection in natural populations of <i>Dictyophara europaea</i>, an alternative vector of grapevine Flavescence dorÃ©e phytoplasma: effects and interactions. <i>Annals of Applied Biology</i> , 2018, 172, 47-64.	2.5	20
17	A new species of <i>Aculus</i> mite (Acari: Eriophyidae), a potential biocontrol agent for Australian swamp stoncrop, <i>Crassula helmsii</i> (Crassulaceae). <i>Zootaxa</i> , 2018, 4497, 573.	0.5	2
18	A new <i>Aculodes</i> species (Prostigmata: Eriophyoidea: Eriophyidae) associated with medusahead, <i>Taeniatherum caput-medusae</i> (L.) Nevski (Poaceae). <i>Systematic and Applied Acarology</i> , 2018, 23, 1217.	0.5	9

#	ARTICLE	IF	CITATIONS
19	Widespread plant specialization in the polyphagous planthopper <i>Hyalesthes obsoletus</i> (Cixiidae), a major vector of stolbur phytoplasma: Evidence of cryptic speciation. <i>PLoS ONE</i> , 2018, 13, e0196969.	2.5	20
20	New species and records of phytoptids (Eriophyoidea, Phytoptidae) from cinquefoils (Rosaceae), <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7</i>	0.5	2
21	A new <i>Aceria</i> species (Acari:Trombidiformes: Eriophyoidea) from West Asia, a potential biological control agent for the invasive weed camelthorn, <i>Alhagi maurorum</i> Medik. (Leguminosae). <i>Acarologia</i> , 2018, 58, 302-312.	0.6	4
22	Phenetic and phylogenetic relationships among <i>Aceria</i> spp. (Acari: Eriophyoidea) inhabiting species within the family Brassicaceae in Serbia. <i>Experimental and Applied Acarology</i> , 2017, 71, 329-343.	1.6	9
23	<i>Dictyophara europaea</i> (Hemiptera: Fulgoromorpha: Dictyopharidae): description of immatures, biology and host plant associations. <i>Bulletin of Entomological Research</i> , 2016, 106, 395-405.	1.0	15
24	<i>Eriophyid mite Aceria artemisiifoliae</i> sp.nov. (Acari: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7) <i>artemisiifolia</i> L. (Asteraceae) in Serbia. <i>Systematic and Applied Acarology</i> , 2016, 21, 919.	0.5	4
25	â€ˆCandidatus phytoplasma solaniâ€™™ genotypes associated with potato stolbur in Serbia and the role of <i>Hyalesthes obsoletus</i> and <i>Reptalus panzeri</i> (hemiptera, cixiidae) as natural vectors. <i>European Journal of Plant Pathology</i> , 2016, 144, 619-630.	1.7	32
26	Cryptic speciation within <i>Phytoptus avellanae</i> s.l. (Eriophyoidea: Phytoptidae) revealed by molecular data and observations on molting Tegenotus-like nymphs. <i>Experimental and Applied Acarology</i> , 2016, 68, 83-96.	1.6	30
27	<i>Euscelis incisus</i> (Cicadellidae, Deltocephalinae), a natural vector of 16SrIII-B phytoplasma causing multiple inflorescence disease of <i>Cirsium arvense</i> . <i>Annals of Applied Biology</i> , 2015, 167, 406-419.	2.5	8
28	<i>Eriocaenus</i> (Acari: Trombidiformes: Eriophyoidea), a new genus from <i>Equisetum</i> spp. (Equisetaceae): morphological and molecular delimitation of two morphologically similar species. <i>Zootaxa</i> , 2015, 4013, 51-66.	0.5	11
29	Basal divergence of Eriophyoidea (Acariformes, Eupodina) inferred from combined partial COI and 28S gene sequences and CLSM genital anatomy. <i>Experimental and Applied Acarology</i> , 2015, 67, 219-245.	1.6	56
30	The molecular epidemiology of bois noir grapevine yellows caused by â€ˆCandidatus Phytoplasma solaniâ€™™ in the Republic of Macedonia. <i>European Journal of Plant Pathology</i> , 2015, 142, 759-770.	1.7	34
31	A New <i>Metaculus</i> Species (Acari: Eriophyoidea) on <i>Diplotaxis tenuifolia</i> (Brassicaceae) From Serbia: A Combined Description Using Morphology and DNA Barcode Data. <i>Annals of the Entomological Society of America</i> , 2015, 108, 922-931.	2.5	7
32	First Report of â€ˆCandidatus Phytoplasma solaniâ€™™ Infecting Garden Bean <i>Phaseolus vulgaris</i> in Serbia. <i>Plant Disease</i> , 2015, 99, 551-551.	1.4	6
33	Potential Hemipteran vectors of â€ˆstolburâ€™-phytoplasma in potato fields in Serbia. <i>Phytopathogenic Mollicutes</i> , 2015, 5, S49.	0.1	4
34	Experimental and molecular evidence of <i>Reptalus panzeri</i> as a natural vector of bois noir. <i>Plant Pathology</i> , 2014, 63, 42-53.	2.4	92
35	First Report of Alder Yellows Phytoplasma Associated with Common Alder (<i>Alnus glutinosa</i>) in the Republic of Macedonia. <i>Plant Disease</i> , 2014, 98, 1268-1268.	1.4	11
36	<i>Crepis foetida</i> L.: New host plant of cixiid planthopper <i>Hyalesthes obsoletus</i> Signoret 1865 (Hemiptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7)	0.2	4

#	ARTICLE	IF	CITATIONS
37	<i>Drosophila suzukii</i> (Matsumura, 1931) (Siptera: Srosophilidae): A new invasive pest in Serbia. <i>Zastita Bilja</i> , 2014, 65, 99-104.	0.2	18
38	Description of a new relict eriophyoid mite, <i>Loboquintus subsquamatus</i> n. gen. & n. sp. (Eriophyoidea,) Tj ETQq0 0 0 rgBT /Overlock 10 T anatomy of internal genitalia. <i>Experimental and Applied Acarology</i> , 2013, 61, 1-30.	1.6	26
39	First Report of Alder Yellows Phytoplasma Infecting Common and Grey Alder (<i>Alnus glutinosa</i>) Tj ETQq1 1 0.784314 rgBT /Ove	1.4	11
40	Characterisation of a 16SrII phytoplasma strain associated with bushy stunt of hawkweed oxtongue (<i>Picris hieracioides</i>) in south-eastern Serbia and the role of the leafhopper <i>Neoaliturus fenestratus</i> (Deltocephalinae) as a natural vector. <i>European Journal of Plant Pathology</i> , 2012, 134, 647-660.	1.7	25
41	<i>Oziella sibirica</i> (Acari: Eriophyoidea: Phytoptidae), a new eriophyoid mite species described using confocal microscopy, COI barcoding and 3D surface reconstruction. <i>Zootaxa</i> , 2012, 3560, 41.	0.5	37
42	Multigene sequence data and genetic diversity among <i>Candidatus</i> Phytoplasma ulmi™ strains infecting <i>Ulmus</i> spp. in Serbia. <i>Plant Pathology</i> , 2011, 60, 356-368.	2.4	22
43	<i>Tuta absoluta</i> (Meyrick, 1917) (Lepidoptera, Gelechiidae): A new pest of tomato in Serbia. <i>Pesticidi i Fitomedicina = Pesticides and Phytomedicine</i> , 2011, 26, 197-204.	0.2	7
44	Molecular characteristics of phytoplasmas associated with <i>Flavescence dorée</i> in clematis and grapevine and preliminary results on the role of <i>Dictyophara europaea</i> as a vector. <i>Plant Pathology</i> , 2009, 58, 826-837.	2.4	74
45	Occurrence and Distribution of Grapevine Yellows Caused by Stolbur Phytoplasma in Montenegro. <i>Journal of Phytopathology</i> , 2009, 157, 682-685.	1.0	14
46	Stolbur Phytoplasma Transmission to Maize by <i>Reptalus panzeri</i> and the Disease Cycle of Maize Redness in Serbia. <i>Phytopathology</i> , 2009, 99, 1053-1061.	2.2	44
47	First report of alder yellows phytoplasma on common alder (<i>Alnus glutinosa</i>) in Serbia. <i>Plant Pathology</i> , 2008, 57, 773-773.	2.4	8
48	<i>Candidatus</i> Phytoplasma ulmi™ causing yellows in <i>Zelkova serrata</i> newly reported in Italy. <i>Plant Pathology</i> , 2008, 57, 1174-1174.	2.4	4
49	New strain of <i>Candidatus</i> Phytoplasma ulmi™ infecting <i>Ulmus minor</i> and <i>U. laevis</i> in Serbia. <i>Plant Pathology</i> , 2008, 57, 1174-1174.	2.4	9
50	Roles of stolbur phytoplasma and <i>Reptalus panzeri</i> (Cixiinae, Auchenorrhyncha) in the epidemiology of Maize redness in Serbia. <i>European Journal of Plant Pathology</i> , 2007, 118, 85-89.	1.7	43
51	Framework for risk assessment of <i>Candidatus</i> Phytoplasma solani™ associated diseases outbreaks in agroecosystems in Serbia. <i>Journal of Plant Pathology</i> , 0, , 1.	1.2	3