

Raquel Gomez de Heras

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5194071/publications.pdf>

Version: 2024-02-01

25
papers

1,615
citations

516710

16
h-index

552781

26
g-index

27
all docs

27
docs citations

27
times ranked

1894
citing authors

#	ARTICLE	IF	CITATIONS
1	Abrupt cessation of reboxetine along alcohol deprivation results in alcohol intake escalation after reinstatement of drinking. <i>Addiction Biology</i> , 2021, 26, e12957.	2.6	3
2	Bupropion, a possible antidepressant without negative effects on alcohol relapse. <i>European Neuropsychopharmacology</i> , 2019, 29, 756-765.	0.7	2
3	Perinatal free choice of a high-calorie low-protein diet affects leptin signaling through IRS1 and AMPK dephosphorylation in the hypothalamus of female rat offspring in adulthood. <i>Acta Physiologica</i> , 2019, 226, e13244.	3.8	11
4	Oleylethanolamide prevents neuroimmune HMGB1/TLR4/NF- κ B danger signaling in rat frontal cortex and depressive-like behavior induced by ethanol binge administration. <i>Addiction Biology</i> , 2017, 22, 724-741.	2.6	88
5	A moderate diet restriction during pregnancy alters the levels of endocannabinoids and endocannabinoid-related lipids in the hypothalamus, hippocampus and olfactory bulb of rat offspring in a sex-specific manner. <i>PLoS ONE</i> , 2017, 12, e0174307.	2.5	15
6	Exposure to a Highly Caloric Palatable Diet During Pregestational and Gestational Periods Affects Hypothalamic and Hippocampal Endocannabinoid Levels at Birth and Induces Adiposity and Anxiety-Like Behaviors in Male Rat Offspring. <i>Frontiers in Behavioral Neuroscience</i> , 2016, 9, 339.	2.0	25
7	Maternal Caloric Restriction Implemented during the Preconceptional and Pregnancy Period Alters Hypothalamic and Hippocampal Endocannabinoid Levels at Birth and Induces Overweight and Increased Adiposity at Adulthood in Male Rat Offspring. <i>Frontiers in Behavioral Neuroscience</i> , 2016, 10, 208.	2.0	22
8	Long-Term Effects of Prenatal Exposure to Undernutrition on Cannabinoid Receptor-Related Behaviors: Sex and Tissue-Specific Alterations in the mRNA Expression of Cannabinoid Receptors and Lipid Metabolic Regulators. <i>Frontiers in Behavioral Neuroscience</i> , 2016, 10, 241.	2.0	20
9	Role of the satiety factor oleylethanolamide in alcoholism. <i>Addiction Biology</i> , 2016, 21, 859-872.	2.6	58
10	Exposure to a Highly Caloric Palatable Diet during the Perinatal Period Affects the Expression of the Endogenous Cannabinoid System in the Brain, Liver and Adipose Tissue of Adult Rat Offspring. <i>PLoS ONE</i> , 2016, 11, e0165432.	2.5	24
11	The administration of atomoxetine during alcohol deprivation induces a time-limited increase in alcohol consumption after relapse. <i>International Journal of Neuropsychopharmacology</i> , 2014, 17, 1905-1910.	2.1	8
12	Cannabinoid Receptors and Cholecystokinin in Feeding Inhibition. <i>Vitamins and Hormones</i> , 2013, 92, 165-196.	1.7	13
13	Increased alcohol consumption in rats after subchronic antidepressant treatment. <i>International Journal of Neuropsychopharmacology</i> , 2013, 16, 1809-1818.	2.1	17
14	Ghrelin-Induced Orexigenic Effect in Rats Depends on the Metabolic Status and Is Counteracted by Peripheral CB1 Receptor Antagonism. <i>PLoS ONE</i> , 2013, 8, e60918.	2.5	33
15	Effects of the anandamide uptake blocker AM404 on food intake depend on feeding status and route of administration. <i>Pharmacology Biochemistry and Behavior</i> , 2012, 101, 1-7.	2.9	17
16	Additive effects of cannabinoid CB1 receptors blockade and cholecystokinin on feeding inhibition. <i>Pharmacology Biochemistry and Behavior</i> , 2011, 98, 220-226.	2.9	18
17	Nicotine causes opposite effects on alcohol intake: Evidence in an animal experimental model of abstinence and relapse from alcohol. <i>Nicotine and Tobacco Research</i> , 2009, 11, 1304-1311.	2.6	12
18	Pretreatment with subeffective doses of Rimonabant attenuates orexigenic actions of orexin A-hypocretin 1. <i>Neuropharmacology</i> , 2008, 54, 219-225.	4.1	48

#	ARTICLE	IF	CITATIONS
19	Antiobesity effects of the novel in vivo neutral cannabinoid receptor antagonist 5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-3-hexyl-1H-1,2,4-triazole "LH 21". <i>Neuropharmacology</i> , 2006, 51, 358-366.	4.1	116
20	Nicotine in alcohol deprivation increases alcohol operant self-administration during reinstatement. <i>Neuropharmacology</i> , 2004, 47, 1036-1044.	4.1	39
21	Stimulation of voluntary ethanol intake by cannabinoid receptor agonists in ethanol-preferring sP rats. <i>Psychopharmacology</i> , 2002, 159, 181-187.	3.1	156
22	An anorexic lipid mediator regulated by feeding. <i>Nature</i> , 2001, 414, 209-212.	27.8	646
23	The cannabinoid receptor antagonist SR 141716 prevents acquisition of drinking behavior in alcohol-preferring rats. <i>European Journal of Pharmacology</i> , 2001, 430, 369-371.	3.5	71
24	Peripheral versus central effects of glucagon-like peptide-1 receptor agonists on satiety and body weight loss in Zucker obese rats. <i>Metabolism: Clinical and Experimental</i> , 2000, 49, 709-717.	3.4	144
25	Rethinking the Use of Antidepressants to Treat Alcohol Use Disorders and Depression Comorbidity: The Role of Neurogenesis. , 0, , .		0