

Hans-Georg Joost

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5200897/publications.pdf>

Version: 2024-02-01

223
papers

14,813
citations

18479

62
h-index

22161

113
g-index

240
all docs

240
docs citations

240
times ranked

17415
citing authors

#	ARTICLE	IF	CITATIONS
1	Salivary nitrate/nitrite and acetaldehyde in humans: potential combination effects in the upper gastrointestinal tract and possible consequences for the in vivo formation of N-nitroso compounds—a hypothesis. Archives of Toxicology, 2022, 96, 1905-1914.	4.2	5
2	Comparison of points of departure between subchronic and chronic toxicity studies on food additives, food contaminants and natural food constituents. Food and Chemical Toxicology, 2020, 146, 111784.	3.6	4
3	Identification of Novel Potential Type 2 Diabetes Genes Mediating β -Cell Loss and Hyperglycemia Using Positional Cloning. Frontiers in Genetics, 2020, 11, 567191.	2.3	5
4	Immunity-related GTPase induces lipophagy to prevent excess hepatic lipid accumulation. Journal of Hepatology, 2020, 73, 771-782.	3.7	34
5	Potential effects of reduced red meat compared with increased fiber intake on glucose metabolism and liver fat content: a randomized and controlled dietary intervention study. American Journal of Clinical Nutrition, 2019, 109, 288-296.	4.7	15
6	Increased Irf202b/IFI16 expression stimulates adipogenesis in mice and humans. Diabetologia, 2018, 61, 1167-1179.	6.3	21
7	Derivation and external validation of a clinical version of the German Diabetes Risk Score (GDRS) including measures of HbA1c. BMJ Open Diabetes Research and Care, 2018, 6, e000524.	2.8	8
8	A collective diabetes cross in combination with a computational framework to dissect the genetics of human obesity and Type 2 diabetes. Human Molecular Genetics, 2018, 27, 3099-3112.	2.9	21
9	The role of dual leucine zipper kinase (DLK) in β -cell apoptosis: a potential target for the prevention and treatment of type 2 diabetes?. Naunyn-Schmiedeberg's Archives of Pharmacology, 2017, 390, 767-768.	3.0	1
10	Early hypermethylation of hepatic <i>Igf2bp2</i> results in its reduced expression preceding fatty liver in mice. Human Molecular Genetics, 2016, 25, ddw121.	2.9	46
11	Identification of Four Mouse Diabetes Candidate Genes Altering β -Cell Proliferation. PLoS Genetics, 2015, 11, e1005506.	3.5	37
12	Amino acids, lipid metabolites, and ferritin as potential mediators linking red meat consumption to type 2 diabetes. American Journal of Clinical Nutrition, 2015, 101, 1241-1250.	4.7	95
13	Caloric restriction and intermittent fasting alter hepatic lipid droplet proteome and diacylglycerol species and prevent diabetes in NZO mice. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2015, 1851, 566-576.	2.4	98
14	GLP-1—oestrogen attenuates hyperphagia and protects from beta cell failure in diabetes-prone New Zealand obese (NZO) mice. Diabetologia, 2015, 58, 604-614.	6.3	32
15	The diabetes gene Zfp69 modulates hepatic insulin sensitivity in mice. Diabetologia, 2015, 58, 2403-2413.	6.3	20
16	Skeletal muscle mitochondrial uncoupling prevents diabetes but not obesity in NZO mice, a model for polygenic diabetes. Genes and Nutrition, 2015, 10, 57.	2.5	10
17	Nitrate and nitrite in the diet: How to assess their benefit and risk for human health. Molecular Nutrition and Food Research, 2015, 59, 106-128.	3.3	170
18	Deletion of Both Rab-GTPase—Activating Proteins TBC1D1 and TBC1D4 in Mice Eliminates Insulin- and AICAR-Stimulated Glucose Transport. Diabetes, 2015, 64, 746-759.	0.6	69

#	ARTICLE	IF	CITATIONS
19	Diabetes and cancer: Epidemiology and potential mechanisms. Diabetes and Vascular Disease Research, 2014, 11, 390-394.	2.0	44
20	Differential Transcriptome Analysis of Diabetes-Resistant and -Sensitive Mouse Islets Reveals Significant Overlap With Human Diabetes Susceptibility Genes. Diabetes, 2014, 63, 4230-4238.	0.6	40
21	The genetic basis of obesity-associated type 2 diabetes (diabesity) in polygenic mouse models. Mammalian Genome, 2014, 25, 401-412.	2.2	53
22	Update of the German Diabetes Risk Score and external validation in the German MONICA/KORA study. Diabetes Research and Clinical Practice, 2014, 104, 459-466.	2.8	48
23	Impact of the Adipokine Adiponectin and the Hepatokine Fetuin-A on the Development of Type 2 Diabetes: Prospective Cohort- and Cross-Sectional Phenotyping Studies. PLoS ONE, 2014, 9, e92238.	2.5	63
24	Microsomal triglyceride transfer protein -164 Tâ€™&â€™C gene polymorphism and risk of cardiovascular disease: results from the EPIC-Potsdam case-cohort study. BMC Medical Genetics, 2013, 14, 19.	2.1	6
25	Red meat and T2DMâ€™the difficult path to a proof of causality. Nature Reviews Endocrinology, 2013, 9, 509-511.	9.6	2
26	Consumption of red meat and whole-grain bread in relation to biomarkers of obesity, inflammation, glucose metabolism and oxidative stress. European Journal of Nutrition, 2013, 52, 337-345.	3.9	177
27	Identification of Serum Metabolites Associated With Risk of Type 2 Diabetes Using a Targeted Metabolomic Approach. Diabetes, 2013, 62, 639-648.	0.6	820
28	Assessing improvement in disease prediction using net reclassification improvement: impact of risk cut-offs and number of risk categories. European Journal of Epidemiology, 2013, 28, 25-33.	5.7	27
29	Conventional Knockout of Tbc1d1 in Mice Impairs Insulin- and AICAR-Stimulated Glucose Uptake in Skeletal Muscle. Endocrinology, 2013, 154, 3502-3514.	2.8	61
30	Estrogen Deficiency Aggravates Insulin Resistance and Induces Î²-Cell Loss and Diabetes in Female New Zealand Obese Mice. Hormone and Metabolic Research, 2013, 45, 430-435.	1.5	43
31	An Interval of the Obesity QTL Nob3.38 within a QTL Hotspot on Chromosome 1 Modulates Behavioral Phenotypes. PLoS ONE, 2013, 8, e53025.	2.5	8
32	The Value of Genetic Information for Diabetes Risk Prediction â€™ Differences According to Sex, Age, Family History and Obesity. PLoS ONE, 2013, 8, e64307.	2.5	33
33	Loss of function of Ifi202b by a microdeletion on chromosome 1 of C57BL/6J mice suppresses 11Î²-hydroxysteroid dehydrogenase type 1 expression and development of obesity. Human Molecular Genetics, 2012, 21, 3845-3857.	2.9	29
34	GTPase ARFRP1 Is Essential for Normal Hepatic Glycogen Storage and Insulin-Like Growth Factor 1 Secretion. Molecular and Cellular Biology, 2012, 32, 4363-4374.	2.3	24
35	The GTPase ARFRP1 controls the lipidation of chylomicrons in the Golgi of the intestinal epithelium. Human Molecular Genetics, 2012, 21, 3128-3142.	2.9	26
36	Pathophysiology and Genetics of Obesity and Diabetes in the New Zealand Obese Mouse: A Model of the Human Metabolic Syndrome. , 2012, 933, 59-73.		35

#	ARTICLE	IF	CITATIONS
37	Novel biomarkers for pre-diabetes identified by metabolomics. <i>Molecular Systems Biology</i> , 2012, 8, 615.	7.2	605
38	Body iron stores and risk of type 2 diabetes: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam study. <i>Diabetologia</i> , 2012, 55, 2613-2621.	6.3	102
39	Gamma-glutamyltransferase, cardiovascular disease and mortality in individuals with diabetes mellitus. <i>Diabetes/Metabolism Research and Reviews</i> , 2012, 28, 284-288.	4.0	21
40	The HPA axis modulates the CNS melanocortin control of liver triacylglyceride metabolism. <i>Physiology and Behavior</i> , 2012, 105, 791-799.	2.1	16
41	Heterogeneity of the Stearoyl-CoA desaturase-1 (SCD1) Gene and Metabolic Risk Factors in the EPIC-Potsdam Study. <i>PLoS ONE</i> , 2012, 7, e48338.	2.5	13
42	Erythrocyte membrane phospholipid fatty acids, desaturase activity, and dietary fatty acids in relation to risk of type 2 diabetes in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. <i>American Journal of Clinical Nutrition</i> , 2011, 93, 127-142.	4.7	218
43	Ghrelin-induced hypothermia: A physiological basis but no clinical risk. <i>Physiology and Behavior</i> , 2011, 105, 43-51.	2.1	18
44	Estimation of the contribution of biomarkers of different metabolic pathways to risk of type 2 diabetes. <i>European Journal of Epidemiology</i> , 2011, 26, 29-38.	5.7	41
45	Non-fasting lipids and risk of cardiovascular disease in patients with diabetes mellitus. <i>Diabetologia</i> , 2011, 54, 73-77.	6.3	28
46	Dissociation of lipotoxicity and glucotoxicity in a mouse model of obesity associated diabetes: role of forkhead box O1 (FOXO1) in glucose-induced beta cell failure. <i>Diabetologia</i> , 2011, 54, 605-616.	6.3	77
47	Role of Medium- and Short-Chain L-3-Hydroxyacyl-CoA Dehydrogenase in the Regulation of Body Weight and Thermogenesis. <i>Endocrinology</i> , 2011, 152, 4641-4651.	2.8	33
48	A Two-Step Association Study Identifies CAV2 rs2270188 Single Nucleotide Polymorphism Interaction with Fat Intake in Type 2 Diabetes Risk. <i>Journal of Nutrition</i> , 2011, 141, 177-181.	2.9	26
49	Diet Dependence of Diabetes in the New Zealand Obese (NZO) Mouse: Total Fat, But not Fat Quality or Sucrose Accelerates and Aggravates Diabetes. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2011, 119, 167-171.	1.2	19
50	The Genetic Basis of Obesity and Type 2 Diabetes: Lessons from the New Zealand Obese Mouse, a Polygenic Model of the Metabolic Syndrome. <i>Results and Problems in Cell Differentiation</i> , 2011, 52, 1-11.	0.7	19
51	Role of Zinc Finger Transcription Factor Zfp69 in Body Fat Storage and Diabetes Susceptibility of Mice. <i>Results and Problems in Cell Differentiation</i> , 2011, 52, 57-68.	0.7	12
52	Reliability of Serum Metabolite Concentrations over a 4-Month Period Using a Targeted Metabolomic Approach. <i>PLoS ONE</i> , 2011, 6, e21103.	2.5	131
53	Diet-induced gene expression of isolated pancreatic islets from a polygenic mouse model of the metabolic syndrome. <i>Diabetologia</i> , 2010, 53, 309-320.	6.3	44
54	Fasting plasma glucose and Type 2 diabetes risk: a non-linear relationship. <i>Diabetic Medicine</i> , 2010, 27, 473-476.	2.3	14

#	ARTICLE	IF	CITATIONS
55	The ARF-Like GTPase ARFRP1 Is Essential for Lipid Droplet Growth and Is Involved in the Regulation of Lipolysis. <i>Molecular and Cellular Biology</i> , 2010, 30, 1231-1242.	2.3	55
56	Altered GLUT4 trafficking in adipocytes in the absence of the GTPase Arfrp1. <i>Biochemical and Biophysical Research Communications</i> , 2010, 394, 896-903.	2.1	28
57	Positional Cloning of Zinc Finger Domain Transcription Factor Zfp69, a Candidate Gene for Obesity-Associated Diabetes Contributed by Mouse Locus Nidd/SJL. <i>PLoS Genetics</i> , 2009, 5, e1000541.	3.5	68
58	Association of <i>AHSG</i> Gene Polymorphisms With Fetuin-A Plasma Levels and Cardiovascular Diseases in the EPIC-Potsdam Study. <i>Circulation: Cardiovascular Genetics</i> , 2009, 2, 607-613.	5.1	83
59	Whole-grain consumption and transcription factor-7-like 2 (<i>TCF7L2</i>) rs7903146: gene×diet interaction in modulating type 2 diabetes risk. <i>British Journal of Nutrition</i> , 2009, 101, 478-481.	2.3	98
60	Medical Antihyperglycaemic Treatment of Type 2 Diabetes Mellitus. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2009, 117, 522-557.	1.2	84
61	Essential role of glucose transporter GLUT3 for post-implantation embryonic development. <i>Journal of Endocrinology</i> , 2009, 200, 23-33.	2.6	51
62	GLUT8, the enigmatic intracellular hexose transporter. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2009, 296, E614-E618.	3.5	95
63	A dual role of the N-terminal FQQI motif in GLUT4 trafficking. <i>Biological Chemistry</i> , 2009, 390, 883-92.	2.5	10
64	Association of a diabetes risk score with risk of myocardial infarction, stroke, specific types of cancer, and mortality: a prospective study in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam cohort. <i>European Journal of Epidemiology</i> , 2009, 24, 281-288.	5.7	49
65	GOAT links dietary lipids with the endocrine control of energy balance. <i>Nature Medicine</i> , 2009, 15, 741-745.	30.7	359
66	Association of the <i>FTO</i> rs9939609 Single Nucleotide Polymorphism With C-reactive Protein Levels. <i>Obesity</i> , 2009, 17, 330-334.	3.0	37
67	Lysosomal localization of GLUT8 in the testis – the EXXXLL motif of GLUT8 is sufficient for its intracellular sorting via AP1- and AP2-mediated interaction. <i>FEBS Journal</i> , 2009, 276, 3729-3743.	4.7	26
68	Monitoring detaching murals in the Convent of M ¹ /4stair (Switzerland) by optical metrology. <i>Journal of Cultural Heritage</i> , 2009, 10, 94-105.	3.3	24
69	Use of Multiple Metabolic and Genetic Markers to Improve the Prediction of Type 2 Diabetes: the EPIC-Potsdam Study. <i>Diabetes Care</i> , 2009, 32, 2116-2119.	8.6	125
70	Angiogenesis in the New Zealand obese mouse model fed with high fat diet. <i>Lipids in Health and Disease</i> , 2009, 8, 13.	3.0	16
71	Characterization of <i>Nob3</i> , a major quantitative trait locus for obesity and hyperglycemia on mouse chromosome 1. <i>Physiological Genomics</i> , 2009, 38, 226-232.	2.3	31
72	Genetische Disposition f ¹ /4r die Entwicklung von Adipositas und Typ-2-Diabetes im Mausmodell und beim Menschen. <i>Journal Fur Verbraucherschutz Und Lebensmittelsicherheit</i> , 2008, 3, 86-88.	1.4	1

#	ARTICLE	IF	CITATIONS
73	Impaired leptin activity in New Zealand Obese mice: model of angiogenesis. <i>Genes and Nutrition</i> , 2008, 3, 177-180.	2.5	5
74	Deletion of Glucose Transporter GLUT8 in Mice Increases Locomotor Activity. <i>Behavior Genetics</i> , 2008, 38, 396-406.	2.1	35
75	Metabolic syndrome and risk of incident diabetes: findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. <i>Cardiovascular Diabetology</i> , 2008, 7, 35.	6.8	80
76	Tbc1d1 mutation in lean mouse strain confers leanness and protects from diet-induced obesity. <i>Nature Genetics</i> , 2008, 40, 1354-1359.	21.4	174
77	Plasma Fetuin-A Levels and the Risk of Myocardial Infarction and Ischemic Stroke. <i>Circulation</i> , 2008, 118, 2555-2562.	1.6	277
78	Liver Enzymes and Incident Diabetes. <i>Diabetes Care</i> , 2008, 31, 1138-1143.	8.6	84
79	Pathogenesis, Risk Assessment and Prevention of Type 2 Diabetes mellitus. <i>Obesity Facts</i> , 2008, 1, 128-137.	3.4	24
80	Targeted disruption of <i>Slc2a8</i> (GLUT8) reduces motility and mitochondrial potential of spermatozoa. <i>Molecular Membrane Biology</i> , 2008, 25, 224-235.	2.0	40
81	Uncoupling protein 1 expression in murine skeletal muscle increases AMPK activation, glucose turnover, and insulin sensitivity in vivo. <i>Physiological Genomics</i> , 2008, 33, 333-340.	2.3	53
82	Simultaneous deletion of ghrelin and its receptor increases motor activity and energy expenditure. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 294, G610-G618.	3.4	153
83	Neuronal functions, feeding behavior, and energy balance in <i>Slc2a3</i> ^{+/-} mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2008, 295, E1084-E1094.	3.5	30
84	ADP-ribosylation Factor-like GTPase ARFRP1 Is Required for Trans-Golgi to Plasma Membrane Trafficking of E-cadherin. <i>Journal of Biological Chemistry</i> , 2008, 283, 27179-27188.	3.4	31
85	High-Fat, Carbohydrate-Free Diet Markedly Aggravates Obesity but Prevents β^2 -Cell Loss and Diabetes in the Obese, Diabetes-Susceptible <i>db/db</i> Strain. <i>Obesity Facts</i> , 2008, 1, 292-297.	3.4	12
86	A meta-analysis of QTL for diabetes-related traits in rodents. <i>Physiological Genomics</i> , 2008, 34, 42-53.	2.3	40
87	Plasma Fetuin-A Levels and the Risk of Type 2 Diabetes. <i>Diabetes</i> , 2008, 57, 2762-2767.	0.6	326
88	Ablation of the Cholesterol Transporter Adenosine Triphosphate-Binding Cassette Transporter G1 Reduces Adipose Cell Size and Protects against Diet-Induced Obesity. <i>Endocrinology</i> , 2007, 148, 1561-1573.	2.8	74
89	NO to Obesity: Does Nitric Oxide Regulate Fat Oxidation and Insulin Sensitivity?. <i>Endocrinology</i> , 2007, 148, 4545-4547.	2.8	16
90	Personalised nutrition: status and perspectives. <i>British Journal of Nutrition</i> , 2007, 98, 26-31.	2.3	72

#	ARTICLE	IF	CITATIONS
91	An Accurate Risk Score Based on Anthropometric, Dietary, and Lifestyle Factors to Predict the Development of Type 2 Diabetes. <i>Diabetes Care</i> , 2007, 30, 510-515.	8.6	341
92	Effects of Obestatin on Energy Balance and Growth Hormone Secretion in Rodents. <i>Endocrinology</i> , 2007, 148, 21-26.	2.8	228
93	A meta-analysis of quantitative trait loci associated with body weight and adiposity in mice. <i>International Journal of Obesity</i> , 2007, 31, 829-841.	3.4	78
94	Prevention of complex diseases by genotype-based nutrition: realistic concept or fiction?. <i>Journal of Molecular Medicine</i> , 2007, 85, 103-105.	3.9	1
95	Development of diabetes in obese, insulin-resistant mice: essential role of dietary carbohydrate in beta cell destruction. <i>Diabetologia</i> , 2007, 50, 1481-1489.	6.3	51
96	Variation in the HHEX/IDE gene region predisposes to type 2 diabetes in the prospective, population-based EPIC-Potsdam cohort. <i>Diabetologia</i> , 2007, 50, 2405-2407.	6.3	24
97	The central melanocortin system directly controls peripheral lipid metabolism. <i>Journal of Clinical Investigation</i> , 2007, 117, 3475-3488.	8.2	341
98	Personalized Prevention of Type 2 Diabetes. , 2007, , 61-74.		0
99	Hyperphagia, lower body temperature, and reduced running wheel activity precede development of morbid obesity in New Zealand obese mice. <i>Physiological Genomics</i> , 2006, 25, 234-241.	2.3	80
100	Gene Variants and Obesity. , 2006, , 266-299.		0
101	Knockout of Arfrp1 leads to disruption of ARF-like1 (ARL1) targeting to the trans-Golgi in mouse embryos and HeLa cells. <i>Molecular Membrane Biology</i> , 2006, 23, 475-485.	2.0	51
102	Endocytosis of the glucose transporter GLUT8 is mediated by interaction of a dileucine motif with the β 2-adaptin subunit of the AP-2 adaptor complex. <i>Journal of Cell Science</i> , 2006, 119, 2321-2331.	2.0	36
103	PYY3-36 as an anti-obesity drug target. <i>Obesity Reviews</i> , 2005, 6, 307-322.	6.5	109
104	Consuming Fructose-sweetened Beverages Increases Body Adiposity in Mice. <i>Obesity</i> , 2005, 13, 1146-1156.	4.0	255
105	Mice without the Regulator Gene Rsc1A1 Exhibit Increased Na ⁺ - d -Glucose Cotransport in Small Intestine and Develop Obesity. <i>Molecular and Cellular Biology</i> , 2005, 25, 78-87.	2.3	49
106	The case for strategic international alliances to harness nutritional genomics for public and personal health. <i>British Journal of Nutrition</i> , 2005, 94, 623-632.	2.3	137
107	Characterization of the human SLC2A11 (GLUT11) gene: alternative promoter usage, function, expression, and subcellular distribution of three isoforms, and lack of mouse orthologue. <i>Molecular Membrane Biology</i> , 2005, 22, 339-351.	2.0	63
108	Nutrition-/diet-induced changes in gene expression in white adipose tissue. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2005, 19, 589-603.	4.7	63

#	ARTICLE	IF	CITATIONS
109	Differential hepatic gene expression in a polygenic mouse model with insulin resistance and hyperglycemia: evidence for a combined transcriptional dysregulation of gluconeogenesis and fatty acid synthesis. <i>Journal of Molecular Endocrinology</i> , 2004, 32, 195-208.	2.5	28
110	Pleiotropy of leptin receptor signalling is defined by distinct roles of the intracellular tyrosines. <i>FEBS Journal</i> , 2004, 272, 109-119.	4.7	93
111	Does gut hormone PYY36 decrease food intake in rodents?. <i>Nature</i> , 2004, 430, 1-3.	27.8	207
112	GLUT11, but not GLUT8 or GLUT12, is expressed in human skeletal muscle in a fibre type-specific pattern. <i>Pflügers Archiv European Journal of Physiology</i> , 2004, 448, 105-113.	2.8	25
113	The glucose transporter families SGLT and GLUT: molecular basis of normal and aberrant function. <i>Journal of Parenteral and Enteral Nutrition</i> , 2004, 28, 364-371.	2.6	370
114	Cellular models for the analysis of signaling by protein kinase B and the forkhead transcription factor FKHR (Foxo1a). <i>Regulatory Peptides</i> , 2004, 121, 19-24.	1.9	3
115	Arf-Like Proteins. , 2004, , 325-350.		0
116	DYRK1 is a co-activator of FKHR (FOXO1a)-dependent glucose-6-phosphatase gene expression. <i>Biochemical and Biophysical Research Communications</i> , 2003, 300, 764-769.	2.1	45
117	Unusual function of the activation loop in the protein kinase DYRK1A. <i>Biochemical and Biophysical Research Communications</i> , 2003, 302, 403-408.	2.1	39
118	Diet-dependent obesity and hypercholesterolemia in the New Zealand obese mouse: identification of a quantitative trait locus for elevated serum cholesterol on the distal mouse chromosome 5. <i>Biochemical and Biophysical Research Communications</i> , 2003, 304, 812-817.	2.1	33
119	Akt Modulates STAT3-mediated Gene Expression through a FKHR (FOXO1a)-dependent Mechanism. <i>Journal of Biological Chemistry</i> , 2003, 278, 5242-5249.	3.4	68
120	Alternative splicing variants of dual specificity tyrosine phosphorylated and regulated kinase 1B exhibit distinct patterns of expression and functional properties. <i>Biochemical Journal</i> , 2003, 372, 881-888.	3.7	47
121	Effect of Hyperinsulinemia and Type 2 Diabetes-Like Hyperglycemia on Expression of Hepatic Cytochrome P450 and GlutathioneS-Transferase Isoforms in a New Zealand Obese-Derived Mouse Backcross Population. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2002, 302, 442-450.	2.5	35
122	Reduced Sperm Count and Normal Fertility in Male Mice with Targeted Disruption of the ADP-Ribosylation Factor-Like 4 (Arl4) Gene. <i>Molecular and Cellular Biology</i> , 2002, 22, 2761-2768.	2.3	60
123	Regulation of the Forkhead Transcription Factor FKHR (FOXO1a) by Glucose Starvation and AICAR, an Activator of AMP-Activated Protein Kinase. <i>Endocrinology</i> , 2002, 143, 3183-3186.	2.8	74
124	Identification of the Critical Sequence Elements in the Cytoplasmic Domain of Leptin Receptor Isoforms Required for Janus Kinase/Signal Transducer and Activator of Transcription Activation by Receptor Heterodimers. <i>Molecular Endocrinology</i> , 2002, 16, 859-872.	3.7	90
125	Construction And Characterization of a Conditionally Active Construct of The Insulin-Regulated Forkhead Transcription Factor FKHR. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2002, 110, 304-309.	1.2	6
126	Embryonic Lethality Caused by Apoptosis during Gastrulation in Mice Lacking the Gene of the ADP-Ribosylation Factor-Related Protein 1. <i>Molecular and Cellular Biology</i> , 2002, 22, 1488-1494.	2.3	44

#	ARTICLE	IF	CITATIONS
127	Mouse ARF-Related Protein 1: Genomic Organization and Analysis of Its Promoter. Biochemical and Biophysical Research Communications, 2002, 292, 113-120.	2.1	14
128	Nomenclature of the GLUT/SLC2A family of sugar/polyol transport facilitators. American Journal of Physiology - Endocrinology and Metabolism, 2002, 282, E974-E976.	3.5	325
129	Inhibition of the renin-angiotensin system ameliorates genetically determined hyperinsulinemia. European Journal of Pharmacology, 2002, 436, 145-150.	3.5	25
130	Characterisation of the mouse diabetes susceptibility locus Nidd/SJL: islet cell destruction, interaction with the obesity QTL Nob1, and effect of dietary fat. Diabetologia, 2002, 45, 823-830.	6.3	56
131	Concentration-dependent stimulatory and inhibitory effect of troglitazone on insulin-induced fatty acid synthase expression and protein kinase B activity in 3T3-L1 adipocytes. Naunyn-Schmiedeberg's Archives of Pharmacology, 2002, 365, 290-295.	3.0	5
132	The glucose transport facilitator GLUT8 is predominantly associated with the acrosomal region of mature spermatozoa. Cell and Tissue Research, 2002, 307, 237-242.	2.9	73
133	Regulation of the Forkhead Transcription Factor FKHR (FOXO1a) by Glucose Starvation and AICAR, an Activator of AMP-Activated Protein Kinase. Endocrinology, 2002, 143, 3183-3183.	2.8	12
134	Identification of the Critical Sequence Elements in the Cytoplasmic Domain of Leptin Receptor Isoforms Required for Janus Kinase/Signal Transducer and Activator of Transcription Activation by Receptor Heterodimers. Molecular Endocrinology, 2002, 16, 859-872.	3.7	20
135	The extended GLUT-family of sugar/polyol transport facilitators: nomenclature, sequence characteristics, and potential function of its novel members. Molecular Membrane Biology, 2001, 18, 247-256.	2.0	583
136	Differential Regulation of Endogenous Glucose-6-Phosphatase and Phosphoenolpyruvate Carboxykinase Gene Expression by the Forkhead Transcription Factor FKHR in H4IIE-Hepatoma Cells. Biochemical and Biophysical Research Communications, 2001, 285, 897-902.	2.1	97
137	Mouse GLUT8: Genomic Organization and Regulation of Expression in 3T3-L1 Adipocytes by Glucose. Biochemical and Biophysical Research Communications, 2001, 288, 969-974.	2.1	35
138	PSCA expression is regulated by phorbol ester and cell adhesion in the bladder carcinoma cell line RT112. Cancer Letters, 2001, 168, 37-43.	7.2	20
139	Targeting of GLUT6 (formerly GLUT9) and GLUT8 in rat adipose cells. Biochemical Journal, 2001, 358, 517.	3.7	64
140	Characterization of human glucose transporter (GLUT) 11 (encoded by SLC2A11), a novel sugar-transport facilitator specifically expressed in heart and skeletal muscle. Biochemical Journal, 2001, 359, 443.	3.7	59
141	Identification of the autophosphorylation sites and characterization of their effects in the protein kinase DYRK1A. Biochemical Journal, 2001, 359, 497.	3.7	115
142	Phorbol ester-induced activation of mitogen-activated protein kinase/extracellular-signal-regulated kinase kinase and extracellular-signal-regulated protein kinase decreases glucose-6-phosphatase gene expression. Biochemical Journal, 2001, 357, 867-873.	3.7	17
143	Characterization of human glucose transporter (GLUT) 11 (encoded by SLC2A11), a novel sugar-transport facilitator specifically expressed in heart and skeletal muscle. Biochemical Journal, 2001, 359, 443-449.	3.7	110
144	Phorbol ester-induced activation of mitogen-activated protein kinase/extracellular-signal-regulated kinase kinase and extracellular-signal-regulated protein kinase decreases glucose-6-phosphatase gene expression. Biochemical Journal, 2001, 357, 867.	3.7	12

#	ARTICLE	IF	CITATIONS
145	Targeting of GLUT6 (formerly GLUT9) and GLUT8 in rat adipose cells. <i>Biochemical Journal</i> , 2001, 358, 517-522.	3.7	95
146	The vitamin D receptor gene variant is associated with the prevalence of Type 2 diabetes mellitus and coronary artery disease. <i>Diabetic Medicine</i> , 2001, 18, 842-845.	2.3	94
147	Increased Hypothalamic Expression of the p75 Tumor Necrosis Factor Receptor in New Zealand Obese Mice. <i>Hormone and Metabolic Research</i> , 2001, 33, 520-524.	1.5	13
148	Activity and genomic organization of human glucose transporter 9 (GLUT9), a novel member of the family of sugar-transport facilitators predominantly expressed in brain and leucocytes. <i>Biochemical Journal</i> , 2000, 350, 771.	3.7	35
149	Activity and genomic organization of human glucose transporter 9 (GLUT9), a novel member of the family of sugar-transport facilitators predominantly expressed in brain and leucocytes. <i>Biochemical Journal</i> , 2000, 350, 771-776.	3.7	128
150	Glucose deprivation induces Akt-dependent synthesis and incorporation of GLUT1, but not of GLUT4, into the plasma membrane of 3T3-L1 adipocytes. <i>European Journal of Cell Biology</i> , 2000, 79, 943-949.	3.6	28
151	Quantitative trait loci for obesity and insulin resistance (Nob1, Nob2) and their interaction with the leptin receptor allele (Lepr A720T/T1044I) in New Zealand obese mice. <i>Diabetologia</i> , 2000, 43, 1565-1572.	6.3	69
152	GLUT8, a Novel Member of the Sugar Transport Facilitator Family with Glucose Transport Activity. <i>Journal of Biological Chemistry</i> , 2000, 275, 16275-16280.	3.4	211
153	Specificity Determinants of Substrate Recognition by the Protein Kinase DYRK1A. <i>Journal of Biological Chemistry</i> , 2000, 275, 2431-2438.	3.4	219
154	Type 2 diabetes-like hyperglycemia in a backcross model of NZO and SJL mice: characterization of a susceptibility locus on chromosome 4 and its relation with obesity. <i>Diabetes</i> , 2000, 49, 1590-1596.	0.6	57
155	Reduced Expression of PSCA, a Member of the LY-6 Family of Cell Surface Antigens, in Bladder, Esophagus, and Stomach Tumors. <i>Biochemical and Biophysical Research Communications</i> , 2000, 275, 783-788.	2.1	107
156	Association of Impaired Phosphatidylinositol 3-Kinase Activity in GLUT1-Containing Vesicles with Malinsertion of Glucose Transporters into the Plasma Membrane of Fibroblasts from a Patient with Severe Insulin Resistance and Clinical Features of Werner Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2000, 85, 905-918.	3.6	2
157	The ADP-ribosylation Factor (ARF)-related GTPase ARF-related Protein Binds to the ARF-specific Guanine Nucleotide Exchange Factor Cytohesin and Inhibits the ARF-dependent Activation of Phospholipase D. <i>Journal of Biological Chemistry</i> , 1999, 274, 9744-9751.	3.4	41
158	Preparation of Recombinant Histone H3 as a Substrate for Protein Kinase Assays. <i>Analytical Biochemistry</i> , 1999, 274, 138-141.	2.4	3
159	ADP-ribosylation factor (ARF)-like 4, 6, and 7 represent a subgroup of the ARF family characterized by rapid nucleotide exchange and a nuclear localization signal. <i>FEBS Letters</i> , 1999, 456, 384-388.	2.8	63
160	Cloning and Characterization of DYRK1B, a Novel Member of the DYRK Family of Protein Kinases. <i>Biochemical and Biophysical Research Communications</i> , 1999, 254, 474-479.	2.1	70
161	Inhibition of insulin-stimulated glucose transport in 3T3-L1 cells by <i>Clostridium difficile</i> toxin B, <i>Clostridium sordellii</i> lethal toxin, and <i>Clostridium botulinum</i> C2 toxin. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1998, 357, 385-392.	3.0	13
162	Sequence Characteristics, Subcellular Localization, and Substrate Specificity of DYRK-related Kinases, a Novel Family of Dual Specificity Protein Kinases. <i>Journal of Biological Chemistry</i> , 1998, 273, 25893-25902.	3.4	258

#	ARTICLE	IF	CITATIONS
163	Hyperleptinemia and leptin receptor variant Asp600Asn in the obese, hyperinsulinemic KK mouse strain. <i>Journal of Molecular Endocrinology</i> , 1998, 21, 337-345.	2.5	19
164	Structural and Functional Characteristics of Dyrk, a Novel Subfamily of Protein Kinases with Dual Specificity. <i>Progress in Molecular Biology and Translational Science</i> , 1998, 62, 1-17.	1.9	181
165	Serine-294 and threonine-295 in the exofacial loop domain between helices 7 and 8 of glucose transporters (GLUT) are involved in the conformational alterations during the transport process. <i>Biochemical Journal</i> , 1998, 329, 289-293.	3.7	40
166	The mouse ADP-ribosylation factor-like 4 gene: two separate promoters direct specific transcription in tissues and testicular germ cell. <i>Biochemical Journal</i> , 1998, 335, 259-265.	3.7	19
167	Characterization of GLUT5 Domains Responsible for Fructose Transport. <i>Endocrinology</i> , 1998, 139, 827-831.	2.8	4
168	Chronic Effects of a Nonpeptide Corticotropin-Releasing Hormone Type I Receptor Antagonist on Pituitary-Adrenal Function, Body Weight, and Metabolic Regulation. <i>Endocrinology</i> , 1998, 139, 1546-1555.	2.8	37
169	Hyperleptinemia, Leptin Resistance, and Polymorphic Leptin Receptor in the New Zealand Obese Mouse*. <i>Endocrinology</i> , 1997, 138, 4234-4239.	2.8	83
170	Role of Conserved Arginine and Glutamate Residues on the Cytosolic Surface of Glucose Transporters for Transporter Function. <i>Biochemistry</i> , 1997, 36, 12897-12902.	2.5	74
171	Cloning of two splicing variants of the novel Ras-related GTPase Rab29 which is predominately expressed in kidney. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1997, 1352, 48-55.	2.4	5
172	Hyperleptinemia, Leptin Resistance, and Polymorphic Leptin Receptor in the New Zealand Obese Mouse. <i>Endocrinology</i> , 1997, 138, 4234-4239.	2.8	30
173	cDNA cloning and characterization of rat Clk3, a LAMMER kinase predominately expressed in testis. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1996, 1312, 63-67.	4.1	17
174	Distribution of the mRNA for protein phosphatase T in rat brain. <i>Molecular Brain Research</i> , 1996, 36, 23-28.	2.3	13
175	Glucose transport activity and ligand binding (cytochalasin B, IAPS-forskolin) of chimeric constructs of GLUT2 and GLUT4 expressed in COS-7-cells. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1996, 1284, 56-62.	2.6	7
176	Cloning of a novel member (ARL5) of the ARF-family of Ras-related GTPases. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1996, 1308, 1-6.	2.4	15
177		6.3	45
178	Comparison of the effects of insulin, PDGF, interleukin-6, and interferon-g on glucose transport in 3T3-L1 cells: lack of cross-talk between tyrosine kinase receptors and JAK/STAT pathways. <i>Diabetologia</i> , 1996, 39, 1432-1439.	6.3	30
179	Molecular Cloning and Characterization of a Novel Mammalian Protein Kinase Harboring a Homology Domain that Defines a Subfamily of Serine/Threonine Kinases. <i>FEBS Journal</i> , 1996, 235, 736-743.	0.2	35
180	Alternative mRNA Splicing of the Novel GTPase Rab28 Generates Isoforms with Different C-Termini. <i>FEBS Journal</i> , 1996, 237, 833-840.	0.2	29

#	ARTICLE	IF	CITATIONS
181	Apparatefestigkeit - eine bestimmende Größe im Sicherheitskonzept von Chemieanlagen. Chemie-Ingenieur-Technik, 1996, 68, 1442-1447.	0.8	3
182	Dyrk, a Dual Specificity Protein Kinase with Unique Structural Features Whose Activity Is Dependent on Tyrosine Residues between Subdomains VII and VIII. Journal of Biological Chemistry, 1996, 271, 3488-3495.	3.4	231
183	Evidence that Reduced Leptin Levels, but Not an Aberrant Sequence of Leptin or its Receptor, Contribute to the Obesity Syndrome in NON Mice. Hormone and Metabolic Research, 1996, 28, 669-673.	1.5	8
184	Difference in Leptin mRNA Levels Between Omental and Subcutaneous Abdominal Adipose Tissue From Obese Humans. Hormone and Metabolic Research, 1996, 28, 690-693.	1.5	216
185	Structural and functional heterogeneity of insulin receptors. Cellular Signalling, 1995, 7, 85-91.	3.6	30
186	Cloning of a Novel Family of Mammalian GTP-binding Proteins (RagA, RagBs, RagB1) with Remote Similarity to the Ras-related GTPases. Journal of Biological Chemistry, 1995, 270, 28982-28988.	3.4	106
187	ARP Is a Plasma Membrane-associated Ras-related GTPase with Remote Similarity to the Family of ADP-ribosylation Factors. Journal of Biological Chemistry, 1995, 270, 30657-30663.	3.4	41
188	Dissociation of the effects of training on oxidative metabolism, glucose utilisation and GLUT4 levels in skeletal muscle of streptozotocin-diabetic rats. Pflügers Archiv European Journal of Physiology, 1994, 427, 444-449.	2.8	15
189	In vivo glucose uptake and glucose transporter proteins GLUT1 and GLUT4 in heart and various types of skeletal muscle from streptozotocin-diabetic rats. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 1994, 1225, 275-282.	3.8	59
190	Substitution of conserved tyrosine residues in helix 4 (Y143) and 7 (Y293) affects the activity, but not IAPS-forskolin binding, of the glucose transporter GLUT4. FEBS Letters, 1994, 348, 114-118.	2.8	24
191	Abundance and subcellular distribution of GTP-binding proteins in 3T3-L1 cells before and after differentiation to the insulin-sensitive phenotype. FEBS Journal, 1993, 215, 611-617.	0.2	7
192	In vivo glucose uptake and glucose transporter proteins GLUT1 and GLUT3 in brain tissue from streptozotocin-diabetic rats. Acta Physiologica Scandinavica, 1993, 149, 221-225.	2.2	26
193	Biphasic Alteration of Glucose Transport in 3T3-L1 Cells During Differentiation to the Adipocyte-Like Phenotype. Hormone and Metabolic Research, 1993, 25, 71-76.	1.5	12
194	Localization of the binding domain of the inhibitory ligand forskolin in the glucose transporter GLUT-4 by photolabeling, proteolytic cleavage and a site-specific antiserum. Biochimica Et Biophysica Acta - Biomembranes, 1992, 1111, 178-184.	2.6	11
195	Subcellular distribution and activity of glucose transporter isoforms GLUT1 and GLUT4 transiently expressed in COS-7 cells. Biochimica Et Biophysica Acta Gene Regulatory Mechanisms, 1992, 1131, 245-252.	2.4	33
196	Subcellular distribution and phosphorylation state of insulin receptors from insulin- and isoproterenol- treated rat adipose cells. Cellular Signalling, 1991, 3, 51-58.	3.6	11
197	The signaling potential of the receptors for insulin and insulin-like growth factor I (IGF-I) in 3T3-L1 adipocytes: Comparison of glucose transport activity, induction of oncogene-c-fos, glucose transporter mRNA, and DNA-synthesis. Journal of Cellular Physiology, 1991, 149, 428-435.	4.1	56
198	Antagonistic effects of a covalently dimerized insulin derivative on insulin receptors in 3T3-L1 adipocytes.. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 1154-1158.	7.1	21

#	ARTICLE	IF	CITATIONS
199	The regulation of glucose transport in insulin-sensitive cells. <i>Diabetologia</i> , 1989, 32, 831-838.	6.3	32
200	Differential sensitivity to guanine nucleotides of basal and insulin-stimulated glucose transporter activity reconstituted from adipocyte membrane fractions. <i>FEBS Letters</i> , 1989, 255, 259-264.	2.8	34
201	Quantitative dissociation of glucose transport stimulation and insulin receptor tyrosine kinase activation in isolated adipocytes with a covalent insulin dimer (B29,B29 α^2 -sunberoyl-insulin). <i>Biochemical Pharmacology</i> , 1989, 38, 2269-2277.	4.4	6
202	Adipocyte plasma membranes contain two Gisubtypes but are devoid of Go. <i>FEBS Letters</i> , 1988, 238, 191-196.	2.8	46
203	Inhibition of insulin-stimulated glucose transport in rat adipocytes by nucleoside transport inhibitors. <i>FEBS Letters</i> , 1988, 227, 215-219.	2.8	14
204	Prostaglandin E2 differentiates between two forms of glucose transport inhibition by lipolytic agents. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1987, 336, 105-10.	3.0	7
205	The inhibition of rat adipocyte ecto-5 α^2 -nucleotidase by xanthines is not related to lipolysis. <i>Biochemical Pharmacology</i> , 1986, 35, 867-869.	4.4	1
206	Modulation of glucose transport in hamster adipocytes by insulin and by β - and α 2-adrenoceptor agonists. <i>Diabetologia</i> , 1986, 29, 371-377.	6.3	8
207	Extrapaneatic effects of hypoglycemic sulfonylureas: still a controversial issue. <i>Trends in Pharmacological Sciences</i> , 1985, 6, 239-241.	8.7	10
208	Insulin-like stimulation of glucose transport in isolated adipocytes by fatty acids. <i>Biochemical and Biophysical Research Communications</i> , 1985, 128, 1358-1363.	2.1	26
209	Dual effect of isoprenaline on glucose transport and response to insulin in isolated adipocytes. <i>Biochemical Pharmacology</i> , 1985, 34, 649-653.	4.4	20
210	Effects of theophylline on insulin receptors and insulin action in the adipocyte. <i>Molecular and Cellular Biochemistry</i> , 1983, 57, 177-183.	3.1	19
211	In vitro effects of theophylline on insulin receptors in adipocytes: Correlation with the lipolytic action of the agent. <i>Biochemical and Biophysical Research Communications</i> , 1982, 104, 45-51.	2.1	12
212	Effects of tolbutamide on insulin binding to isolated fat cells of the rat. <i>Biochemical Pharmacology</i> , 1982, 31, 1227-1231.	4.4	25
213	On the relation of glucose and insulin secretion in the fasting state. <i>Metabolism: Clinical and Experimental</i> , 1980, 29, 23-27.	3.4	19
214	Insulin release by tolbutamide and glibenclamide. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1979, 306, 185-188.	3.0	17
215	Subcellular B cell calcium and insulin secretion in vitro. <i>Virchows Archiv A, Pathological Anatomy and Histology</i> , 1978, 379, 203-217.	1.3	4
216	Uptake of tolbutamide by islets of Langerhans and other tissues. <i>Experientia</i> , 1978, 34, 1372-1373.	1.2	4

#	ARTICLE	IF	CITATIONS
217	Effects of polymer-linked sulfonylurea derivatives on insulin release. Naunyn-Schmiedeberg's Archives of Pharmacology, 1977, 297, 81-84.	3.0	7
218	Inhibition of insulin and glucagon release from the perfused rat pancreas by cyproheptadine (Periactinoli $\frac{1}{2}$, Nuran $\frac{1}{2}$). Diabetologia, 1976, 12, 201-206.	6.3	20
219	The inhibition of insulin secretion from the perfused rat pancreas after thyroxine treatment. Diabetologia, 1976, 12, 495-500.	6.3	16
220	Effects of bicyclic leucine analogues on insulin release from the perfused rat pancreas and fluorescence of reduced pyridine nucleotides from perfused pancreatic islets. Life Sciences, 1975, 16, 247-253.	4.3	5
221	Inhibition of insulin release from the rat pancreas by cyproheptadine and tricyclic antidepressants. Naunyn-Schmiedeberg's Archives of Pharmacology, 1974, 285, 99-102.	3.0	16
222	GLUT8. The AFCS-nature Molecule Pages, 0, , .	0.2	0
223	Arfrp1. The AFCS-nature Molecule Pages, 0, , .	0.2	0