Patrik Midlöv

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5206251/publications.pdf

Version: 2024-02-01

		126907	2	14800
113	2,914	33		47
papers	citations	h-index		g-index
119	119	119		3853
all docs	docs citations	times ranked		citing authors

#	Article	IF	CITATIONS
1	Health care utilization following "digi-physical―assessment compared to physical assessment for infectious symptoms in primary care. , 2022, 23, 4.		4
2	Disparities in prevalence of heart failure according to age, multimorbidity level and socioeconomic status in southern Sweden: a cross-sectional study. BMJ Open, 2022, 12, e051997.	1.9	5
3	Low adherence to statin treatment during the 1st year after an acute myocardial infarction is associated with increased 2nd-year mortality risk—an inverse probability of treatment weighted study on 54 872 patients. European Heart Journal - Cardiovascular Pharmacotherapy, 2021, 7, 141-147.	3.0	26
4	Antibiotic Prescription Rates After eVisits Versus Office Visits in Primary Care: Observational Study. JMIR Medical Informatics, 2021, 9, e25473.	2.6	10
5	Medication-related hospital readmissions within 30 days of discharge—A retrospective study of risk factors in older adults. PLoS ONE, 2021, 16, e0253024.	2.5	14
6	Patients and Professionals as Partners in Hypertension Care: Qualitative Substudy of a Randomized Controlled Trial Using an Interactive Web-Based System Via Mobile Phone. Journal of Medical Internet Research, 2021, 23, e26143.	4.3	6
7	Lifetime Risk of Visual Impairment Resulting from Glaucoma in Patients Initially Followed up for Elevated Intraocular Pressure. Ophthalmology Glaucoma, 2020, 3, 60-65.	1.9	2
8	Medication Discrepancies in Discharge Summaries and Associated Risk Factors for Elderly Patients with Many Drugs. Drugs - Real World Outcomes, 2020, 7, 53-62.	1.6	24
9	PERson-centredness in hypertension management using information technology (PERHIT): a protocol for a randomised controlled trial in primary health care. Blood Pressure, 2020, 29, 149-156.	1.5	9
10	Risk factors for hospital readmission in older adults within 30 days of discharge – a comparative retrospective study. BMC Geriatrics, 2020, 20, 467.	2.7	50
11	Text message-based lifestyle intervention in primary care patients with hypertension: a randomized controlled pilot trial. Scandinavian Journal of Primary Health Care, 2020, 38, 300-307.	1.5	11
12	Experiences of digital communication with automated patient interviews and asynchronous chat in Swedish primary care: a qualitative study. BMJ Open, 2020, 10, e036585.	1.9	20
13	Use of potentially inappropriate medication and polypharmacy in older adults: a repeated cross-sectional study. BMC Geriatrics, 2020, 20, 73.	2.7	29
14	Human- Versus Machine Learning–Based Triage Using Digitalized Patient Histories in Primary Care: Comparative Study. JMIR Medical Informatics, 2020, 8, e18930.	2.6	9
15	Health Care Staff's Experiences of Engagement When Introducing a Digital Decision Support System for Wound Management: Qualitative Study. JMIR Human Factors, 2020, 7, e23188.	2.0	4
16	Primary care doctor fostering and clinical research training in Sweden: Implications for Japan. Journal of General and Family Medicine, 2019, 20, 4-8.	0.8	6
17	Importance of potentially inappropriate medications, number of chronic conditions and medications for the risk of hospitalisation in elderly in Sweden: a case–control study. BMJ Open, 2019, 9, e029477.	1.9	19
18	Use of non-benzodiazepine hypnotics is associated with falls in nursing home residents: a longitudinal cohort study. Aging Clinical and Experimental Research, 2019, 31, 1087-1095.	2.9	10

#	Article	IF	CITATIONS
19	The use of case management for communityâ€dwelling older people: the effects on loneliness, symptoms of depression and life satisfaction in a randomised controlled trial. Scandinavian Journal of Caring Sciences, 2018, 32, 889-901.	2.1	66
20	Japan can learn from the Swedish primary care doctor fostering system. Journal of General and Family Medicine, 2018, 19, 183-184.	0.8	3
21	Active listing and more consultations in primary care are associated with shorter mean hospitalisation and interacting with psychiatric disorders when adjusting for multimorbidity, age and sex. Scandinavian Journal of Primary Health Care, 2018, 36, 308-316.	1.5	2
22	Elderly at risk in care transitions When discharge summaries are poorly transferred and used –a descriptive study. BMC Health Services Research, 2018, 18, 770.	2.2	19
23	Drugs, distrust and dialogue –a focus group study with Swedish GPs on discharge summary use in primary care. BMC Family Practice, 2018, 19, 127.	2.9	9
24	Sociodemographic Characteristics and Health Profile of the Elderly Seeking Health Care in Kampala, Uganda. Current Gerontology and Geriatrics Research, 2018, 2018, 1-9.	1.6	11
25	Women's Health in the Lund Area (WHILA) study. Health problems and acute myocardial infarction in women – A 17-year follow-up study. Maturitas, 2018, 115, 45-50.	2.4	10
26	Active listing and more consultations in primary care are associated with reduced hospitalisation in a Swedish population. BMC Health Services Research, 2018, 18, 101.	2.2	5
27	Effects of medication reviews on use of potentially inappropriate medications in elderly patients; a cross-sectional study in Swedish primary care. BMC Health Services Research, 2018, 18, 616.	2.2	41
28	Fall-risk-increasing drugs and falls requiring health care among older people with intellectual disability in comparison with the general population: A register study. PLoS ONE, 2018, 13, e0199218.	2.5	11
29	Risk factors and consequences of decreased kidney function in nursing home residents: A longitudinal study. Geriatrics and Gerontology International, 2017, 17, 791-797.	1.5	8
30	Use of antipsychotics, benzodiazepine derivatives, and dementia medication among older people with intellectual disability and/or autism spectrum disorder and dementia. Research in Developmental Disabilities, 2017, 62, 50-57.	2.2	22
31	Socioeconomic status and geographical factors associated with active listing in primary care: a cross-sectional population study accounting for multimorbidity, age, sex and primary care. BMJ Open, 2017, 7, e014984.	1.9	5
32	Effects of an intervention (SÃ,,KLÃ,,K) on prescription of potentially inappropriate medication in elderly patients. Family Practice, 2017, 34, 213-218.	1.9	10
33	Prevention in the elderly: A necessary priority for general practitioners. European Journal of General Practice, 2017, 23, 203-208.	2.0	16
34	Yoga – a laborious way to well-being: patients' experiences of yoga as a treatment for hypertension in primary care. Scandinavian Journal of Primary Health Care, 2017, 35, 360-368.	1.5	5
35	Use of Antipsychotic Drugs by Elderly Primary Care Patients and the Effects of Medication Reviews: A Cross-Sectional Study in Sweden. Drugs - Real World Outcomes, 2017, 4, 159-165.	1.6	5
36	Longitudinal trends in self-reported anxiety. Effects of age and birth cohort during 25 years. BMC Psychiatry, 2017, 17, 119.	2.6	41

#	Article	lF	Citations
37	Prescription of potentially inappropriate medications among older people with intellectual disability: a register study. BMC Pharmacology & Empty Toxicology, 2017, 18, 68.	2.4	25
38	Trends in treatment with antipsychotic medication in relation to national directives, in people with dementia $\hat{a} \in \text{``a review of the Swedish context. BMC Psychiatry, 2017, 17, 251.}$	2.6	5
39	Factors related to falls, weightâ€loss and pressure ulcers – more insight in risk assessment among nursing home residents. Journal of Clinical Nursing, 2016, 25, 940-950.	3.0	25
40	Being in a Bubble: the experience of loneliness among frail older people. Journal of Advanced Nursing, 2016, 72, 631-640.	3.3	50
41	Prevalence and predictive importance of anemia in Swedish nursing home residents – a longitudinal study. BMC Geriatrics, 2016, 16, 206.	2.7	10
42	Blood pressure and all-cause mortality: a prospective study of nursing home residents. Age and Ageing, 2016, 45, 826-832.	1.6	15
43	Safer drug use in primary care - a pilot intervention study to identify improvement needs and make agreements for change in five Swedish primary care units. BMC Family Practice, 2016, 17, 140.	2.9	10
44	Factors associated with increased hospitalisation risk among nursing home residents in Sweden: a prospective study with a threeâ€year followâ€up. International Journal of Older People Nursing, 2016, 11, 130-139.	1.3	25
45	Indication for pharmacological treatment is often lacking: a cross-sectional study on the quality of drug therapy among the elderly. BMC Geriatrics, 2015 , 15 , 117 .	2.7	7
46	Healthcare system intervention for safer use of medicines in elderly patients in primary care—a qualitative study of the participants' perceptions of self-assessment, peer review, feedback and agreement for change. BMC Family Practice, 2015, 16, 117.	2.9	8
47	Women's health in the Lund area (WHILA) - Alcohol consumption and all-cause mortality among women – a 17Âyear follow-up study. BMC Public Health, 2015, 16, 22.	2.9	9
48	The longitudinal age and birth cohort trends of regular exercise among adults aged 16–63 years in Sweden: a 24-year follow-up study. Population Health Metrics, 2015, 13, 18.	2.7	4
49	Cost-utility analysis of case management for frail older people: effects of a randomised controlled trial. Health Economics Review, 2015, 5, 51.	2.0	22
50	Loneliness and health care consumption among older people. Scandinavian Journal of Caring Sciences, 2015, 29, 435-443.	2.1	71
51	"lt is meaningful; I feel that I can make a difference―A qualitative study about GPs' experiences of work at nursing homes in Sweden. BMC Family Practice, 2015, 16, 111.	2.9	13
52	Surgery for urinary incontinence in women 65 years and older: a systematic review. International Urogynecology Journal, 2015, 26, 1095-1102.	1.4	15
53	Longitudinal trends in good self-rated health: effects of age and birth cohort in a 25-year follow-up study in Sweden. International Journal of Public Health, 2015, 60, 363-373.	2.3	33
54	The effect of conservative treatment of urinary incontinence among older and frail older people: a systematic review. Age and Ageing, 2015, 44, 736-744.	1.6	40

#	Article	IF	Citations
55	Yoga's effect on inflammatory biomarkers and metabolic risk factors in a high risk population – a controlled trial in primary care. BMC Cardiovascular Disorders, 2015, 15, 91.	1.7	18
56	Effect of pharmacological treatment for urinary incontinence in the elderly and frail elderly: A systematic review. Geriatrics and Gerontology International, 2015, 15, 521-534.	1.5	45
57	The nutritional situation in Swedish nursing homes – A longitudinal study. Archives of Gerontology and Geriatrics, 2015, 60, 128-133.	3.0	36
58	Effects on healthcare utilization of case management for frail older people: A randomized controlled trial (RCT). Archives of Gerontology and Geriatrics, 2015, 60, 71-81.	3.0	26
59	Depressive symptoms among frail elderly in ordinary living: who is affected and who is treated?. Aging and Mental Health, 2014, 18, 1022-1028.	2.8	4
60	Is use of fall risk-increasing drugs in an elderly population associated with an increased risk of hip fracture, after adjustment for multimorbidity level: a cohort study. BMC Geriatrics, 2014, 14, 131.	2.7	72
61	Drugs prescribed by general practitioners according to age, gender and socioeconomic status after adjustment for multimorbidity level. BMC Family Practice, 2014, 15, 183.	2.9	12
62	Effects of a pharmacist-led structured medication review in primary care on drug-related problems and hospital admission rates: a randomized controlled trial. Scandinavian Journal of Primary Health Care, 2014, 32, 180-186.	1.5	81
63	Importance of healthcare utilization and multimorbidity level in choosing a primary care provider in Sweden. Scandinavian Journal of Primary Health Care, 2014, 32, 99-105.	1.5	12
64	Vitamin D deficiency in elderly people in Swedish nursing homes is associated with increased mortality. European Journal of Endocrinology, 2014, 170, 667-675.	3.7	42
65	Fighting for control in an unpredictable life – a qualitative study of older persons' experiences of living with chronic dizziness. BMC Geriatrics, 2014, 14, 97.	2.7	16
66	The longitudinal exercise trend among older Swedes aged 53–84 years – a 16-year follow-up study. BMC Public Health, 2014, 14, 1327.	2.9	10
67	Swedish general practitioners' attitudes towards treatment guidelines – a qualitative study. BMC Family Practice, 2014, 15, 199.	2.9	13
68	Depression and use of antidepressants in Swedish nursing homes: a 12-month follow-up study. International Psychogeriatrics, 2014, 26, 669-675.	1.0	13
69	Case management for frail older people – a qualitative study of receivers' and providers' experiences of a complex intervention. BMC Health Services Research, 2014, 14, 14.	2.2	55
70	The rate and nature of medication errors among elderly upon admission to hospital after implementation of clinical pharmacist-led medication reconciliation. European Journal of Hospital Pharmacy, 2014, 21, 156-160.	1.1	6
71	Fall risk-increasing drugs and falls: a cross-sectional study among elderly patients in primary care. BMC Geriatrics, 2014, 14, 40.	2.7	113
72	Can gender difference in prescription drug use be explained by gender-related morbidity?: a study on a Swedish population during 2006. BMC Public Health, 2014, 14, 329.	2.9	16

#	Article	IF	Citations
73	Effects of a One-Year Home-Based Case Management Intervention on Falls in Older People: A Randomized Controlled Trial. Journal of Aging and Physical Activity, 2014, 22, 457-464.	1.0	18
74	Swedish hunters' safety behaviour and experience of firearm incidents. Accident Analysis and Prevention, 2013, 60, 64-70.	5.7	6
75	Longitudinal age-and cohort trends in body mass index in Sweden – a 24-year follow-up study. BMC Public Health, 2013, 13, 893.	2.9	34
76	Impact of yoga on blood pressure and quality of life in patients with hypertension – a controlled trial in primary care, matched for systolic blood pressure. BMC Cardiovascular Disorders, 2013, 13, 111.	1.7	51
77	Prevalence and treatment of heart failure in Swedish nursing homes. BMC Geriatrics, 2013, 13, 118.	2.7	9
78	Prevalence and predictors of falls and dizziness in people younger and older than 80 years of ageâ€"A longitudinal cohort study. Archives of Gerontology and Geriatrics, 2013, 56, 160-168.	3.0	49
79	In-hospital medication reviews reduce unidentified drug-related problems. European Journal of Clinical Pharmacology, 2013, 69, 647-655.	1.9	38
80	Improving the Quality of Pharmacotherapy in Elderly Primary Care Patients Through Medication Reviews: A Randomised Controlled Study. Drugs and Aging, 2013, 30, 235-246.	2.7	120
81	Low level of antimicrobial resistance in Escherichia coli among Swedish nursing home residents. Scandinavian Journal of Infectious Diseases, 2013, 45, 117-123.	1.5	8
82	Theory-based interventions to reduce prescription of antibioticsa randomized controlled trial in Sweden. Family Practice, 2013, 30, 634-640.	1.9	9
83	The hospital LIMM-based clinical pharmacy service improves the quality of the patient medication process and saves time. European Journal of Hospital Pharmacy, 2012, 19, 375-377.	1.1	11
84	Colonization with Staphylococcus aureus in Swedish nursing homes: A cross-sectional study. Scandinavian Journal of Infectious Diseases, 2012, 44, 3-8.	1.5	7
85	Lack of adherence to hypertension treatment guidelines among GPs in southern Sweden-A case report-based survey. BMC Family Practice, 2012, 13, 34.	2.9	10
86	Frail elderly patients' experiences of information on medication. A qualitative study. BMC Geriatrics, 2012, 12, 46.	2.7	35
87	GP's Adherence to Guidelines for Cardiovascular Disease among Elderly: A Quality Development Study. Scientific World Journal, The, 2012, 2012, 1-7.	2.1	11
88	Acceptance and importance of clinical pharmacists' LIMM-based recommendations. International Journal of Clinical Pharmacy, 2012, 34, 272-276.	2.1	33
89	Prevalence and predictors of healthcare utilization among older people (60+): Focusing on ADL dependency and risk of depression. Archives of Gerontology and Geriatrics, 2012, 54, e349-e363.	3.0	20
90	The effect of medication reconciliation in elderly patients at hospital discharge. International Journal of Clinical Pharmacy, 2012, 34, 113-119.	2.1	39

#	Article	IF	Citations
91	How can we Improve Drug Safety When Treating Frail Elderly Patients?. Advances in Pharmacoepidemiology & Drug Safety, 2012, 01, .	0.1	О
92	The process of identifying, solving and preventing drug related problems in the LIMM-study. International Journal of Clinical Pharmacy, 2011, 33, 1010-1018.	2.1	36
93	Impact of the Lund Integrated Medicines Management (LIMM) model on medication appropriateness and drug-related hospital revisits. European Journal of Clinical Pharmacology, 2011, 67, 741-752.	1.9	130
94	The assessment of renal function in relation to the use of drugs in elderly in nursing homes; a cohort study. BMC Geriatrics, $2011, 11, 1$.	2.7	108
95	Psychosocial perspectives on health care utilization among frail elderly people: An explorative study. Archives of Gerontology and Geriatrics, 2011, 52, 290-294.	3.0	15
96	How to assess frailty and the need for care? Report from the Study of Health and Drugs in the Elderly (SHADES) in community dwellings in Sweden. Archives of Gerontology and Geriatrics, 2011, 53, 40-45.	3.0	56
97	Healthcare utilisation and knowledge concerning prescribed drugs among older people. European Journal of Clinical Pharmacology, 2010, 66, 1047-1054.	1.9	6
98	Case managers for frail older people: a randomised controlled pilot study. Scandinavian Journal of Caring Sciences, 2010, 24, 755-763.	2.1	17
99	Frail elderly patients in primary care—their medication knowledge and beliefs about prescribed medicines. European Journal of Clinical Pharmacology, 2009, 65, 151-155.	1.9	54
100	Improved quality in the hospital discharge summary reduces medication errorsâ€"LIMM: Landskrona Integrated Medicines Management. European Journal of Clinical Pharmacology, 2009, 65, 1037-1046.	1.9	90
101	A multiâ€intervention approach on drug therapy can lead to a more appropriate drug use in the elderly. LIMMâ€Landskrona Integrated Medicines Management. Journal of Evaluation in Clinical Practice, 2009, 15, 660-667.	1.8	61
102	Aging and Drugs. , 2009, , 1-7.		2
103	Strategies and Interventions for Improvement. , 2009, , 119-127.		0
104	Quality of Care in the Elderly. , 2009, , 21-35.		0
105	Clinical outcomes from the use of Medication Report when elderly patients are discharged from hospital. International Journal of Clinical Pharmacy, 2008, 30, 840-845.	1.4	32
106	Barriers to adherence to hypertension guidelines among GPs in southern Sweden: A survey. Scandinavian Journal of Primary Health Care, 2008, 26, 154-159.	1.5	42
107	Medication report reduces number of medication errors when elderly patients are discharged from hospital. International Journal of Clinical Pharmacy, 2007, 30, 92-98.	1.4	54
108	High blood pressure despite treatment: Results from a cross-sectional primary healthcare-based study in southern Sweden. Scandinavian Journal of Primary Health Care, 2006, 24, 224-230.	1,5	24

Patrik Midlöv

#	Article	IF	CITATIONS
109	Effects of educational outreach visits on prescribing of benzodiazepines and antipsychotic drugs to elderly patients in primary health care in southern Sweden. Family Practice, 2006, 23, 60-64.	1.9	35
110	Medication errors when transferring elderly patients between primary health care and hospital care. International Journal of Clinical Pharmacy, 2005, 27, 116-120.	1.4	78
111	Pharmacotherapeutic interventions by a multi-speciality team: opinions of the general practitioners and nurses. European Journal of Clinical Pharmacology, 2003, 59, 65-69.	1.9	4
112	Descriptive study and pharmacotherapeutic intervention in patients with epilepsy or Parkinson's disease at nursing homes in southern Sweden. European Journal of Clinical Pharmacology, 2002, 57, 903-910.	1.9	34
113	Obstacles and Opportunities in Information Transfer Regarding Medications at Discharge – A Focus Group Study with Hospital Physicians. Drug, Healthcare and Patient Safety, 0, Volume 14, 61-73.	2.5	4