

Maarten H Vermeer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5206533/publications.pdf>

Version: 2024-02-01

213
papers

15,468
citations

23879

60
h-index

20625

120
g-index

216
all docs

216
docs citations

216
times ranked

8276
citing authors

#	ARTICLE	IF	CITATIONS
1	Deregulation of JAK2 signaling underlies primary cutaneous CD8 ⁺ aggressive epidermotropic cytotoxic T-cell lymphoma. <i>Haematologica</i> , 2022, 107, 702-714.	1.7	20
2	Whole-genome profiling of primary cutaneous anaplastic large cell lymphoma. <i>Haematologica</i> , 2022, 107, 1619-1632.	1.7	9
3	Genetic and epigenetic insights into cutaneous T-cell lymphoma. <i>Blood</i> , 2022, 139, 15-33.	0.6	28
4	Dermatologic Events Associated with the Anti-CCR4 Antibody Mogamulizumab: Characterization and Management. <i>Dermatology and Therapy</i> , 2022, 12, 29-40.	1.4	17
5	Primary cutaneous lymphoma: recommendations for clinical trial design and staging update from the ISCL, USCLC, and EORTC. <i>Blood</i> , 2022, 140, 419-437.	0.6	58
6	Cell-of-origin classification using the Hans and Lymph2Cx algorithms in primary cutaneous large B-cell lymphomas. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022, 480, 667-675.	1.4	12
7	Total skin electron beam therapy for cutaneous T-cell lymphomas in the Netherlands: A retrospective analysis of treatment outcomes and selection for high or low dose schedule. <i>Clinical and Translational Radiation Oncology</i> , 2022, 33, 77-82.	0.9	3
8	Flow cytometry for the assessment of blood tumour burden in cutaneous T-cell lymphoma: towards a standardized approach. <i>British Journal of Dermatology</i> , 2022, 187, 21-28.	1.4	9
9	Mass Cytometric Analysis of Early-Stage Mycosis Fungoides. <i>Cells</i> , 2022, 11, 1062.	1.8	1
10	Topical Bimiralisib Shows Meaningful Cutaneous Drug Levels in Healthy Volunteers and Mycosis Fungoides Patients but No Clinical Activity in a First-in-Human, Randomized Controlled Trial. <i>Cancers</i> , 2022, 14, 1510.	1.7	1
11	Tumor Clone Frequency Calculation Using High-Throughput Sequencing of the TCR β Gene in Patients with Folliculotropic Mycosis Fungoides. <i>Journal of Investigative Dermatology</i> , 2022, 142, 2544-2546.e2.	0.3	0
12	Treatment of early-stage mycosis fungoides: results from the PROspective Cutaneous Lymphoma International Prognostic Index (PROCLIP) study*. <i>British Journal of Dermatology</i> , 2021, 184, 722-730.	1.4	39
13	Should we be imaging lymph nodes at initial diagnosis of early-stage mycosis fungoides? Results from the PROspective Cutaneous Lymphoma International Prognostic Index (PROCLIP) international study*. <i>British Journal of Dermatology</i> , 2021, 184, 524-531.	1.4	18
14	The importance of assessing blood tumour burden in cutaneous T-cell lymphoma*. <i>British Journal of Dermatology</i> , 2021, 185, 19-25.	1.4	12
15	Phenotypical Markers, Molecular Mutations, and Immune Microenvironment as Targets for New Treatments in Patients with Mycosis Fungoides and/or S α zary Syndrome. <i>Journal of Investigative Dermatology</i> , 2021, 141, 484-495.	0.3	31
16	Epidemiology of cutaneous lymphoma. <i>British Journal of Dermatology</i> , 2021, 184, 993-994.	1.4	4
17	Incidence of mycosis fungoides and S α zary syndrome in the Netherlands between 2000 and 2020. <i>British Journal of Dermatology</i> , 2021, 185, 434-435.	1.4	15
18	Serum and cutaneous transcriptional expression levels of IL31 are minimal in cutaneous T cell lymphoma variants. <i>Biochemistry and Biophysics Reports</i> , 2021, 26, 101007.	0.7	2

#	ARTICLE	IF	CITATIONS
19	Improved SÅ©zary cell detection and novel insights into immunophenotypic and molecular heterogeneity in SÅ©zary syndrome. <i>Blood</i> , 2021, 138, 2539-2554.	0.6	28
20	Cutaneous T cell lymphoma. <i>Nature Reviews Disease Primers</i> , 2021, 7, 61.	18.1	70
21	Quality of life in patients with Mycosis Fungoides and SÅ©zary Syndrome: a systematic review of the literature. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2021, 35, 2377-2387.	1.3	20
22	Primary cutaneous peripheral Tâ€cell lymphoma, not otherwise specified: results of a multicentre European Organization for Research and Treatment of Cancer (EORTC) cutaneous lymphoma taskforce study on the clinicoâ€pathological and prognostic features. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2021, 35, 658-668.	1.3	12
23	Characterization and outcomes in patients with mogamulizumab-associated skin reactions in the MAVORIC trial. <i>European Journal of Cancer</i> , 2021, 156, S46.	1.3	5
24	Frequency and prognosis of associated malignancies in 504 patients with lymphomatoid papulosis. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2020, 34, 260-266.	1.3	31
25	Wholeâ€genome analysis uncovers recurrent <i>IKZF1</i> inactivation and aberrant cell adhesion in blastic plasmacytoid dendritic cell neoplasm. <i>Genes Chromosomes and Cancer</i> , 2020, 59, 295-308.	1.5	14
26	Plaque stage folliculotropic mycosis fungoides: histopathologic features and prognostic factors in a series of 40 patients. <i>Journal of Cutaneous Pathology</i> , 2020, 47, 241-250.	0.7	20
27	Human papillomavirus 16â€positive supraclavicular cutaneous squamous cell carcinoma metastatic to the level IV supraclavicular lymph nodes. <i>JAAD Case Reports</i> , 2020, 6, 822-825.	0.4	0
28	Botulinum toxin type A in the treatment of Raynaud's phenomenon. <i>Dermatologic Therapy</i> , 2020, 33, e14182.	0.8	6
29	Cucurbitacin E and I target the JAK/STAT pathway and induce apoptosis in SÅ©zary cells. <i>Biochemistry and Biophysics Reports</i> , 2020, 24, 100832.	0.7	12
30	A baby with red plaques on the face and a first-degree heart block: neonatal lupus. <i>Lancet, The</i> , 2020, 396, 1432.	6.3	0
31	Management of primary cutaneous lymphoma patients during COVIDâ€19 pandemic: EORTC CLTF guidelines. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2020, 34, 1633-1636.	1.3	14
32	Staphylococcus aureus enterotoxins induce FOXP3 in neoplastic T cells in SÅ©zary syndrome. <i>Blood Cancer Journal</i> , 2020, 10, 57.	2.8	24
33	Outcomes of rare patients with a primary cutaneous CD30+ lymphoproliferative disorder developing extracutaneous disease. <i>Blood</i> , 2020, 135, 769-773.	0.6	9
34	Clinical and pathogenic aspects of the severe cutaneous adverse reaction epidermal necrolysis (EN). <i>Journal of the European Academy of Dermatology and Venereology</i> , 2020, 34, 1957-1971.	1.3	25
35	Clinical, Histologic, and Molecular Characteristics of Anaplastic Lymphoma Kinase-positive Primary Cutaneous Anaplastic Large Cell Lymphoma. <i>American Journal of Surgical Pathology</i> , 2020, 44, 776-781.	2.1	25
36	Sentinel node biopsy in cutaneous melanoma patients with germline CDKN2A mutations. <i>Melanoma Research</i> , 2020, 30, 630-631.	0.6	1

#	ARTICLE	IF	CITATIONS
37	Characterization and Outcomes in Patients with Mogamulizumab-Associated Skin Reactions in the MAVORIC Trial. <i>Blood</i> , 2020, 136, 23-24.	0.6	5
38	Acquired N-Linked Glycosylation Motifs in B-Cell Receptors of Primary Cutaneous B-Cell Lymphoma and the Normal B-Cell Repertoire. <i>Journal of Investigative Dermatology</i> , 2019, 139, 2195-2203.	0.3	12
39	A rare case of cutaneous Epstein-Barr virus-negative intravascular cytotoxic T-cell lymphoma. <i>JAAD Case Reports</i> , 2019, 5, 548-551.	0.4	3
40	Cell adhesion molecule 1 (CADM1) can be a biomarker for SÅ©zary syndrome. <i>European Journal of Cancer</i> , 2019, 119, S7.	1.3	0
41	Prognostic factors in mycosis fungoides: the PROCLIP study. <i>European Journal of Cancer</i> , 2019, 119, S26.	1.3	1
42	Treatment of rare patients with a primary cutaneous CD30+ lymphoproliferation developing extracutaneous disease: a retrospective study of 43 patients. <i>European Journal of Cancer</i> , 2019, 119, S28-S29.	1.3	0
43	PD-1 overexpression in SÅ©zary syndrome is epigenetically regulated. <i>European Journal of Cancer</i> , 2019, 119, S2.	1.3	0
44	Molecular analysis of primary cutaneous diffuse large B-cell lymphoma, leg type at diagnosis and relapse. <i>European Journal of Cancer</i> , 2019, 119, S8-S9.	1.3	0
45	Whole-genome analysis uncovers recurrent IKZF1 inactivation and aberrant cell adhesion in blastic plasmacytoid dendritic cell neoplasm. <i>European Journal of Cancer</i> , 2019, 119, S9.	1.3	0
46	Clinical, histological and molecular characteristics of ALK-positive primary cutaneous anaplastic large cell lymphoma. <i>European Journal of Cancer</i> , 2019, 119, S18.	1.3	0
47	Treatment of early-phase mycosis fungoides: results from the Prospective Cutaneous Lymphoma International (PROCLIP) study. <i>European Journal of Cancer</i> , 2019, 119, S27.	1.3	0
48	Time to Next Treatment in Patients with Previously Treated Cutaneous T-Cell Lymphoma (CTCL) Receiving Mogamulizumab or Vorinostat: A MAVORIC Post-Hoc Analysis. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2019, 19, S361.	0.2	0
49	T-cell receptor rearrangements in the skin and blood of patients in the PROCLIP study: detection of clonal rearrangements in the skin (and blood) correlates with the B-class of MF and SS patients. <i>European Journal of Cancer</i> , 2019, 119, S25.	1.3	0
50	IPH4102, a first-in-class anti-KIR3DL2 monoclonal antibody, in patients with relapsed or refractory cutaneous T-cell lymphoma: an international, first-in-human, open-label, phase 1 trial. <i>Lancet Oncology</i> , 2019, 20, 1160-1170.	5.1	119
51	Generalized Molluscum Contagiosum Successfully Treated with Interferon-Alpha in a Patient with Folliculotropic Mycosis Fungoides. <i>Case Reports in Dermatology</i> , 2019, 11, 52-56.	0.3	6
52	Antibiotics can improve CTCL. <i>Blood</i> , 2019, 134, 1000-1001.	0.6	4
53	TIME TO NEXT TREATMENT IN PATIENTS WITH PREVIOUSLY TREATED CUTANEOUS T-CELL LYMPHOMA (CTCL) RECEIVING MOGAMULIZUMAB OR VORINOSTAT: A POST-HOC ANALYSIS OF THE MAVORIC STUDY. <i>Hematological Oncology</i> , 2019, 37, 285-286.	0.8	3
54	Corresponding anaplastic lymphoma kinase-tropomyosin 3 (ALK-TPM3) fusion in a patient with a primary cutaneous anaplastic large-cell lymphoma and a Spitz nevus. <i>JAAD Case Reports</i> , 2019, 5, 970-972.	0.4	4

#	ARTICLE	IF	CITATIONS
55	Loss of the GPIâ€ anchor in Bâ€ lymphoblastic leukemia by epigenetic downregulation of <i>PIGH</i> expression. American Journal of Hematology, 2019, 94, 93-102.	2.0	8
56	Pitfalls in diagnosing primary cutaneous aggressive epidermotropic CD8 ⁺ Tâ€ cell lymphoma. British Journal of Dermatology, 2019, 180, 411-412.	1.4	8
57	The PROCLIP international registry of earlyâ€ stage mycosis fungoides identifies substantial diagnostic delay in most patients. British Journal of Dermatology, 2019, 181, 350-357.	1.4	127
58	Safety of Mogamulizumab in Mycosis Fungoides and SÃ©zary Syndrome: Final Results from the Phase 3 Mavoric Study. Blood, 2019, 134, 5300-5300.	0.6	3
59	Time to next treatment in patients with previously treated cutaneous T-cell lymphoma (CTCL) receiving mogamulizumab or vorinostat: A MAVORIC post-hoc analysis.. Journal of Clinical Oncology, 2019, 37, 7539-7539.	0.8	4
60	Evaluation of treatment results in multifocal primary cutaneous anaplastic large cell lymphoma: report of the Dutch Cutaneous Lymphoma Group. British Journal of Dermatology, 2018, 179, 724-731.	1.4	19
61	Blood classification and blood response criteria in mycosis fungoides and SÃ©zary syndrome using flow cytometry: recommendations from the EORTC cutaneous lymphoma task force. European Journal of Cancer, 2018, 93, 47-56.	1.3	105
62	High prevalence of MYD88 and CD79B mutations in intravascular large B-cell lymphoma. Blood, 2018, 131, 2086-2089.	0.6	69
63	Classification and recommended treatment options for folliculotropic mycosis fungoides. Expert Opinion on Orphan Drugs, 2018, 6, 35-45.	0.5	1
64	516 Global collaboration for establishment of a prognostic index in mycosis fungoides & Sezary Syndrome. Journal of Investigative Dermatology, 2018, 138, S88.	0.3	3
65	Folliculotropic mycosis fungoides presenting with a solitary lesion: Clinicopathological features and longâ€ term followâ€ up data in a series of 9 cases. Journal of Cutaneous Pathology, 2018, 45, 122-128.	0.7	12
66	High Incidence and Clinical Significance of MYC Rearrangements in Primary Cutaneous Diffuse Large B-Cell Lymphoma, Leg Type. American Journal of Surgical Pathology, 2018, 42, 1488-1494.	2.1	42
67	Remission of psoriasis during treatment with sorafenib. JAAD Case Reports, 2018, 4, 1065-1067.	0.4	10
68	Arrayâ€ based CGH of primary cutaneous CD8 ⁺ aggressive EPIDERMOTROPIC cytotoxic Tâ€ cell lymphoma. Genes Chromosomes and Cancer, 2018, 57, 622-629.	1.5	11
69	Levels of IL-31 in different variants of cutaneous T cell lymphomas. European Journal of Cancer, 2018, 101, S17.	1.3	1
70	Lymph node imaging in patch/plaque mycosis fungoides; enlarged LN are infrequent but lymphomatous nodal involvement may occur and upstage patients to advanced disease. European Journal of Cancer, 2018, 101, S25-S26.	1.3	0
71	An Integrated Data Resource for Genomic Analysis of Cutaneous T-Cell Lymphoma. Journal of Investigative Dermatology, 2018, 138, 2681-2683.	0.3	38
72	Genomic analysis reveals recurrent deletion of HNRNPK and SOCS1 in mycosis fungoides. European Journal of Cancer, 2018, 101, S7.	1.3	0

#	ARTICLE	IF	CITATIONS
73	Treatment of early-stage mycosis fungoides: Results from the PROCLIP study. <i>European Journal of Cancer</i> , 2018, 101, S34.	1.3	0
74	The PROCLIP study; a prototype registry for rare disease with global collaboration for establishment of a prognostic index in mycosis fungoides and Sezary syndrome. <i>European Journal of Cancer</i> , 2018, 101, S6.	1.3	0
75	Frequency of associated haematologic malignancies in lymphomatoid papulosis: A preliminary report of 505 patients from the Dutch Cutaneous Lymphoma Group. <i>European Journal of Cancer</i> , 2018, 101, S6.	1.3	0
76	IPH4102 in relapsed/refractory cutaneous T cell lymphoma (CTCL): Results of the first-in-human multicenter phase 1 study. <i>European Journal of Cancer</i> , 2018, 101, S29.	1.3	3
77	Standardized flow cytometry (EuroFlow) demonstrates heterogeneous T-cell origin of Sezary lymphoma cells. <i>European Journal of Cancer</i> , 2018, 101, S8.	1.3	0
78	Cardiomyopathy in patients with epidermolysis bullosa simplex with mutations in <i>KLHL24</i> . <i>British Journal of Dermatology</i> , 2018, 179, 1181-1183.	1.4	23
79	Developments in the understanding of blood involvement and stage in mycosis fungoides/Sezary syndrome. <i>European Journal of Cancer</i> , 2018, 101, 278-280.	1.3	10
80	Genomic analysis reveals recurrent deletion of JAK-STAT signaling inhibitors <i>HNRNPK</i> and <i>SOCS1</i> in mycosis fungoides. <i>Genes Chromosomes and Cancer</i> , 2018, 57, 653-664.	1.5	56
81	Mogamulizumab versus vorinostat in previously treated cutaneous T-cell lymphoma (MAVORIC): an international, open-label, randomised, controlled phase 3 trial. <i>Lancet Oncology</i> , The, 2018, 19, 1192-1204.	5.1	398
82	IPH4102; An Anti-KIR3DL2 Monoclonal Antibody in Refractory Sezary Syndrome: Results from a Multicenter Phase 1 Trial. <i>Blood</i> , 2018, 132, 684-684.	0.6	2
83	Molecular advances in cutaneous T-cell lymphoma. <i>Seminars in Cutaneous Medicine and Surgery</i> , 2018, 37, 81-86.	1.6	22
84	Primary Cutaneous Follicle Center Lymphomas (PCFCL) Express Heavily Mutated B-Cell Receptors with Acquired N-Glycosylation Motifs and Lack Ongoing Somatic Hypermutation. <i>Blood</i> , 2018, 132, 1573-1573.	0.6	3
85	Increased Expression of PLS3 Correlates with Better Outcome in Sezary Syndrome. <i>Journal of Investigative Dermatology</i> , 2017, 137, 754-757.	0.3	7
86	Recommendations for treatment in folliculotropic mycosis fungoides: report of the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2017, 177, 223-228.	1.4	43
87	European Organisation for Research and Treatment of Cancer consensus recommendations for the treatment of mycosis fungoides/Sezary syndrome – Update 2017. <i>European Journal of Cancer</i> , 2017, 77, 57-74.	1.3	363
88	Accurate Quantification of T Cells by Measuring Loss of Germline T-Cell Receptor Loci with Generic Single Duplex Droplet Digital PCR Assays. <i>Journal of Molecular Diagnostics</i> , 2017, 19, 236-243.	1.2	19
89	A phase III study of lenalidomide maintenance after debulking therapy in patients with advanced cutaneous T-cell lymphoma - EORTC 21081 (NCT01098656): results and lessons learned for future trial designs. <i>European Journal of Dermatology</i> , 2017, 27, 286-294.	0.3	16
90	Global patterns of care in advanced stage mycosis fungoides/Sezary syndrome: a multicenter retrospective follow-up study from the Cutaneous Lymphoma International Consortium. <i>Annals of Oncology</i> , 2017, 28, 2517-2525.	0.6	98

#	ARTICLE	IF	CITATIONS
91	Micro RNA miR-155 potentiates tumour development in mycosis fungoides. British Journal of Dermatology, 2017, 177, 618-620.	1.4	6
92	Recommendations for the Optimal Radiation Dose in Patients With Primary Cutaneous Anaplastic Large Cell Lymphoma: A Report of the Dutch Cutaneous Lymphoma Group. International Journal of Radiation Oncology Biology Physics, 2017, 99, 1279-1285.	0.4	19
93	The clinical spectrum of mycosis fungoides in Tanzania, East Africa. British Journal of Dermatology, 2017, 176, 1653-1656.	1.4	6
94	A novel keratin 13 variant in a four-generation family with white sponge nevus. Clinical Case Reports (discontinued), 2017, 5, 1503-1509.	0.2	12
95	Special variant of histiocytosis. BMJ Case Reports, 2017, 2017, bcr-2017-221538.	0.2	2
96	Anti-CCR4 Monoclonal Antibody, Mogamulizumab, Demonstrates Significant Improvement in PFS Compared to Vorinostat in Patients with Previously Treated Cutaneous T-Cell Lymphoma (CTCL): Results from the Phase III MAVORIC Study. Blood, 2017, 130, 817-817.	0.6	15
97	Genetic rearrangements result in altered gene expression and novel fusion transcripts in S�azary syndrome. Oncotarget, 2017, 8, 39627-39639.	0.8	41
98	Evaluation of Immunophenotypic and Molecular Biomarkers for S�azary Syndrome Using Standard Operating Procedures: A Multicenter Study of 59 Patients. Journal of Investigative Dermatology, 2016, 136, 1364-1372.	0.3	78
99	183 Cutaneous lymphoma international consortium (CLIC): Uniting worldwide experts to develop and validate a prognostic index model. Journal of Investigative Dermatology, 2016, 136, S32.	0.3	0
100	Epigenomic Analysis of S�azary Syndrome Defines Patterns of Aberrant DNA Methylation and Identifies Diagnostic Markers. Journal of Investigative Dermatology, 2016, 136, 1876-1884.	0.3	46
101	Clinical Staging and Prognostic Factors in Folliculotropic Mycosis Fungoides. JAMA Dermatology, 2016, 152, 992.	2.0	119
102	Score test for association between recurrent events and a terminal event. Statistics in Medicine, 2016, 35, 3037-3048.	0.8	9
103	Abstract LB-012: Autonomous, antigen-independent B-cell receptor signalling as a novel pathogenetic mechanism in non-GCB DLBCL. , 2016, , .		1
104	First-in-Human, Multicenter Phase I Study of IPH4102, First-in-Class Humanized Anti-KIR3DL2 Monoclonal Antibody, in Relapsed/Refractory Cutaneous T-Cell Lymphomas: Preliminary Safety, Exploratory and Clinical Activity Results. Blood, 2016, 128, 1826-1826.	0.6	6
105	First-in-human, open label, multicenter phase I of IPH4102, first-in-class humanized anti-KIR3DL2 monoclonal antibody, in relapsed/refractory cutaneous T-cell lymphomas.. Journal of Clinical Oncology, 2016, 34, TPS2591-TPS2591.	0.8	0
106	The B-Cell Receptor of Primary Cutaneous Follicle Center Lymphoma: Implications for Pathogenesis. Blood, 2016, 128, 4136-4136.	0.6	0
107	Antigen-Independent, Autonomous B-Cell Receptor Signaling As a Dominant Candidate Oncogenic Mechanism in ABC DLBCL. Blood, 2016, 128, 778-778.	0.6	1
108	Limited effect of chemotherapy in cutaneous lymphoma. Blood, 2015, 125, 4-5.	0.6	0

#	ARTICLE	IF	CITATIONS
109	Recommendations for treatment of lymphomatoid papulosis with methotrexate: a report from the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2015, 173, 1319-1322.	1.4	30
110	Histopathological and immunophenotypical criteria for the diagnosis of SÅ©zary syndrome in differentiation from other erythrodermic skin diseases: a European Organisation for Research and Treatment of Cancer (EORTC) Cutaneous Lymphoma Task Force Study of 9. <i>British Journal of Dermatology</i> , 2015, 173, 93-105.	1.4	67
111	Differential expression of <sc>TOX</sc> by skinâ€infiltrating T cells in SÅ©zary syndrome and erythrodermic dermatitis. <i>Journal of Cutaneous Pathology</i> , 2015, 42, 604-609.	0.7	17
112	Mutation in PIGA Results in a CD52-Negative Escape Variant in a SÅ©zary Syndrome Patient during Alemtuzumab Treatment. <i>Journal of Investigative Dermatology</i> , 2015, 135, 1199-1202.	0.3	4
113	Genetic ablation of macrohistone H2A1 leads to increased leanness, glucose tolerance and energy expenditure in mice fed a high-fat diet. <i>International Journal of Obesity</i> , 2015, 39, 331-338.	1.6	20
114	Microarray Techniques to Analyze Copy-Number Alterations in Genomic DNA: Array Comparative Genomic Hybridization and Single-Nucleotide Polymorphism Array. <i>Journal of Investigative Dermatology</i> , 2015, 135, 1-5.	0.3	6
115	Cutaneous Lymphoma International Consortium Study of Outcome in Advanced Stages of Mycosis Fungoides and SÅ©zary Syndrome: Effect of Specific Prognostic Markers on Survival and Development of a Prognostic Model. <i>Journal of Clinical Oncology</i> , 2015, 33, 3766-3773.	0.8	328
116	The mutational landscape of cutaneous T cell lymphoma and SÅ©zary syndrome. <i>Nature Genetics</i> , 2015, 47, 1465-1470.	9.4	322
117	EPHA4 is overexpressed but not functionally active in SÅ©zary syndrome. <i>Oncotarget</i> , 2015, 6, 31868-31876.	0.8	6
118	The Mutational Landscape of CTCL and Sezary Syndrome. <i>Blood</i> , 2015, 126, 573-573.	0.6	17
119	Loss of Pigh Expression Frequently Results in a GPI-Negative Subclone Lacking CD52 Membrane Expression, Conferring Alemtuzumab Resistance to B Cell Acute Lymphoblastic Leukemia. <i>Blood</i> , 2015, 126, 1435-1435.	0.6	0
120	Nuclear Factor-ÎB Pathwayâ€Activating Gene Aberrancies in Primary Cutaneous Large B-Cell Lymphoma, Leg Type. <i>Journal of Investigative Dermatology</i> , 2014, 134, 290-292.	0.3	54
121	Exploring the IL-21â€STAT3 Axis as Therapeutic Target for SÅ©zary Syndrome. <i>Journal of Investigative Dermatology</i> , 2014, 134, 2639-2647.	0.3	23
122	Quantitation of tumour development correlates with prognosis in tumour stage (stage IIB) mycosis fungoides. <i>British Journal of Dermatology</i> , 2014, 170, 1080-1086.	1.4	5
123	Recent advances in primary cutaneous B-cell lymphomas. <i>Current Opinion in Oncology</i> , 2014, 26, 230-236.	1.1	13
124	Methotrexate-associated B-cell Lymphoproliferative Disorders Presenting in the Skin. <i>American Journal of Surgical Pathology</i> , 2014, 38, 999-1006.	2.1	42
125	Prognostic factors, prognostic indices and staging in mycosis fungoides and SÅ©zary syndrome: where are we now?. <i>British Journal of Dermatology</i> , 2014, 170, 1226-1236.	1.4	121
126	A Cutaneous Lymphoma International Consortium â€CLICâ€™ study of Prognostic Parameters in Advanced Stages of Mycosis Fungoides and Sezary Syndrome: Progress Towards Establishing a Prognostic Index to Augment Clinical Staging. <i>Blood</i> , 2014, 124, 1621-1621.	0.6	2

#	ARTICLE	IF	CITATIONS
127	MicroRNA Profiling of Primary Cutaneous Large B-Cell Lymphomas. PLoS ONE, 2013, 8, e82471.	1.1	20
128	Autocrine IL-21 Stimulation Is Involved in the Maintenance of Constitutive STAT3 Activation in S \tilde{A} zary Syndrome. Journal of Investigative Dermatology, 2012, 132, 440-447.	0.3	37
129	Deep-Sequencing Analysis Reveals that the miR-199a2/214 Cluster within DN3os Represents the Vast Majority of Aberrantly Expressed MicroRNAs in S \tilde{A} zary Syndrome. Journal of Investigative Dermatology, 2012, 132, 1520-1522.	0.3	42
130	NOTCH1 Signaling as a Therapeutic Target in S \tilde{A} zary Syndrome. Journal of Investigative Dermatology, 2012, 132, 2810-2817.	0.3	18
131	A novel mouse model for S \tilde{A} zary syndrome using xenotransplantation of S \tilde{A} zary cells into immunodeficient RAG2 \sup mice. Experimental Dermatology, 2012, 21, 706-709.	1.4	18
132	A Meta-Analysis of Gene Expression Data Identifies a Molecular Signature Characteristic for Tumor-Stage Mycosis Fungoides. Journal of Investigative Dermatology, 2012, 132, 2050-2059.	0.3	75
133	Differential Expression of Programmed Death-1 (PD-1) in S \tilde{A} zary Syndrome and Mycosis Fungoides. Archives of Dermatology, 2012, 148, 1379.	1.7	113
134	A Delayed Granulomatous Reaction to a Cosmetic Tattoo of the Eyebrows: A Report of Total Regression After Intralesional Corticosteroid Injections. Dermatologic Surgery, 2012, 38, 951-953.	0.4	15
135	Langerhans cell histiocytosis first presenting in the skin in adults: frequent association with a second haematological malignancy. British Journal of Dermatology, 2012, 167, 1287-1294.	1.4	57
136	Major achievements of the EORTC Cutaneous Lymphoma Task Force (CLTF). European Journal of Cancer, Supplement, 2012, 10, 46-50.	2.2	1
137	Prognostic factors in transformed mycosis fungoides: a retrospective analysis of 100 cases. Blood, 2012, 119, 1643-1649.	0.6	186
138	Primary cutaneous anaplastic large cell lymphoma shows a distinct miRNA expression profile and reveals differences from tumor-stage mycosis fungoides. Experimental Dermatology, 2012, 21, 632-634.	1.4	47
139	miRNA expression profiling of mycosis fungoides. Molecular Oncology, 2011, 5, 273-280.	2.1	91
140	EORTC, ISCL, and USCLC consensus recommendations for the treatment of primary cutaneous CD30-positive lymphoproliferative disorders: lymphomatoid papulosis and primary cutaneous anaplastic large-cell lymphoma*. Blood, 2011, 118, 4024-4035.	0.6	365
141	A restricted clonal T-cell receptor repertoire in S \tilde{A} zary syndrome is indicative of superantigenic stimulation. British Journal of Dermatology, 2011, 165, 78-84.	1.4	11
142	Erythematous and purpuric plaques of the face and scalp in Sweet syndrome. British Journal of Haematology, 2011, 154, 159-159.	1.2	1
143	The histone deacetylase inhibitors vorinostat and romidepsin downmodulate IL10 expression in cutaneous T-cell lymphoma cells. British Journal of Pharmacology, 2011, 162, 1590-1602.	2.7	78
144	MicroRNA-21 Expression in CD4+ T Cells Is Regulated by STAT3 and Is Pathologically Involved in S \tilde{A} zary Syndrome. Journal of Investigative Dermatology, 2011, 131, 762-768.	0.3	116

#	ARTICLE	IF	CITATIONS
145	Alentuzumab Treatment Can Affect CD52 Positive As Well As CD52 Negative GPI-Deficient, Sezary Cells Allowing Long-Term Disease Control. <i>Blood</i> , 2011, 118, 2491-2491.	0.6	0
146	IgM Expression on Paraffin Sections Distinguishes Primary Cutaneous Large B-cell Lymphoma, Leg Type From Primary Cutaneous Follicle Center Lymphoma. <i>American Journal of Surgical Pathology</i> , 2010, 34, 1043-1048.	2.1	86
147	MicroRNA expression in SÅ©zary syndrome: identification, function, and diagnostic potential. <i>Blood</i> , 2010, 116, 1105-1113.	0.6	131
148	Carcinoma erysipeloides in breast cancer. <i>Surgery</i> , 2010, 148, 1040-1041.	1.0	2
149	Keratosis Follicularis Spinulosa Decalvans is caused by mutations in MBTPS2. <i>Human Mutation</i> , 2010, 31, 1125-1133.	1.1	67
150	Chemokine/chemokine receptor interactions in extramedullary leukaemia of the skin in childhood AML: Differential roles for CCR2, CCR5, CXCR4 and CXCR7. <i>Pediatric Blood and Cancer</i> , 2010, 55, 344-348.	0.8	45
151	Oligonucleotide Array-CGH Identifies Genomic Subgroups and Prognostic Markers for Tumor Stage Mycosis Fungoides. <i>Journal of Investigative Dermatology</i> , 2010, 130, 1126-1135.	0.3	71
152	Cutaneous Anaplastic Large Cell Lymphoma and Peripheral T-Cell Lymphoma NOS Show Distinct Chromosomal Alterations and Differential Expression of Chemokine Receptors and Apoptosis Regulators. <i>Journal of Investigative Dermatology</i> , 2010, 130, 563-575.	0.3	62
153	Bullous systemic lupus erythematosus responding to mycophenolate mofetil. <i>European Journal of Dermatology</i> , 2010, 20, 844-5.	0.3	9
154	The EORTC Cutaneous T-Cell Lymphoma (CTCL) Platform. <i>Blood</i> , 2010, 116, 4896-4896.	0.6	0
155	Cutaneous Gamma/Delta T-cell Lymphoma During Treatment with Etanercept for Rheumatoid Arthritis. <i>Acta Dermato-Venereologica</i> , 2009, 89, 653-654.	0.6	30
156	Antipleptin autoantibodies in subepidermal blistering diseases. <i>British Journal of Dermatology</i> , 2009, 161, 762-771.	1.4	16
157	Fine-Mapping Chromosomal Loss at 9p21: Correlation with Prognosis in Primary Cutaneous Diffuse Large B-Cell Lymphoma, Leg Type. <i>Journal of Investigative Dermatology</i> , 2009, 129, 1149-1155.	0.3	84
158	Low-Dose Palliative Radiotherapy for Cutaneous B- and T-Cell Lymphomas. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009, 74, 154-158.	0.4	142
159	Oncogenomic analysis of mycosis fungoides reveals major differences with SÅ©zary syndrome. <i>Blood</i> , 2009, 113, 127-136.	0.6	188
160	Profiling of apoptosis genes identifies distinct types of primary cutaneous large B cell lymphoma. <i>Journal of Pathology</i> , 2008, 215, 340-346.	2.1	20
161	A genomic and expression study of APâ€1 in primary cutaneous Tâ€cell lymphoma: evidence for dysregulated expression of JUNB and JUND in MF and SS. <i>Journal of Cutaneous Pathology</i> , 2008, 35, 899-910.	0.7	57
162	Cucurbitacin I Inhibits Stat3 and Induces Apoptosis in SÅ©zary Cells. <i>Journal of Investigative Dermatology</i> , 2008, 128, 1691-1695.	0.3	74

#	ARTICLE	IF	CITATIONS
163	Ultrastructural studies bearing on the mechanism of UVB-impaired induction of contact hypersensitivity to DNCB in man. <i>Clinical and Experimental Immunology</i> , 2008, 92, 487-493.	1.1	26
164	Coincidence of cutaneous follicle center lymphoma and diffuse large B-cell lymphoma. <i>International Journal of Dermatology</i> , 2008, 47, 21-24.	0.5	5
165	Subcutaneous panniculitis-like T-cell lymphoma: definition, classification, and prognostic factors: an EORTC Cutaneous Lymphoma Group Study of 83 cases. <i>Blood</i> , 2008, 111, 838-845.	0.6	617
166	Novel and Highly Recurrent Chromosomal Alterations in Sezary Syndrome. <i>Cancer Research</i> , 2008, 68, 2689-2698.	0.4	176
167	European Organization for Research and Treatment of Cancer and International Society for Cutaneous Lymphoma consensus recommendations for the management of cutaneous B-cell lymphomas. <i>Blood</i> , 2008, 112, 1600-1609.	0.6	415
168	Genome Wide Analysis of 41 Mycosis Fungoides Tumor Stage Using Array Comparative Genomic Hybridization Technology.. <i>Blood</i> , 2008, 112, 1769-1769.	0.6	0
169	Reclassification of 300 Primary Cutaneous B-Cell Lymphomas According to the New WHO/EORTC Classification for Cutaneous Lymphomas: Comparison With Previous Classifications and Identification of Prognostic Markers. <i>Journal of Clinical Oncology</i> , 2007, 25, 1581-1587.	0.8	278
170	Results of Radiotherapy in 153 Primary Cutaneous B-Cell Lymphomas Classified According to the WHO-EORTC Classification. <i>Archives of Dermatology</i> , 2007, 143, 1520.	1.7	120
171	Gene-expression profiling and array-based CGH classify CD4+CD56+ hematodermic neoplasm and cutaneous myelomonocytic leukemia as distinct disease entities. <i>Blood</i> , 2007, 109, 1720-1727.	0.6	137
172	Acitretin induces capillary leak syndrome in a patient with pustular psoriasis. <i>Journal of the American Academy of Dermatology</i> , 2007, 56, 339-342.	0.6	25
173	Array-Based Comparative Genomic Hybridization Analysis Reveals Recurrent Chromosomal Alterations and Prognostic Parameters in Primary Cutaneous Large B-Cell Lymphoma. <i>Journal of Clinical Oncology</i> , 2006, 24, 296-305.	0.8	125
174	Annular erythema of Sjögren's syndrome. <i>Lancet</i> , The, 2006, 367, 1604.	6.3	8
175	Primary cutaneous follicle center lymphoma and primary cutaneous large B-cell lymphoma, leg type, are both targeted by aberrant somatic hypermutation but demonstrate differential expression of AID. <i>Blood</i> , 2006, 107, 4926-4929.	0.6	51
176	Expression of B-cell transcription factors in primary cutaneous B-cell lymphoma. <i>Modern Pathology</i> , 2006, 19, 1270-1276.	2.9	48
177	First Line Treatment of Aggressive Cutaneous NK/T Cell Lymphomas Based on High Dose Cytarabine and Stem Cell Transplantation Does Not Lead to Durable Remissions.. <i>Blood</i> , 2006, 108, 4669-4669.	0.6	0
178	Primary Cutaneous Marginal Zone B-Cell Lymphoma. <i>Archives of Dermatology</i> , 2005, 141, 1139.	1.7	173
179	Distinct types of primary cutaneous large B-cell lymphoma identified by gene expression profiling. <i>Blood</i> , 2005, 105, 3671-3678.	0.6	266
180	WHO/EORTC classification of cutaneous lymphomas 2005: histological and molecular aspects. <i>Journal of Cutaneous Pathology</i> , 2005, 32, 647-674.	0.7	313

#	ARTICLE	IF	CITATIONS
181	A 2-bp deletion in the CJA1 gene is associated with oculo-dento-digital dysplasia with palmoplantar keratoderma. American Journal of Medical Genetics, Part A, 2005, 132A, 171-174.	0.7	73
182	High numbers of tumour-infiltrating activated cytotoxic T lymphocytes, and frequent loss of HLA class I and II expression, are features of aggressive B cell lymphomas of the brain and testis. Journal of Pathology, 2005, 206, 328-336.	2.1	57
183	Epigenetic Profiling of Cutaneous T-Cell Lymphoma: Promoter Hypermethylation of Multiple Tumor Suppressor Genes Including BCL7a, PTPRG, and p73. Journal of Clinical Oncology, 2005, 23, 3886-3896.	0.8	224
184	A missense mutation in the type II hair keratin hHb3 is associated with monilethrix. Journal of Medical Genetics, 2005, 42, e19-e19.	1.5	70
185	WHO-EORTC classification for cutaneous lymphomas. Blood, 2005, 105, 3768-3785.	0.6	3,529
186	Aberrant Expression of the Tyrosine Kinase Receptor EphA4 and the Transcription Factor Twist in SÅazary Syndrome Identified by Gene Expression Analysis. Cancer Research, 2004, 64, 5578-5586.	0.4	155
187	Chromosomal Aberration Patterns Differ in Subtypes of Primary Cutaneous B Cell Lymphomas. Journal of Investigative Dermatology, 2004, 122, 1495-1502.	0.3	67
188	Molecular Cytogenetic Analysis of Chromosomal Breakpoints in the IGH, MYC, BCL6, and MALT1 Gene Loci in Primary Cutaneous B-cell Lymphomas. Journal of Investigative Dermatology, 2004, 123, 213-219.	0.3	105
189	Site-Specific Expression of Polycomb-Group Genes Encoding the HPC-HPH/PRC1 Complex in Clinically Defined Primary Nodal and Cutaneous Large B-Cell Lymphomas. American Journal of Pathology, 2004, 164, 533-542.	1.9	35
190	Bcl-2, Bcl-6 and CD10 expression in cutaneous B-cell lymphoma: further support for a follicle centre cell origin and differential diagnostic significance. British Journal of Dermatology, 2003, 149, 1183-1191.	1.4	155
191	EBV-positive cutaneous B-cell lymphoproliferative disease after imatinib mesylate. Blood, 2003, 102, 4243-4243.	0.6	34
192	Peripheral T-cell lymphomas unspecified presenting in the skin: analysis of prognostic factors in a group of 82 patients. Blood, 2003, 102, 2213-2219.	0.6	221
193	Differential Expression of Thymus and Activation Regulated Chemokine and Its Receptor CCR4 in Nodal and Cutaneous Anaplastic Large-Cell Lymphomas and Hodgkin's Disease. Modern Pathology, 2002, 15, 838-844.	2.9	38
194	Acute hemorrhagic edema of infancy (AHEI). Clinics in Dermatology, 2002, 20, 2-3.	0.8	25
195	A novel splice variant of the Fas gene in patients with cutaneous T-cell lymphoma. Cancer Research, 2002, 62, 5389-92.	0.4	48
196	Mouse fingers, a new computer-related skin disorder. Journal of the American Academy of Dermatology, 2001, 45, 477.	0.6	16
197	Should primary cutaneous Ki-1(CD30)-positive anaplastic large cell lymphoma in childhood be treated with multiple-agent chemotherapy?. Journal of the American Academy of Dermatology, 2001, 45, 638-639.	0.6	6
198	CD8+ T Cells in Cutaneous T-Cell Lymphoma: Expression of Cytotoxic Proteins, Fas Ligand, and Killing Inhibitory Receptors and Their Relationship With Clinical Behavior. Journal of Clinical Oncology, 2001, 19, 4322-4329.	0.8	105

#	ARTICLE	IF	CITATIONS
199	Absence of T(H)2 cytokine messenger RNA expression in CD30-negative primary cutaneous large T-cell lymphomas. <i>Archives of Dermatology</i> , 2001, 137, 901-5.	1.7	5
200	Expression of Fas and Fas-ligand in primary cutaneous T-cell lymphoma (CTCL): association between lack of Fas expression and aggressive types of CTCL. <i>British Journal of Dermatology</i> , 2000, 143, 313-319.	1.4	103
201	Treatment of Multifocal Primary Cutaneous B-Cell Lymphoma: A Clinical Follow-Up Study of 29 Patients. <i>Journal of Clinical Oncology</i> , 1999, 17, 2471-2471.	0.8	105
202	Primary Cutaneous CD8-Positive Epidermotropic Cytotoxic T Cell Lymphomas. <i>American Journal of Pathology</i> , 1999, 155, 483-492.	1.9	476
203	Expression of Cytotoxic Proteins by Neoplastic T Cells in Mycosis Fungoides Increases with Progression from Plaque Stage to Tumor Stage Disease. <i>American Journal of Pathology</i> , 1999, 154, 1203-1210.	1.9	82
204	Epidermal Interferon- γ Inducible Protein-10 (IP-10) and Monokine Induced by γ -Interferon (Mig) but not IL-8 mRNA Expression is Associated with Epidermotropism in Cutaneous T Cell Lymphomas. <i>Journal of Investigative Dermatology</i> , 1998, 111, 222-226.	0.3	67
205	Progression of mycosis fungoides is associated with expression of granzyme B and TIA-1 in neoplastic cells. <i>Journal of Dermatological Science</i> , 1998, 16, S53.	1.0	0
206	bcl-2 protein expression in primary cutaneous large B-cell lymphoma is site-related.. <i>Journal of Clinical Oncology</i> , 1998, 16, 2080-2085.	0.8	136
207	p53 and bcl-2 expression do not correlate with prognosis in primary cutaneous large T-cell lymphomas. <i>Journal of Cutaneous Pathology</i> , 1997, 24, 462-467.	0.7	20
208	Most Primary Cutaneous CD30-Positive Lymphoproliferative Disorders Have a CD4-Positive Cytotoxic T-Cell Phenotype. <i>Journal of Investigative Dermatology</i> , 1997, 109, 636-640.	0.3	131
209	Is mycosis fungoides exacerbated by fluoxetine?. <i>Journal of the American Academy of Dermatology</i> , 1996, 35, 635-636.	0.6	18
210	Primary Cutaneous Large B-Cell Lymphomas of the Legs. <i>Archives of Dermatology</i> , 1996, 132, 1304.	1.7	167
211	Effects of Ultraviolet B Light on Cutaneous Immune Responses of Humans with Deeply Pigmented Skin. <i>Journal of Investigative Dermatology</i> , 1991, 97, 729-734.	0.3	90
212	Functional Dichotomy between Langerhans Cells that Present Antigen to Naive and to Memory/Effector T Lymphocytes. <i>Immunological Reviews</i> , 1990, 117, 159-183.	2.8	111
213	Ultraviolet B light-induced alterations in epidermal Langerhans cells are mediated in part by tumor necrosis factor-alpha. <i>Photodermatology Photoimmunology and Photomedicine</i> , 1990, 7, 258-65.	0.7	101