

Tim Snijders

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5207323/publications.pdf>

Version: 2024-02-01

71
papers

4,721
citations

81743

39
h-index

98622

67
g-index

75
all docs

75
docs citations

75
times ranked

5078
citing authors

#	ARTICLE	IF	CITATIONS
1	Ingestion of an ample amount of meat substitute based on a lysine-enriched, plant-based protein blend stimulates postprandial muscle protein synthesis to a similar extent as an isonitrogenous amount of chicken in healthy, young men. <i>British Journal of Nutrition</i> , 2022, 128, 1955-1965.	1.2	12
2	Muscle fiber capillarization is associated with various indices of skeletal muscle mass in healthy, older men. <i>Experimental Gerontology</i> , 2021, 143, 111161.	1.2	13
3	Myonuclear content and domain size in small versus larger muscle fibres in response to 12 weeks of resistance exercise training in older adults. <i>Acta Physiologica</i> , 2021, 231, e13599.	1.8	15
4	No differences in muscle protein synthesis rates following ingestion of wheat protein, milk protein, and their protein blend in healthy, young males. <i>British Journal of Nutrition</i> , 2021, 126, 1832-1842.	1.2	34
5	Skeletal Muscle Adaptive Responses to Different Types of Short-Term Exercise Training and Detraining in Middle-Age Men. <i>Medicine and Science in Sports and Exercise</i> , 2021, 53, 2023-2036.	0.2	8
6	The Importance of Muscle Capillarization for Optimizing Satellite Cell Plasticity. <i>Exercise and Sport Sciences Reviews</i> , 2021, 49, 284-290.	1.6	17
7	The Anabolic Response to Plant-Based Protein Ingestion. <i>Sports Medicine</i> , 2021, 51, 59-74.	3.1	48
8	Moderate Intensity Exercise Training Improves Skeletal Muscle Performance in Asymptomatic and Asymptomatic Statin Users. <i>Journal of the American College of Cardiology</i> , 2021, 78, 2023-2037.	1.2	13
9	Brain-derived neurotrophic factor is associated with human muscle satellite cell differentiation in response to muscle-damaging exercise. <i>Applied Physiology, Nutrition and Metabolism</i> , 2020, 45, 581-590.	0.9	19
10	Capillary facilitation of skeletal muscle function in health and disease. <i>Applied Physiology, Nutrition and Metabolism</i> , 2020, 45, 453-462.	0.9	7
11	Age-related changes to the satellite cell niche are associated with reduced activation following exercise. <i>FASEB Journal</i> , 2020, 34, 8975-8989.	0.2	15
12	The concept of skeletal muscle memory: Evidence from animal and human studies. <i>Acta Physiologica</i> , 2020, 229, e13465.	1.8	52
13	Variability in skeletal muscle fibre characteristics during repeated muscle biopsy sampling in human vastus lateralis. <i>Applied Physiology, Nutrition and Metabolism</i> , 2020, 45, 368-375.	0.9	21
14	Consistent expression pattern of myogenic regulatory factors in whole muscle and isolated human muscle satellite cells after eccentric contractions in humans. <i>Journal of Applied Physiology</i> , 2019, 127, 1419-1426.	1.2	13
15	Nandrolone decanoate administration does not attenuate muscle atrophy during a short period of disuse. <i>PLoS ONE</i> , 2019, 14, e0210823.	1.1	8
16	A Multi-Ingredient Nutritional Supplement in Combination With Resistance Exercise and High-Intensity Interval Training Improves Cognitive Function and Increases N-3 Index in Healthy Older Men: A Randomized Controlled Trial. <i>Frontiers in Aging Neuroscience</i> , 2019, 11, 107.	1.7	14
17	Integrated Myofibrillar Protein Synthesis in Recovery From Unaccustomed and Accustomed Resistance Exercise With and Without Multi-ingredient Supplementation in Overweight Older Men. <i>Frontiers in Nutrition</i> , 2019, 6, 40.	1.6	14
18	Muscle mass and strength gains following 6 months of resistance type exercise training are only partly preserved within one year with autonomous exercise continuation in older adults. <i>Experimental Gerontology</i> , 2019, 121, 71-78.	1.2	41

#	ARTICLE	IF	CITATIONS
19	The Impact of Pre-sleep Protein Ingestion on the Skeletal Muscle Adaptive Response to Exercise in Humans: An Update. <i>Frontiers in Nutrition</i> , 2019, 6, 17.	1.6	45
20	Exercise training impacts skeletal muscle gene expression related to the kynurenine pathway. <i>American Journal of Physiology - Cell Physiology</i> , 2019, 316, C444-C448.	2.1	37
21	Prolonged exercise training improves the acute type II muscle fibre satellite cell response in healthy older men. <i>Journal of Physiology</i> , 2019, 597, 105-119.	1.3	45
22	The Impact of Aerobic Exercise on the Muscle Stem Cell Response. <i>Exercise and Sport Sciences Reviews</i> , 2018, 46, 180-187.	1.6	25
23	A multi-ingredient nutritional supplement enhances exercise training-related reductions in markers of systemic inflammation in healthy older men. <i>Applied Physiology, Nutrition and Metabolism</i> , 2018, 43, 299-302.	0.9	13
24	The influence of capillarization on satellite cell pool expansion and activation following exercise-induced muscle damage in healthy young men. <i>Journal of Physiology</i> , 2018, 596, 1063-1078.	1.3	50
25	Presleep dietary protein-derived amino acids are incorporated in myofibrillar protein during postexercise overnight recovery. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2018, 314, E457-E467.	1.8	56
26	Temporal Response of Angiogenesis and Hypertrophy to Resistance Training in Young Men. <i>Medicine and Science in Sports and Exercise</i> , 2018, 50, 36-45.	0.2	59
27	Leucine Supplementation Does Not Attenuate Skeletal Muscle Loss during Leg Immobilization in Healthy, Young Men. <i>Nutrients</i> , 2018, 10, 635.	1.7	37
28	Skeletal muscle fiber characteristics in patients with chronic heart failure: impact of disease severity and relation with muscle oxygenation during exercise. <i>Journal of Applied Physiology</i> , 2018, 125, 1266-1276.	1.2	11
29	Ingestion of a Multi-Ingredient Supplement Does Not Alter Exercise-Induced Satellite Cell Responses in Older Men. <i>Journal of Nutrition</i> , 2018, 148, 891-899.	1.3	13
30	Early- and later-phases satellite cell responses and myonuclear content with resistance training in young men. <i>PLoS ONE</i> , 2018, 13, e0191039.	1.1	42
31	The First Characterization of a Novel Stem Cell Population and the Temporal Relationship with Satellite Cells in Human Skeletal Muscle. <i>FASEB Journal</i> , 2018, 32, 615.2.	0.2	0
32	Muscle fiber capillarization as determining factor on indices of insulin sensitivity in humans. <i>Physiological Reports</i> , 2017, 5, e13278.	0.7	22
33	Extensive Type II Muscle Fiber Atrophy in Elderly Female Hip Fracture Patients. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2017, 72, 1369-1375.	1.7	50
34	Role of muscle stem cells in sarcopenia. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2017, 20, 186-190.	1.3	45
35	Intramyocellular lipid content and lipogenic gene expression responses following a single bout of resistance type exercise differ between young and older men. <i>Experimental Gerontology</i> , 2017, 93, 36-45.	1.2	12
36	Altered muscle satellite cell activation following 16 wk of resistance training in young men. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2017, 312, R85-R92.	0.9	45

#	ARTICLE	IF	CITATIONS
37	Creatine Loading Does Not Preserve Muscle Mass or Strength During Leg Immobilization in Healthy, Young Males: A Randomized Controlled Trial. <i>Sports Medicine</i> , 2017, 47, 1661-1671.	3.1	36
38	The Influence and Delivery of Cytokines and their Mediating Effect on Muscle Satellite Cells. <i>Current Stem Cell Reports</i> , 2017, 3, 192-201.	0.7	5
39	Skeletal Muscle Regeneration, Repair and Remodelling in Aging: The Importance of Muscle Stem Cells and Vascularization. <i>Gerontology</i> , 2017, 63, 91-100.	1.4	82
40	Muscle fibre capillarization is a critical factor in muscle fibre hypertrophy during resistance exercise training in older men. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2017, 8, 267-276.	2.9	114
41	A whey protein-based multi-ingredient nutritional supplement stimulates gains in lean body mass and strength in healthy older men: A randomized controlled trial. <i>PLoS ONE</i> , 2017, 12, e0181387.	1.1	87
42	Resistance training-induced changes in integrated myofibrillar protein synthesis are related to hypertrophy only after attenuation of muscle damage. <i>Journal of Physiology</i> , 2016, 594, 5209-5222.	1.3	236
43	Changes in myonuclear domain size do not precede muscle hypertrophy during prolonged resistance-type exercise training. <i>Acta Physiologica</i> , 2016, 216, 231-239.	1.8	50
44	Resistance Training Increases Skeletal Muscle Capillarization in Healthy Older Men. <i>Medicine and Science in Sports and Exercise</i> , 2016, 48, 2157-2164.	0.2	63
45	Skeletal muscle satellite cells are located at a closer proximity to capillaries in healthy young compared with older men. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2016, 7, 547-554.	2.9	91
46	Presleep protein ingestion does not compromise the muscle protein synthetic response to protein ingested the following morning. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 311, E964-E973.	1.8	30
47	Are satellite cells lost during short-term disuse-induced muscle fiber atrophy?. <i>Journal of Applied Physiology</i> , 2016, 120, 1490-1490.	1.2	5
48	Exercise conditioning in old mice improves skeletal muscle regeneration. <i>FASEB Journal</i> , 2016, 30, 3256-3268.	0.2	56
49	Short-term muscle disuse lowers myofibrillar protein synthesis rates and induces anabolic resistance to protein ingestion. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 310, E137-E147.	1.8	103
50	Satellite cells in human skeletal muscle plasticity. <i>Frontiers in Physiology</i> , 2015, 6, 283.	1.3	236
51	Ingestion of Casein in a Milk Matrix Modulates Dietary Protein Digestion and Absorption Kinetics but Does Not Modulate Postprandial Muscle Protein Synthesis in Older Men. <i>Journal of Nutrition</i> , 2015, 145, 1438-1445.	1.3	41
52	Protein Ingestion before Sleep Increases Muscle Mass and Strength Gains during Prolonged Resistance-Type Exercise Training in Healthy Young Men. <i>Journal of Nutrition</i> , 2015, 145, 1178-1184.	1.3	129
53	Myostatin inhibition for treatment of sarcopenia. <i>Lancet Diabetes and Endocrinology</i> , 2015, 3, 917-918.	5.5	12
54	Differences in postprandial protein handling after beef compared with milk ingestion during postexercise recovery: a randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2015, 102, 828-836.	2.2	99

#	ARTICLE	IF	CITATIONS
55	Short-term muscle disuse atrophy is not associated with increased intramuscular lipid deposition or a decline in the maximal activity of key mitochondrial enzymes in young and older males. <i>Experimental Gerontology</i> , 2015, 61, 76-83.	1.2	39
56	Acute Dietary Protein Intake Restriction Is Associated with Changes in Myostatin Expression after a Single Bout of Resistance Exercise in Healthy Young Men. <i>Journal of Nutrition</i> , 2014, 144, 137-145.	1.3	24
57	Muscle disuse atrophy is not accompanied by changes in skeletal muscle satellite cell content. <i>Clinical Science</i> , 2014, 126, 557-566.	1.8	55
58	Skeletal muscle capillary density and microvascular function are compromised with aging and type 2 diabetes. <i>Journal of Applied Physiology</i> , 2014, 116, 998-1005.	1.2	151
59	Satellite cells in human skeletal muscle; from birth to old age. <i>Age</i> , 2014, 36, 545-557.	3.0	280
60	Substantial skeletal muscle loss occurs during only 5 days of disuse. <i>Acta Physiologica</i> , 2014, 210, 600-611.	1.8	222
61	The skeletal muscle satellite cell response to a single bout of resistance-type exercise is delayed with aging in men. <i>Age</i> , 2014, 36, 9699.	3.0	87
62	Neuromuscular electrical stimulation prevents muscle disuse atrophy during leg immobilization in humans. <i>Acta Physiologica</i> , 2014, 210, 628-641.	1.8	177
63	The decline in skeletal muscle mass with aging is mainly attributed to a reduction in type II muscle fiber size. <i>Experimental Gerontology</i> , 2013, 48, 492-498.	1.2	522
64	Disuse Impairs the Muscle Protein Synthetic Response to Protein Ingestion in Healthy Men. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 4872-4881.	1.8	127
65	Eccentric Exercise Increases Satellite Cell Content in Type II Muscle Fibers. <i>Medicine and Science in Sports and Exercise</i> , 2013, 45, 230-237.	0.2	76
66	Neuromuscular electrical stimulation increases muscle protein synthesis in elderly type 2 diabetic men. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012, 303, E614-E623.	1.8	72
67	Reduced Satellite Cell Numbers with Spinal Cord Injury and Aging in Humans. <i>Medicine and Science in Sports and Exercise</i> , 2012, 44, 2322-2330.	0.2	82
68	A single bout of exercise activates skeletal muscle satellite cells during subsequent overnight recovery. <i>Experimental Physiology</i> , 2012, 97, 762-773.	0.9	51
69	Continuous endurance-type exercise training does not modulate satellite cell content in obese type 2 diabetes patients. <i>Muscle and Nerve</i> , 2011, 43, 393-401.	1.0	33
70	Characteristics of Muscle Fiber Type Are Predictive of Skeletal Muscle Mass and Strength in Elderly Men. <i>Journal of the American Geriatrics Society</i> , 2010, 58, 2069-2075.	1.3	86
71	The impact of sarcopenia and exercise training on skeletal muscle satellite cells. <i>Ageing Research Reviews</i> , 2009, 8, 328-338.	5.0	190