

Zhiguo Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5208757/publications.pdf>

Version: 2024-02-01

43
papers

1,201
citations

304743

22
h-index

377865

34
g-index

43
all docs

43
docs citations

43
times ranked

2216
citing authors

#	ARTICLE	IF	CITATIONS
1	Difference in the regulation of biological rhythm symptoms of Major depressive disorder between escitalopram and mirtazapine. <i>Journal of Affective Disorders</i> , 2022, 296, 258-264.	4.1	3
2	Evaluating the efficacy and moderators of algorithm-guided antidepressant treatments of major depressive disorder. <i>Journal of Affective Disorders</i> , 2022, 297, 68-75.	4.1	1
3	Lower Health Literacy of Mania Than Depression Among Older People: A Random Survey of a Community Healthcare Service Center. <i>Frontiers in Psychiatry</i> , 2021, 12, 512689.	2.6	2
4	Clinical features of the patients with major depressive disorder co-occurring insomnia and hypersomnia symptoms: a report of NSSD study. <i>Sleep Medicine</i> , 2021, 81, 375-381.	1.6	13
5	Predictors and moderators of quality of life in patients with major depressive disorder: An AGTs-MDD study report. <i>Journal of Psychiatric Research</i> , 2021, 138, 96-102.	3.1	5
6	Schizophrenia, bipolar disorder, or intracranial aneurysm? A case report. <i>Brain and Behavior</i> , 2021, 11, e2245.	2.2	2
7	Symptomatology differences of major depression in psychiatric versus general hospitals: A machine learning approach. <i>Journal of Affective Disorders</i> , 2020, 260, 349-360.	4.1	7
8	Causes of drug discontinuation in patients with major depressive disorder in China. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2020, 96, 109755.	4.8	23
9	Disagreement and factors between symptom on self-report and clinician rating of major depressive disorder: A report of a national survey in China. <i>Journal of Affective Disorders</i> , 2019, 253, 141-146.	4.1	10
10	Subtypes of treatment-resistant depression determined by a latent class analysis in a Chinese clinical population. <i>Journal of Affective Disorders</i> , 2019, 249, 82-89.	4.1	12
11	The association between somatic symptoms and suicidal ideation in Chinese first-episode major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 245, 17-21.	4.1	30
12	Advance in Diagnosis of Depressive Disorder. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1180, 179-191.	1.6	4
13	Prevalence, risk factors and clinical characteristics of suicidal ideation in Chinese patients with depression. <i>Journal of Affective Disorders</i> , 2018, 235, 135-141.	4.1	40
14	Somatic symptoms vary in major depressive disorder in China. <i>Comprehensive Psychiatry</i> , 2018, 87, 32-37.	3.1	37
15	The clinical correlates of comorbid anxiety symptoms and syndromal anxiety in patients with major depressive disorder. <i>Psychiatry Research</i> , 2018, 269, 251-257.	3.3	18
16	Different levels of pro- and anti-inflammatory cytokines in patients with unipolar and bipolar depression. <i>Journal of Affective Disorders</i> , 2018, 237, 65-72.	4.1	47
17	Reduced ENA78 levels as novel biomarker for major depressive disorder and venlafaxine efficiency: Result from a prospective longitudinal study. <i>Psychoneuroendocrinology</i> , 2017, 81, 113-121.	2.7	21
18	Identification of plasma biomarkers for distinguishing bipolar depression from major depressive disorder by iTRAQ-coupled LC-MS/MS and bioinformatics analysis. <i>Psychoneuroendocrinology</i> , 2017, 86, 17-24.	2.7	51

#	ARTICLE	IF	CITATIONS
19	Association analysis between mitogen-activated protein/extracellular signal-regulated kinase (MEK) gene polymorphisms and depressive disorder in the Han Chinese population. <i>Journal of Affective Disorders</i> , 2017, 222, 120-125.	4.1	6
20	Ratio of mBDNF to proBDNF for Differential Diagnosis of Major Depressive Disorder and Bipolar Depression. <i>Molecular Neurobiology</i> , 2017, 54, 5573-5582.	4.0	62
21	Evaluating the association between the SHANK3 gene and bipolar disorder. <i>Psychiatry Research</i> , 2016, 244, 284-288.	3.3	10
22	Identification of IL6 as a susceptibility gene for major depressive disorder. <i>Scientific Reports</i> , 2016, 6, 31264.	3.3	35
23	A haplotype in the 5'-upstream region of the NDUFV2 gene is associated with major depressive disorder in Han Chinese. <i>Journal of Affective Disorders</i> , 2016, 190, 329-332.	4.1	13
24	ZNF804A Genetic Variation Confers Risk to Bipolar Disorder. <i>Molecular Neurobiology</i> , 2016, 53, 2936-2943.	4.0	21
25	IL-23 and TGF- β 1 levels as potential predictive biomarkers in treatment of bipolar I disorder with acute manic episode. <i>Journal of Affective Disorders</i> , 2015, 174, 361-366.	4.1	50
26	Down-regulation of PRKCB1 expression in Han Chinese patients with subsyndromal symptomatic depression. <i>Journal of Psychiatric Research</i> , 2015, 69, 1-6.	3.1	6
27	Guidelines concordance of maintenance treatment in euthymic patients with bipolar disorder: Data from the national bipolar mania pathway survey (BIPAS) in mainland China. <i>Journal of Affective Disorders</i> , 2015, 182, 101-105.	4.1	6
28	Dissociated large-scale functional connectivity networks of the precuneus in medication-naïve first-episode depression. <i>Psychiatry Research - Neuroimaging</i> , 2015, 232, 250-256.	1.8	65
29	Validation of the Chinese Version of the Short TEMPS-A and its application in patients with mood disorders. <i>Journal of Affective Disorders</i> , 2015, 170, 178-184.	4.1	9
30	Significantly decreased mRNA levels of BDNF and MEK1 genes in treatment-resistant depression. <i>NeuroReport</i> , 2014, 25, 753-755.	1.2	32
31	A study of N-methyl-D-aspartate receptor gene (GRIN2B) variants as predictors of treatment-resistant major depression. <i>Psychopharmacology</i> , 2014, 231, 685-693.	3.1	65
32	Brain-derived neurotrophic factor levels and bipolar disorder in patients in their first depressive episode: 3-year prospective longitudinal study. <i>British Journal of Psychiatry</i> , 2014, 205, 29-35.	2.8	54
33	Influence of BCL2 gene in major depression susceptibility and antidepressant treatment outcome. <i>Journal of Affective Disorders</i> , 2014, 155, 288-294.	4.1	27
34	Guidelines Disconcordance in Acute Bipolar Depression: Data from the National Bipolar Mania Pathway Survey (BIPAS) in Mainland China. <i>PLoS ONE</i> , 2014, 9, e96096.	2.5	11
35	Comorbidity of depressive and anxiety disorders: challenges in diagnosis and assessment. <i>Shanghai Archives of Psychiatry</i> , 2014, 26, 227-31.	0.7	53
36	Influence of polymorphisms in genes SLC1A1, GRIN2B, and GRIK2 on clozapine-induced obsessive-compulsive symptoms. <i>Psychopharmacology</i> , 2013, 230, 49-55.	3.1	53

#	ARTICLE	IF	CITATIONS
37	Difference in remission in a Chinese population with anxious versus nonanxious treatment-resistant depression: A report of OPERATION study. <i>Journal of Affective Disorders</i> , 2013, 150, 834-839.	4.1	58
38	Association of genetic variation in CACNA1C with bipolar disorder in Han Chinese. <i>Journal of Affective Disorders</i> , 2013, 150, 261-265.	4.1	27
39	Association Study of Val66Met Polymorphism in Brain-Derived Neurotrophic Factor Gene with Clozapine-Induced Metabolic Syndrome: Preliminary Results. <i>PLoS ONE</i> , 2013, 8, e72652.	2.5	36
40	Number and characteristics of medical professionals working in Chinese mental health facilities. <i>Shanghai Archives of Psychiatry</i> , 2013, 25, 277-85.	0.7	39
41	A Pilot Study of the Efficacy and Safety of Paroxetine Augmented With Risperidone, Valproate, Buspirone, Trazodone, or Thyroid Hormone in Adult Chinese Patients With Treatment-Resistant Major Depression. <i>Journal of Clinical Psychopharmacology</i> , 2011, 31, 638-642.	1.4	47
42	Lack of effect of brain derived neurotrophic factor (BDNF) Val66Met polymorphism on early onset schizophrenia in Chinese Han population. <i>Brain Research</i> , 2011, 1417, 146-150.	2.2	42
43	Comparisons of the Efficacy and Tolerability of Extended-Release Venlafaxine, Mirtazapine, and Paroxetine in Treatment-Resistant Depression. <i>Journal of Clinical Psychopharmacology</i> , 2010, 30, 357-364.	1.4	48