

Adel Ahmed Elshafei

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/521240/publications.pdf>

Version: 2024-02-01

11
papers

90
citations

1684188

5
h-index

1474206

9
g-index

12
all docs

12
docs citations

12
times ranked

109
citing authors

#	ARTICLE	IF	CITATIONS
1	Assessment of the salt tolerance of wheat genotypes during the germination stage based on germination ability parameters and associated SSR markers. <i>Journal of Plant Interactions</i> , 2019, 14, 151-163.	2.1	39
2	Morphological variability and genetic diversity of wheat genotypes grown on saline soil and identification of new promising molecular markers associated with salinity tolerance. <i>Journal of Plant Interactions</i> , 2019, 14, 564-571.	2.1	13
3	Identification of new SSR markers linked to leaf chlorophyll content, flag leaf senescence and cell membrane stability traits in wheat under water stressed condition. <i>Acta Biologica Hungarica</i> , 2015, 66, 93-102.	0.7	12
4	The genetic basis of spectral reflectance indices in drought-stressed wheat. <i>Acta Physiologiae Plantarum</i> , 2016, 38, 1.	2.1	6
5	Doubled haploid wheat lines with high molecular weight glutenin alleles derived from microspore cultures. <i>New Zealand Journal of Crop and Horticultural Science</i> , 2018, 46, 198-211.	1.3	5
6	Molecular breeding for rust resistance in wheat genotypes. <i>Molecular Biology Reports</i> , 2021, 48, 731-742.	2.3	5
7	Regeneration of rice somaclones tolerant to high level of abscisic acid and their characterization via RAPD markers. <i>Bulletin of the National Research Centre</i> , 2019, 43, .	1.8	4
8	Phenotyping and validation of molecular markers associated with rust resistance genes in wheat cultivars in Egypt. <i>Molecular Biology Reports</i> , 2022, 49, 1903-1915.	2.3	3
9	Analysis of diversity using simple sequence repeat (SSR): distinctions between original <i>Parmentiera cereifera</i> tree and somaclones. <i>Bulletin of the National Research Centre</i> , 2018, 42, .	1.8	2
10	Genetic diversity of <i>Cucurbita moschata</i> inbred lines selected from six different populations using HFO-TAG markers. <i>Plant OMICS</i> , 2020, , 86-93.	0.4	1
11	Using high frequency oligonucleotides-targeting active gene (HFO-TAG) markers for genetic evaluation among genotypes (<i>Cucurbita pepo</i> L. and <i>C. maxima</i> L.). <i>Bulletin of the National Research Centre</i> , 2019, 43, .	1.8	0