

# Changhua Zhuo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5220134/publications.pdf>

Version: 2024-02-01

32  
papers

846  
citations

516710

16  
h-index

477307

29  
g-index

32  
all docs

32  
docs citations

32  
times ranked

1636  
citing authors

#	ARTICLE	IF	CITATIONS
1	Nuclear-enriched abundant transcript 1 as a diagnostic and prognostic biomarker in colorectal cancer. <i>Molecular Cancer</i> , 2015, 14, 191.	19.2	115
2	PIM1 Kinase Phosphorylates the Human Transcription Factor FOXP3 at Serine 422 to Negatively Regulate Its Activity under Inflammation. <i>Journal of Biological Chemistry</i> , 2014, 289, 26872-26881.	3.4	89
3	Low expression of novel lncRNA RP11-462C24.1 suggests a biomarker of poor prognosis in colorectal cancer. <i>Medical Oncology</i> , 2014, 31, 31.	2.5	68
4	Higher FOXP3-TSDR demethylation rates in adjacent normal tissues in patients with colon cancer were associated with worse survival. <i>Molecular Cancer</i> , 2014, 13, 153.	19.2	59
5	FOXP3+ Tregs: heterogeneous phenotypes and conflicting impacts on survival outcomes in patients with colorectal cancer. <i>Immunologic Research</i> , 2015, 61, 338-347.	2.9	57
6	Better Long-Term Survival in Young Patients with Non-Metastatic Colorectal Cancer after Surgery, an Analysis of 69,835 Patients in SEER Database. <i>PLoS ONE</i> , 2014, 9, e93756.	2.5	56
7	The influence of obesity on short- and long-term outcomes after laparoscopic surgery for colon cancer: A case-matched study of 152 patients. <i>Surgery</i> , 2014, 156, 661-668.	1.9	52
8	Chemokine (C-X-C motif) ligand 1 is associated with tumor progression and poor prognosis in patients with colorectal cancer. <i>Bioscience Reports</i> , 2018, 38, .	2.4	50
9	Overexpression of forkhead Box C2 promotes tumor metastasis and indicates poor prognosis in colon cancer via regulating epithelial-mesenchymal transition. <i>American Journal of Cancer Research</i> , 2015, 5, 2022-34.	1.4	34
10	Increased number of negative lymph nodes is associated with improved cancer specific survival in pathological IIIB and IIIC rectal cancer treated with preoperative radiotherapy. <i>Oncotarget</i> , 2014, 5, 12459-12471.	1.8	27
11	Experience of a specialist centre in the management of anastomotic sinus following leaks after low rectal or ileal pouch-anal anastomosis with diverting stoma. <i>Colorectal Disease</i> , 2013, 15, 1429-1435.	1.4	23
12	USP4 interacts and positively regulates IRF8 function via K48-linked deubiquitination in regulatory T cells. <i>FEBS Letters</i> , 2017, 591, 1677-1686.	2.8	23
13	Lymph Node Count After Preoperative Radiotherapy Is an Independently Prognostic Factor For Pathologically Lymph Node-Negative Patients With Rectal Cancer. <i>Medicine (United States)</i> , 2015, 94, e395.	1.0	21
14	TYMS serves as a prognostic indicator to predict the lymph node metastasis in Chinese patients with colorectal cancer. <i>Clinical Biochemistry</i> , 2013, 46, 1478-1483.	1.9	20
15	Unfavorable effect of small tumor size on cause-specific survival in stage IIA colon cancer, a SEER-based study. <i>International Journal of Colorectal Disease</i> , 2015, 30, 131-137.	2.2	20
16	Real-time noninvasive optical diagnosis for colorectal cancer using multiphoton microscopy. <i>Scanning</i> , 2012, 34, 181-185.	1.5	18
17	Laparoscopic Low Anterior Resection and Eversion Technique Combined With a Nondog Ear Anastomosis for Mid- and Distal Rectal Neoplasms. <i>Medicine (United States)</i> , 2015, 94, e2285.	1.0	17
18	Metastatic lymph node ratio can further stratify prognosis in rectal cancer patients treated with preoperative radiotherapy: a population-based analysis. <i>Tumor Biology</i> , 2014, 35, 6389-6395.	1.8	16

#	ARTICLE	IF	CITATIONS
19	Histological characteristics of collagen denaturation and injuries in bipolar radiofrequency-induced colonic anastomoses. <i>Pathology Research and Practice</i> , 2015, 211, 214-218.	2.3	15
20	Negative lymph node count is a significant prognostic factor in patient with stage IV gastric cancer after palliative gastrectomy. <i>Oncotarget</i> , 2017, 8, 71197-71205.	1.8	13
21	Pathological Features and Survival Outcomes of Young Patients with Operable Colon Cancer: Are They Homogeneous?. <i>PLoS ONE</i> , 2014, 9, e102004.	2.5	10
22	Downregulation of Activin A Receptor Type 2A Is Associated with Metastatic Potential and Poor Prognosis of Colon Cancer. <i>Journal of Cancer</i> , 2018, 9, 3626-3633.	2.5	10
23	Laparoscopic segmental colectomy for colonic lymphangiomas: A definitive, minimally invasive surgical option. <i>World Journal of Gastroenterology</i> , 2014, 20, 8745.	3.3	9
24	LINE-1 hypomethylation in normal colon mucosa is associated with poor survival in Chinese patients with sporadic colon cancer. <i>Oncotarget</i> , 2015, 6, 23820-23836.	1.8	7
25	Systematic Video Documentation in Laparoscopic Colon Surgery Using a Checklist: A Feasibility and Compliance Pilot Study. <i>Journal of Laparoendoscopic and Advanced Surgical Techniques - Part A</i> , 2015, 25, 737-743.	1.0	6
26	Association between the expression of carbonic anhydrase $\alpha$ 2II and clinicopathological features of hepatocellular carcinoma. <i>Oncology Letters</i> , 2019, 17, 5721-5728.	1.8	6
27	Feasibility and Safety of Laparoscopic Colon Surgery Performed Under Intravenous Sedation and Local Anesthesia Using Microinvasive (<3 mm) Instruments. <i>Surgical Innovation</i> , 2015, 22, 131-136.	0.9	2
28	Contribution of FOXC1 to the progression and metastasis and prognosis of human colon cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, 636-636.	1.6	2
29	Experience of a specialist center in the management of anastomotic sinus following leaks after low rectal or ileal pouch-anal anastomosis with diverting stoma. <i>Colorectal Disease</i> , 2014, 16, 565-565.	1.4	1
30	Optimization on cationic liposome-mediated cell transfection of plasmid DNA. <i>Chinese-German Journal of Clinical Oncology</i> , 2011, 10, 290-292.	0.1	0
31	Role of LINE-1 methylation status of adjacent normal colonic mucus in the survival of Chinese patients with colon cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, e14514-e14514.	1.6	0
32	Down expression of activin A receptor type 2A in relation to metastatic potential and prognosis of patients with colon cancer.. <i>Journal of Clinical Oncology</i> , 2018, 36, 712-712.	1.6	0