## Changhua Zhuo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5220134/publications.pdf

Version: 2024-02-01

516710 477307 32 846 16 29 citations g-index h-index papers 32 32 32 1636 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Association between the expression of carbonic anhydraseë;½II and clinicopathological features of hepatocellular carcinoma. Oncology Letters, 2019, 17, 5721-5728.	1.8	6
2	Downregulation of Activin A Receptor Type 2A Is Associated with Metastatic Potential and Poor Prognosis of Colon Cancer. Journal of Cancer, 2018, 9, 3626-3633.	2.5	10
3	Chemokine (C-X-C motif) ligand $1$ is associated with tumor progression and poor prognosis in patients with colorectal cancer. Bioscience Reports, 2018, 38, .	2.4	50
4	Down expression of activin A receptor type 2A in relation to metastatic potential and prognosis of patients with colon cancer Journal of Clinical Oncology, 2018, 36, 712-712.	1.6	0
5	<scp>USP</scp> 4 interacts and positively regulates <scp>IRF</scp> 8 function via K48â€linked deubiquitination in regulatory T cells. FEBS Letters, 2017, 591, 1677-1686.	2.8	23
6	Negative lymph node count is a significant prognostic factor in patient with stage IV gastric cancer after palliative gastrectomy. Oncotarget, 2017, 8, 71197-71205.	1.8	13
7	Laparoscopic Low Anterior Resection and Eversion Technique Combined With a Nondog Ear Anastomosis for Mid- and Distal Rectal Neoplasms. Medicine (United States), 2015, 94, e2285.	1.0	17
8	Nuclear-enriched abundant transcript $1$ as a diagnostic and prognostic biomarker in colorectal cancer. Molecular Cancer, 2015, 14, 191.	19.2	115
9	FOXP3+ Tregs: heterogeneous phenotypes and conflicting impacts on survival outcomes in patients with colorectal cancer. Immunologic Research, 2015, 61, 338-347.	2.9	57
10	Histological characteristics of collagen denaturation and injuries in bipolar radiofrequency-induced colonic anastomoses. Pathology Research and Practice, 2015, 211, 214-218.	2.3	15
11	Unfavorable effect of small tumor size on cause-specific survival in stage IIA colon cancer, a SEER-based study. International Journal of Colorectal Disease, 2015, 30, 131-137.	2.2	20
12	Lymph Node Count After Preoperative Radiotherapy Is an Independently Prognostic Factor For Pathologically Lymph Node-Negative Patients With Rectal Cancer. Medicine (United States), 2015, 94, e395.	1.0	21
13	Systematic Video Documentation in Laparoscopic Colon Surgery Using a Checklist: A Feasibility and Compliance Pilot Study. Journal of Laparoendoscopic and Advanced Surgical Techniques - Part A, 2015, 25, 737-743.	1.0	6
14	Feasibility and Safety of Laparoscopic Colon Surgery Performed Under Intravenous Sedation and Local Anesthesia Using Microinvasive (<3 mm) Instruments. Surgical Innovation, 2015, 22, 131-136.	0.9	2
15	Contribution of FOXC1 to the progression and metastasis and prognosis of human colon cancer Journal of Clinical Oncology, 2015, 33, 636-636.	1.6	2
16	LINE-1 hypomethylation in normal colon mucosa is associated with poor survival in Chinese patients with sporadic colon cancer. Oncotarget, 2015, 6, 23820-23836.	1.8	7
17	Role of LINE-1 methylation status of adjacent normal colonic mucus in the survival of Chinese patients with colon cancer Journal of Clinical Oncology, 2015, 33, e14514-e14514.	1.6	0
18	Overexpression of forkhead Box C2 promotes tumor metastasis and indicates poor prognosis in colon cancer via regulating epithelial-mesenchymal transition. American Journal of Cancer Research, 2015, 5, 2022-34.	1.4	34

#	Article	IF	CITATIONS
19	Better Long-Term Survival in Young Patients with Non-Metastatic Colorectal Cancer after Surgery, an Analysis of 69,835 Patients in SEER Database. PLoS ONE, 2014, 9, e93756.	2.5	56
20	Pathological Features and Survival Outcomes of Young Patients with Operable Colon Cancer: Are They Homogeneous?. PLoS ONE, 2014, 9, e102004.	2.5	10
21	Increased number of negative lymph nodes is associated with improved cancer specific survival in pathological IIIB and IIIC rectal cancer treated with preoperative radiotherapy. Oncotarget, 2014, 5, 12459-12471.	1.8	27
22	Experience of a specialist center in the management of anastomotic sinus following leaks after low rectal or ileal pouch-anal anastomosis with diverting stoma. Colorectal Disease, 2014, 16, 565-565.	1.4	1
23	The influence of obesity on short- and long-term outcomes after laparoscopic surgery for colon cancer: A case-matched study of 152 patients. Surgery, 2014, 156, 661-668.	1.9	52
24	PIM1 Kinase Phosphorylates the Human Transcription Factor FOXP3 at Serine 422 to Negatively Regulate Its Activity under Inflammation. Journal of Biological Chemistry, 2014, 289, 26872-26881.	3.4	89
25	Higher FOXP3-TSDR demethylation rates in adjacent normal tissues in patients with colon cancer were associated with worse survival. Molecular Cancer, 2014, 13, 153.	19.2	59
26	Low expression of novel lncRNA RP11-462C24.1 suggests a biomarker of poor prognosis in colorectal cancer. Medical Oncology, 2014, 31, 31.	2.5	68
27	Metastatic lymph node ratio can further stratify prognosis in rectal cancer patients treated with preoperative radiotherapy: a population-based analysis. Tumor Biology, 2014, 35, 6389-6395.	1.8	16
28	Laparoscopic segmental colectomy for colonic lymphangiomas: A definitive, minimally invasive surgical option. World Journal of Gastroenterology, 2014, 20, 8745.	3.3	9
29	TYMS serves as a prognostic indicator to predict the lymph node metastasis in Chinese patients with colorectal cancer. Clinical Biochemistry, 2013, 46, 1478-1483.	1.9	20
30	Experience of a specialist centre in the management of anastomotic sinus following leaks after low rectal or ileal pouch–anal anastomosis with diverting stoma. Colorectal Disease, 2013, 15, 1429-1435.	1.4	23
31	Realâ€time noninvasive optical diagnosis for colorectal cancer using multiphoton microscopy. Scanning, 2012, 34, 181-185.	1.5	18
32	Optimization on cationic liposome-mediated cell transfection of plasmid DNA. Chinese-German Journal of Clinical Oncology, 2011, 10, 290-292.	0.1	0