Alexander Presciutti

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5225213/publications.pdf

Version: 2024-02-01

933447 996975 19 241 10 15 citations g-index h-index papers 19 19 19 287 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Determinants of Long-Term Neurological Recovery Patterns Relative to Hospital Discharge Among Cardiac Arrest Survivors. Critical Care Medicine, 2018, 46, e141-e150.	0.9	29
2	Cardiac Arrest and Subsequent Hospitalization–Induced Posttraumatic Stress Is Associated With 1-Year Risk of Major Adverse Cardiovascular Events and All-Cause Mortality. Critical Care Medicine, 2019, 47, e502-e505.	0.9	23
3	Women have worse cognitive, functional, and psychiatric outcomes at hospital discharge after cardiac arrest. Resuscitation, 2018, 125, 12-15.	3.0	22
4	Posttraumatic stress and depressive symptoms characterize cardiac arrest survivors' perceived recovery at hospital discharge. General Hospital Psychiatry, 2018, 53, 108-113.	2.4	22
5	The impact of psychological distress on long-term recovery perceptions in survivors of cardiac arrest. Journal of Critical Care, 2019, 50, 227-233.	2.2	22
6	The Impact of Resilience Factors and Anxiety During Hospital Admission on Longitudinal Anxiety Among Dyads of Neurocritical Care Patients Without Major Cognitive Impairment and Their Family Caregivers. Neurocritical Care, 2020, 33, 468-478.	2.4	21
7	Early myoclonus following anoxic brain injury. Neurology: Clinical Practice, 2018, 8, 249-256.	1.6	18
8	Associations between posttraumatic stress symptoms and quality of life in cardiac arrest survivors and informal caregivers: A pilot survey study. Resuscitation Plus, 2021, 5, 100085.	1.7	17
9	Qualitative Study of Longâ€∓erm Cardiac Arrest Survivors' Challenges and Recommendations for Improving Survivorship. Journal of the American Heart Association, 2022, 11, .	3.7	13
10	Hyperarousal Symptoms in Survivors of Cardiac Arrest Are Associated With 13 Month Risk of Major Adverse Cardiovascular Events and All-Cause Mortality. Annals of Behavioral Medicine, 2020, 54, 413-422.	2.9	12
11	Dimensional structure of posttraumatic stress disorder symptoms after cardiac arrest. Journal of Affective Disorders, 2019, 251, 213-217.	4.1	7
12	Gaps in the Provision of Cognitive and Psychological Resources in Cardiac Arrest Survivors with Good Neurologic Recovery. Therapeutic Hypothermia and Temperature Management, 2022, 12, 61-67.	0.9	7
13	The evolution of hypothermia for neuroprotection after cardiac arrest: a history in the making. Annals of the New York Academy of Sciences, 2022, 1507, 60-69.	3.8	6
14	Associations between depression and anxiety symptoms with quality of life in cardiac arrest survivors with good neurologic recovery and informal caregivers of cardiac arrest survivors. Journal of Affective Disorders Reports, 2020, 2, 100046.	1.7	6
15	Associations Between Baseline Total PTSD Symptom Severity, Specific PTSD Symptoms, and 3-Month Quality of Life in Neurologically Intact Neurocritical Care Patients and Informal Caregivers. Neurocritical Care, 2021, 34, 54-63.	2.4	5
16	Modifiable provider-patient relationship factors and illness perceptions are associated with quality of life in survivors of cardiac arrest with good neurologic recovery. Resuscitation Plus, 2020, 3, 100008.	1.7	4
17	Abstract 302: NeuroCardiac Comprehensive Care Clinic Seeks to Define and Detect Neurological, Psychiatric, and Functional Sequelae in Cardiac Arrest Survivors. Circulation, 2018, 138, .	1.6	3
18	Tracheostomy use, long-term survival, and neurological outcomes among cardiac arrest survivors. Resuscitation, 2018, 129, e19-e20.	3.0	2

#	Article	IF	CITATIONS
19	Mindfulness is inversely associated with psychological symptoms in long-term cardiac arrest survivors. Journal of Behavioral Medicine, 2022, , 1.	2.1	2