

Hironori Masuko

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5228131/publications.pdf>

Version: 2024-02-01

27
papers

820
citations

759233

12
h-index

526287

27
g-index

28
all docs

28
docs citations

28
times ranked

1640
citing authors

#	ARTICLE	IF	CITATIONS
1	Common exacerbation-prone phenotypes across asthma and chronic obstructive pulmonary disease (COPD). <i>PLoS ONE</i> , 2022, 17, e0264397.	2.5	4
2	Identification of whole blood gene expressions correlated with responsiveness to benralizumab. <i>Journal of Allergy and Clinical Immunology</i> , 2021, 147, 772-775.	2.9	14
3	ORMDL3/GSDMB genotype as a risk factor for early-onset adult asthma is linked to total serum IgE levels but not to allergic sensitization. <i>Allergology International</i> , 2021, 70, 55-60.	3.3	12
4	Dust mite-dominant sensitization pattern as a causal factor for adult-onset asthma. <i>Allergology International</i> , 2021, 70, 368-369.	3.3	0
5	Identification of distinct phenotypes related to benralizumab responsiveness in patients with severe eosinophilic asthma. <i>PLoS ONE</i> , 2021, 16, e0248305.	2.5	12
6	Expression quantitative trait loci for ETV4 and MEOX1 are associated with adult asthma in Japanese populations. <i>Scientific Reports</i> , 2021, 11, 18791.	3.3	2
7	ORMDL3/GSDMB genotype is associated with distinct phenotypes of adult asthma. <i>Allergology International</i> , 2021, 70, 495-497.	3.3	3
8	Effects of Lung Function-Related Genes and <i>TSLP</i> on COPD Phenotypes. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2020, 17, 59-64.	1.6	5
9	Genetic impact of <i>CDHR3</i> on the adult onset of asthma and COPD. <i>Clinical and Experimental Allergy</i> , 2020, 50, 1223-1229.	2.9	10
10	High-Yield Production of the Major Birch Pollen Allergen Bet v 1 With Allergen Immunogenicity in <i>Nicotiana benthamiana</i> . <i>Frontiers in Plant Science</i> , 2020, 11, 344.	3.6	13
11	Association analyses of eQTLs of the TYRO3 gene and allergic diseases in Japanese populations. <i>Allergology International</i> , 2019, 68, 77-81.	3.3	8
12	The Concavity of the Maximal Expiratory Flow–Volume Curve Reflects the Extent of Emphysema in Obstructive Lung Diseases. <i>Scientific Reports</i> , 2019, 9, 13159.	3.3	4
13	A cis-eQTL allele regulating reduced expression of CHI3L1 is associated with late-onset adult asthma in Japanese cohorts. <i>BMC Medical Genetics</i> , 2019, 20, 58.	2.1	6
14	How important is allergic sensitization as a cause of atopic asthma?. <i>Allergology International</i> , 2018, 67, 292-294.	3.3	4
15	Genetic association of the functional CDHR3 genotype with early-onset adult asthma in Japanese populations. <i>Allergology International</i> , 2017, 66, 563-567.	3.3	24
16	Role of Lung Function Genes in the Development of Asthma. <i>PLoS ONE</i> , 2016, 11, e0145832.	2.5	13
17	Variants near the HLA complex group 22 gene (HCG22) confer increased susceptibility to late-onset asthma in Japanese populations. <i>Journal of Allergy and Clinical Immunology</i> , 2016, 138, 281-283.e13.	2.9	28
18	Heritability of pulmonary function estimated from genome-wide SNPs in healthy Japanese adults. <i>Respiratory Investigation</i> , 2015, 53, 60-67.	1.8	10

#	ARTICLE	IF	CITATIONS
19	Influence of MILR1 promoter polymorphism on expression levels and the phenotype of atopy. <i>Journal of Human Genetics</i> , 2014, 59, 480-483.	2.3	2
20	A Distinct Sensitization Pattern Associated with Asthma and the Thymic Stromal Lymphopoietin (TSLP) Genotype. <i>Allergology International</i> , 2013, 62, 123-130.	3.3	13
21	Genome-Wide Association Study for Levels of Total Serum IgE Identifies HLA-C in a Japanese Population. <i>PLoS ONE</i> , 2013, 8, e80941.	2.5	40
22	Asthma Phenotypes in Japanese Adults - Their Associations with the CCL5 ADRB2 Genotypes. <i>Allergology International</i> , 2013, 62, 113-121.	3.3	41
23	Genome-wide association study identifies three new susceptibility loci for adult asthma in the Japanese population. <i>Nature Genetics</i> , 2011, 43, 893-896.	21.4	296
24	Lower FEV1 in non-COPD, nonasthmatic subjects: association with smoking, annual decline in FEV1, total IgE levels, and TSLP genotypes. <i>International Journal of COPD</i> , 2011, 6, 181.	2.3	31
25	An interaction between Nrf2 polymorphisms and smoking status affects annual decline in FEV1: a longitudinal retrospective cohort study. <i>BMC Medical Genetics</i> , 2011, 12, 97.	2.1	33
26	Thymic Stromal Lymphopoietin Gene Promoter Polymorphisms Are Associated with Susceptibility to Bronchial Asthma. <i>American Journal of Respiratory Cell and Molecular Biology</i> , 2011, 44, 787-793.	2.9	187
27	Pulmonary Sequestration with Elevated Serum Level of Progastrin-releasing Peptide. <i>Internal Medicine</i> , 2008, 47, 157-160.	0.7	4