

Xiaomu Kong

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5229418/publications.pdf>

Version: 2024-02-01

24
papers

522
citations

759190

12
h-index

677123

22
g-index

26
all docs

26
docs citations

26
times ranked

1293
citing authors

#	ARTICLE	IF	CITATIONS
1	Association of AGTR1 A1166C and CYP2C9*3 Gene Polymorphisms with the Antihypertensive Effect of Valsartan. International Journal of Hypertension, 2022, 2022, 1-7.	1.3	7
2	A Novel Stool Methylation Test for the Non-Invasive Screening of Gastric and Colorectal Cancer. Frontiers in Oncology, 2022, 12, 860701.	2.8	4
3	Maternal and paternal histories differentially influence risks for diabetes, insulin secretion and insulin resistance in a Chinese population. Journal of Diabetes Investigation, 2021, 12, 434-445.	2.4	7
4	Acarbose Reduces Low-Grade Albuminuria Compared to Metformin in Chinese Patients with Newly Diagnosed Type 2 Diabetes. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2021, Volume 14, 4451-4458.	2.4	3
5	Early-onset type 2 diabetes is associated with genetic variants of β -cell function in the Chinese Han population. Diabetes/Metabolism Research and Reviews, 2020, 36, e3214.	4.0	14
6	Glucose outcomes of a learning-type artificial pancreas with an unannounced meal in type 1 diabetes. Computer Methods and Programs in Biomedicine, 2020, 191, 105416.	4.7	10
7	Genotypic and phenotypic spectra of hemojuvelin mutations in primary hemochromatosis patients: a systematic review. Orphanet Journal of Rare Diseases, 2019, 14, 171.	2.7	20
8	Sexual Dimorphism of a Genetic Risk Score for Obesity and Related Traits among Chinese Patients with Type 2 Diabetes. Obesity Facts, 2019, 12, 328-343.	3.4	7
9	Membrane-Spanning Protein Genetic Polymorphisms Related to Methotrexate Therapeutic Outcomes in a Chinese Rheumatoid Arthritis Population. Journal of Clinical Pharmacology, 2019, 59, 1471-1476.	2.0	4
10	Interaction of MTHFR C677T polymorphism with smoking in susceptibility to diabetic nephropathy in Chinese men with type 2 diabetes. Journal of Human Genetics, 2019, 64, 23-28.	2.3	9
11	Association of <i>EPHX2</i> R287Q Polymorphism with Diabetic Nephropathy in Chinese Type 2 Diabetic Patients. Journal of Diabetes Research, 2018, 2018, 1-6.	2.3	10
12	Diabetes remission and relapse after metabolic surgery. Journal of Diabetes Investigation, 2018, 9, 1237-1238.	2.4	14
13	Synergistic Effect of the <i>MTHFR</i> C677T and <i>EPHX2</i> G860A Polymorphism on the Increased Risk of Ischemic Stroke in Chinese Type 2 Diabetic Patients. Journal of Diabetes Research, 2017, 2017, 1-8.	2.3	19
14	Genetic variants associated with lean and obese type 2 diabetes in a Han Chinese population. Medicine (United States), 2016, 95, e3841.	1.0	24
15	Mobile Phone Intervention and Weight Loss Among Overweight and Obese Adults: A Meta-Analysis of Randomized Controlled Trials. American Journal of Epidemiology, 2015, 181, 337-348.	3.4	90
16	Genetic Variants Associated with Lipid Profiles in Chinese Patients with Type 2 Diabetes. PLoS ONE, 2015, 10, e0135145.	2.5	12
17	The Association of Type 2 Diabetes Loci Identified in Genome-Wide Association Studies with Metabolic Syndrome and Its Components in a Chinese Population with Type 2 Diabetes. PLoS ONE, 2015, 10, e0143607.	2.5	41
18	Cross-Sectional and Longitudinal Replication Analyses of Genome-Wide Association Loci of Type 2 Diabetes in Han Chinese. PLoS ONE, 2014, 9, e91790.	2.5	17

#	ARTICLE	IF	CITATIONS
19	Obesity-Related Genomic Loci Are Associated with Type 2 Diabetes in a Han Chinese Population. PLoS ONE, 2014, 9, e104486.	2.5	25
20	Liver X receptor activation increases hepatic fatty acid desaturation by the induction of SCD1 expression through an LXR-dependent mechanism (Xia et al., 2014). Journal of Diabetes Research, 2014, 2014, 1-10.	1.8	35
21	Thyroid hormone-responsive SPOT 14 homolog promotes hepatic lipogenesis, and its expression is regulated by Liver X receptor 1 through a sterol regulatory element-binding protein 1c-dependent mechanism in mice. Hepatology, 2013, 58, 617-628.	7.3	72
22	Association of Genetic Variants with Isolated Fasting Hyperglycaemia and Isolated Postprandial Hyperglycaemia in a Han Chinese Population. PLoS ONE, 2013, 8, e71399.	2.5	10
23	Peroxisome proliferator-activated receptor-1 is renoprotective in doxorubicin-induced glomerular injury. Kidney International, 2011, 79, 1302-1311.	5.2	55
24	Rapid detection of the irinotecan-related UGT1A1 *28 polymorphism by asymmetric PCR melting curve analysis using one fluorescent probe. Journal of Clinical Laboratory Analysis, 2009, 24, 1-6.	2.1	2