Katharina Sunnerhagen

List of Publications by Citations

Source: https://exaly.com/author-pdf/5238967/katharina-sunnerhagen-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

10,198 321 51 90 h-index g-index citations papers 12,070 339

ext. papers

ext. citations

3.7 avg, IF

6.59 L-index

#	Paper	IF	Citations
321	Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. <i>Cerebrovascular Diseases</i> , 2008 , 25, 457-507	3.2	1902
320	Person-centered careready for prime time. European Journal of Cardiovascular Nursing, 2011, 10, 248-5	53 .3	841
319	Walking training of patients with hemiparesis at an early stage after stroke: a comparison of walking training on a treadmill with body weight support and walking training on the ground. <i>Clinical Rehabilitation</i> , 2001 , 15, 515-27	3.3	192
318	Kinematic variables quantifying upper-extremity performance after stroke during reaching and drinking from a glass. <i>Neurorehabilitation and Neural Repair</i> , 2011 , 25, 71-80	4.7	178
317	Serum levels of the brain-derived proteins S-100 and NSE predict long-term outcome after cardiac arrest. <i>Resuscitation</i> , 2001 , 49, 183-91	4	158
316	Action Plan for Stroke in Europe 2018-2030. European Stroke Journal, 2018, 3, 309-336	5.6	144
315	Evidence of impaired adipogenesis in insulin resistance. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 317, 1045-51	3.4	137
314	Evidence-based stroke r-ehabilitation: an expanded guidance document from the european stroke organisation (ESO) guidelines for management of ischaemic stroke and transient ischaemic attack 2008. <i>Journal of Rehabilitation Medicine</i> , 2009 , 41, 99-111	3.4	130
313	Virtual reality and haptics as a training device for movement rehabilitation after stroke: a single-case study. <i>Archives of Physical Medicine and Rehabilitation</i> , 2004 , 85, 1247-50	2.8	121
312	Two years of growth hormone (GH) treatment increase isometric and isokinetic muscle strength in GH-deficient adults. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997 , 82, 2877-84	5.6	111
311	Two Years of Growth Hormone (GH) Treatment Increase Isometric and Isokinetic Muscle Strength in GH-Deficient Adults. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997 , 82, 2877-2884	5.6	101
310	Virtual rehabilitation in an activity centre for community-dwelling persons with stroke. The possibilities of 3-dimensional computer games. <i>Cerebrovascular Diseases</i> , 2008 , 26, 289-96	3.2	99
309	Intravenous immunoglobulin for post-polio syndrome: a randomised controlled trial. <i>Lancet Neurology, The</i> , 2006 , 5, 493-500	24.1	93
308	Recovery after stroke: cognition, ADL function and return to work. <i>Acta Neurologica Scandinavica</i> , 2007 , 115, 73-80	3.8	90
307	Baseline characteristics and the effects of five years of GH replacement therapy in adults with GH deficiency of childhood or adulthood onset: a comparative, prospective study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 4693-9	5.6	88
306	Energy expenditure in stroke subjects walking with a carbon composite ankle foot orthosis. <i>Journal of Rehabilitation Medicine</i> , 2004 , 36, 165-8	3.4	87
305	Survivors of out of hospital cardiac arrest: their prognosis, longevity and functional status. <i>Resuscitation</i> , 1997 , 35, 117-21	4	86

(2006-1998)

304	An 8-year longitudinal study of muscle strength, muscle fiber size, and dynamic electromyogram in individuals with late polio. <i>Muscle and Nerve</i> , 1998 , 21, 1428-37	3.4	81	
303	Oxygen consumption during treadmill walking with and without body weight support in patients with hemiparesis after stroke and in healthy subjects. <i>Archives of Physical Medicine and Rehabilitation</i> , 2000 , 81, 953-7	2.8	81	
302	Five years of growth hormone replacement therapy in adults: age- and gender-related changes in isometric and isokinetic muscle strength. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 20	67:69	79	
301	The first year after successful cardiac resuscitation: function, activity, participation and quality of life. <i>Resuscitation</i> , 2005 , 66, 285-9	4	78	
300	Dynamic water exercise in individuals with late poliomyelitis. <i>Archives of Physical Medicine and Rehabilitation</i> , 2001 , 82, 66-72	2.8	73	
299	Poststroke chronic disease management: towards improved identification and interventions for poststroke spasticity-related complications. <i>International Journal of Stroke</i> , 2011 , 6, 42-6	6.3	72	
298	Measurement of energy cost by the physiological cost index in walking after stroke. <i>Archives of Physical Medicine and Rehabilitation</i> , 2007 , 88, 1298-303	2.8	71	
297	Development, expansion, and use of a stroke clinical trials resource for novel exploratory analyses. <i>International Journal of Stroke</i> , 2012 , 7, 133-8	6.3	70	
296	Hydrotherapya new approach to improve function in the older patient with chronic heart failure. <i>European Journal of Heart Failure</i> , 2003 , 5, 527-35	12.3	70	
295	A home-based, self-administered stimulation program to improve selected hand functions of chronic stroke. <i>NeuroRehabilitation</i> , 2003 , 18, 215-225	2	70	
294	Improving the development, monitoring and reporting of stroke rehabilitation research: Consensus-based core recommendations from the Stroke Recovery and Rehabilitation Roundtable. <i>International Journal of Stroke</i> , 2017 , 12, 472-479	6.3	69	
293	Clinical tests performed in acute stroke identify the risk of falling during the first year: postural stroke study in Gothenburg (POSTGOT). <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 348-53	3.4	69	
292	Anxiety after stroke: A systematic review and meta-analysis. <i>Journal of Rehabilitation Medicine</i> , 2018 , 50, 769-778	3.4	69	
291	Can computerized working memory training improve impaired working memory, cognition and psychological health?. <i>Brain Injury</i> , 2013 , 27, 1649-57	2.1	66	
290	MUSCLE PERFORMANCE IN AN URBAN POPULATION SAMPLE OF 40- TO 79-YEAR-OLD MEN AND WOMEN. <i>Journal of Rehabilitation Medicine</i> , 2000 , 32, 159-167		66	
289	Navigating the poststroke continuum of care. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2013 , 22, 1-8	2.8	65	
288	Assessing and treating functional impairment in poststroke spasticity. <i>Neurology</i> , 2013 , 80, S35-44	6.5	65	
287	Three-dimensional kinematic motion analysis of a daily activity drinking from a glass: a pilot study. Journal of NeuroEngineering and Rehabilitation, 2006, 3, 18	5.3	65	

286	Consensus statements and recommendations from the ESO-Karolinska Stroke Update Conference, Stockholm 11-13 November 2018. <i>European Stroke Journal</i> , 2019 , 4, 307-317	5.6	63
285	Outcome and upper extremity function within 72 hours after first occasion of stroke in an unselected population at a stroke unit. A part of the SALGOT study. <i>BMC Neurology</i> , 2012 , 12, 162	3.1	63
284	Assessment and training in a 3-dimensional virtual environment with haptics: a report on 5 cases of motor rehabilitation in the chronic stage after stroke. <i>Neurorehabilitation and Neural Repair</i> , 2007 , 21, 180-9	4.7	63
283	Responsiveness of upper extremity kinematic measures and clinical improvement during the first three months after stroke. <i>Neurorehabilitation and Neural Repair</i> , 2013 , 27, 844-53	4.7	61
282	Stroke in women - from evidence to inequalities. <i>Nature Reviews Neurology</i> , 2017 , 13, 521-532	15	61
281	Life after cardiac arrest; a retrospective study. <i>Resuscitation</i> , 1996 , 31, 135-40	4	61
280	Ten years of growth hormone (GH) replacement normalizes muscle strength in GH-deficient adults. Journal of Clinical Endocrinology and Metabolism, 2009 , 94, 809-16	5.6	59
279	Movement kinematics during a drinking task are associated with the activity capacity level after stroke. <i>Neurorehabilitation and Neural Repair</i> , 2012 , 26, 1106-15	4.7	59
278	Scoring alternatives for FIM in neurological disorders applying Rasch analysis. <i>Acta Neurologica Scandinavica</i> , 2005 , 111, 264-73	3.8	56
277	Fear of falling, balance, and gait velocity in patients with stroke. <i>Physiotherapy Theory and Practice</i> , 2005 , 21, 113-20	1.5	54
276	Safety and efficacy of fluoxetine on functional recovery after acute stroke (EFFECTS): a randomised, double-blind, placebo-controlled trial. <i>Lancet Neurology, The</i> , 2020 , 19, 661-669	24.1	54
275	Upper-limb spasticity during the first year after stroke: stroke arm longitudinal study at the University of Gothenburg. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2014 , 93, 884-96	2.6	53
274	Factors associated with health-related quality of life after stroke: a 1-year prospective cohort study. <i>Neurorehabilitation and Neural Repair</i> , 2012 , 26, 266-74	4.7	53
273	Anxiety, depression, and psychological well-being 2 to 5 years poststroke. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2010 , 19, 364-9	2.8	52
272	Chronic pain and sex-differences; women accept and move, while men feel blue. <i>PLoS ONE</i> , 2017 , 12, e0175737	3.7	52
271	Return to Work after a Stroke in Working Age Persons; A Six-Year Follow Up. <i>PLoS ONE</i> , 2017 , 12, e016	935 , 9	51
270	Post-stroke spasticity: predictors of early development and considerations for therapeutic intervention. <i>PM and R</i> , 2015 , 7, 60-7	2.2	50
269	Skeletal muscle characteristics, muscle strength and thigh muscle area in patients before and after cardiac transplantation. <i>European Journal of Heart Failure</i> , 2001 , 3, 59-67	12.3	50

(2010-2015)

268	Early prediction of long-term upper limb spasticity after stroke: part of the SALGOT study. <i>Neurology</i> , 2015 , 85, 873-80	6.5	49	
267	The functional independence measure in Sweden: experience for outcome measurement in rehabilitation medicine. <i>Journal of Rehabilitation Medicine</i> , 1996 , 28, 51-62		49	
266	Functioning of stroke survivorsA validation of the ICF core set for stroke in Sweden. <i>Disability and Rehabilitation</i> , 2010 , 32, 551-9	2.4	47	
265	Does a short period of rehabilitation in the home setting facilitate functioning after stroke? A randomized controlled trial. <i>Clinical Rehabilitation</i> , 2006 , 20, 1038-49	3.3	46	
264	How well do standard stroke outcome measures reflect quality of life? A retrospective analysis of clinical trial data. <i>Stroke</i> , 2013 , 44, 3161-5	6.7	45	
263	Rising from a chair: a simple screening test for physical function in predialysis patients. <i>Scandinavian Journal of Urology and Nephrology</i> , 2008 , 42, 293-300		45	
262	Is Pain Intensity Really That Important to Assess in Chronic Pain Patients? A Study Based on the Swedish Quality Registry for Pain Rehabilitation (SQRP). <i>PLoS ONE</i> , 2013 , 8, e65483	3.7	44	
261	Neglect assessment as an application of virtual reality. <i>Acta Neurologica Scandinavica</i> , 2007 , 116, 157-63	33.8	40	
260	How is walking speed related to muscle strength? A study of healthy persons and persons with late effects of polio. <i>Archives of Physical Medicine and Rehabilitation</i> , 2004 , 85, 1923-8	2.8	40	
259	Upper motor neuron lesions: their effect on muscle performance and appearance in stroke patients with minor motor impairment. <i>Archives of Physical Medicine and Rehabilitation</i> , 1999 , 80, 155-61	2.8	40	
258	Timed Up & Go as a measure for longitudinal change in mobility after stroke - Postural Stroke Study in Gothenburg (POSTGOT). <i>Journal of NeuroEngineering and Rehabilitation</i> , 2014 , 11, 83	5.3	39	
257	Prestroke physical activity could influence acute stroke severity (part of PAPSIGOT). <i>Neurology</i> , 2018 , 91, e1461-e1467	6.5	39	
256	Baseline characteristics and the effects of two years of growth hormone replacement therapy in adults with growth hormone deficiency previously treated for Cushing's disease. <i>Clinical Endocrinology</i> , 2004 , 60, 550-9	3.4	38	
255	Evidence-based community stroke rehabilitation. <i>Stroke</i> , 2013 , 44, 293-7	6.7	37	
254	Relationship between functional disability and costs one and two years post stroke. <i>PLoS ONE</i> , 2017 , 12, e0174861	3.7	37	
253	SALGOTStroke Arm Longitudinal study at the University of Gothenburg, prospective cohort study protocol. <i>BMC Neurology</i> , 2011 , 11, 56	3.1	36	
252	Walking speed and distance in different environments of subjects in the later stage post-stroke. <i>Physiotherapy Theory and Practice</i> , 2010 , 26, 519-27	1.5	35	
251	Return to work after acquired brain injury: facilitators and hindrances observed in a sub-acute rehabilitation setting. <i>Work</i> , 2010 , 36, 431-9	1.6	34	

250	Muscle and tendon stiffness in patients with upper motor neuron lesion following a stroke. <i>European Journal of Applied Physiology</i> , 2000 , 82, 275-9	3.4	34
249	Evaluation of the Post Stroke Checklist: a pilot study in the United Kingdom and Singapore. <i>International Journal of Stroke</i> , 2014 , 9 Suppl A100, 76-84	6.3	33
248	Is walking endurance associated with activity and participation late after stroke?. <i>Disability and Rehabilitation</i> , 2011 , 33, 2053-7	2.4	33
247	Outcome after severe brain damage, what makes the difference?. <i>Brain Injury</i> , 2005 , 19, 493-503	2.1	33
246	The effects of five-year growth hormone replacement therapy on muscle strength in elderly hypopituitary patients. <i>Clinical Endocrinology</i> , 2005 , 62, 105-13	3.4	33
245	Regional left ventricular wall motion abnormalities in idiopathic dilated cardiomyopathy. <i>American Journal of Cardiology</i> , 1990 , 65, 364-70	3	33
244	More outcomes than trials: a call for consistent data collection across stroke rehabilitation trials. <i>International Journal of Stroke</i> , 2013 , 8, 18-24	6.3	32
243	Facilitators and barriers of stroke survivors in the early post-stroke phase. <i>Disability and Rehabilitation</i> , 2009 , 31, 1584-91	2.4	32
242	Reliability of clinical muscular endurance tests in patients with chronic heart failure. <i>European Journal of Cardiovascular Nursing</i> , 2006 , 5, 122-6	3.3	30
241	Cardiorespiratory effects of warm water immersion in elderly patients with chronic heart failure. <i>Clinical Physiology and Functional Imaging</i> , 2005 , 25, 313-7	2.4	30
240	Two years after cardiac arrest; cognitive status, ADL function and living situation. <i>Brain Injury</i> , 2008 , 22, 972-8	2.1	29
239	Upper Limb Kinematics in Stroke and Healthy Controls Using Target-to-Target Task in Virtual Reality. <i>Frontiers in Neurology</i> , 2018 , 9, 300	4.1	28
238	Aquatic exercise is effective in improving exercise performance in patients with heart failure and type 2 diabetes mellitus. <i>Evidence-based Complementary and Alternative Medicine</i> , 2012 , 2012, 349209	2.3	28
237	Muscular performance in heart failure. <i>Journal of Cardiac Failure</i> , 1998 , 4, 97-104	3.3	28
236	Peripheral muscle training in patients with clinical signs of heart failure. <i>Journal of Rehabilitation Medicine</i> , 1997 , 29, 121-7		28
235	Pituitary function and functional outcome in adults after severe traumatic brain injury: the long-term perspective. <i>Journal of Neurotrauma</i> , 2013 , 30, 271-80	5.4	27
234	Baseline characteristics and the effects of two years of growth hormone (GH) replacement therapy in adults with GH deficiency previously treated for acromegaly. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 2531-8	5.6	27
233	Computer game-based upper extremity training in the home environment in stroke persons: a single subject design. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2014 , 11, 35	5.3	26

(2017-2006)

232	Cross-diagnostic validity in a generic instrument: an example from the Functional Independence Measure in Scandinavia. <i>Health and Quality of Life Outcomes</i> , 2006 , 4, 55	3	26	
231	Stroke survivors' priorities for research related to life after stroke. <i>Topics in Stroke Rehabilitation</i> , 2021 , 28, 153-158	2.6	26	
230	Movement Kinematics of the Ipsilesional Upper Extremity in Persons With Moderate or Mild Stroke. <i>Neurorehabilitation and Neural Repair</i> , 2017 , 31, 376-386	4.7	25	
229	Factors affecting participation after traumatic brain injury. <i>Journal of Rehabilitation Medicine</i> , 2013 , 45, 765-70	3.4	25	
228	Cerebral palsy prevalence, subtypes, and associated impairments: a population-based comparison study of adults and children. <i>Developmental Medicine and Child Neurology</i> , 2019 , 61, 1162-1167	3.3	24	
227	Occupational Therapists[and Patients[Perceptions of ABILHAND, a New Assessment Tool for Measuring Manual Ability. <i>Scandinavian Journal of Occupational Therapy</i> , 2004 , 11, 107-117	2.1	24	
226	Gothenburg very early supported discharge study (GOTVED) NCT01622205: a block randomized trial with superiority design of very early supported discharge for patients with stroke. <i>BMC Neurology</i> , 2013 , 13, 66	3.1	23	
225	Relationship between walking function and 1-legged bicycling test in subjects in the later stage post-stroke. <i>Journal of Rehabilitation Medicine</i> , 2008 , 40, 721-6	3.4	23	
224	Recommendations for Clinical Trials in ICH: The Second Hemorrhagic Stroke Academia Industry Roundtable. <i>Stroke</i> , 2020 , 51, 1333-1338	6.7	22	
223	Perceptions of physical activity and walking in an early stage after stroke or acquired brain injury. <i>PLoS ONE</i> , 2017 , 12, e0173463	3.7	22	
222	Sex as predictor for achieved health outcomes and received care in ischemic stroke and intracerebral hemorrhage: a register-based study. <i>Biology of Sex Differences</i> , 2018 , 9, 11	9.3	22	
221	Participation and autonomy five years after stroke: A longitudinal observational study. <i>PLoS ONE</i> , 2019 , 14, e0219513	3.7	22	
220	Application and validation of the barrow neurological institute screen for higher cerebral functions in a control population and in patient groups commonly seen in neurorehabilitation. <i>Acta Dermato-Venereologica</i> , 2007 , 39, 547-53	2.2	22	
219	Disability in a 4-year follow-up study of people with post-polio syndrome. <i>Acta Dermato-Venereologica</i> , 2007 , 39, 175-80	2.2	22	
218	Muscular effects in late polio. Acta Physiologica Scandinavica, 2001, 171, 335-40		22	
217	Physical activity, muscle performance and quality of life in patients treated with chronic peritoneal dialysis. <i>Scandinavian Journal of Urology and Nephrology</i> , 2001 , 35, 71-8		22	
216	Cardiac function of the leopard shark, Triakis semifasciata. <i>Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology</i> , 1990 , 160, 259-268	2.2	22	
215	Health status and psychological distress among in-hospital cardiac arrest survivors in relation to gender. <i>Resuscitation</i> , 2017 , 114, 27-33	4	21	

214	Muscle strength in elderly adults with GH deficiency after 10 years of GH replacement. <i>European Journal of Endocrinology</i> , 2010 , 163, 207-15	6.5	21
213	Description of functional disability among younger stroke patients: exploration of activity and participation and environmental factors. <i>International Journal of Rehabilitation Research</i> , 2009 , 32, 124-	.3 ¹ 7.8	21
212	Vitality among Swedish patients with post-polio: a physiological phenomenon. <i>Journal of Rehabilitation Medicine</i> , 2008 , 40, 709-14	3.4	21
211	Intra- and inter-rater reliability of Fugl-Meyer Assessment of Upper Extremity in stroke. <i>Journal of Rehabilitation Medicine</i> , 2019 , 51, 652-659	3.4	20
210	A cohort study investigating a simple, early assessment to predict upper extremity function after stroke - a part of the SALGOT study. <i>BMC Neurology</i> , 2015 , 15, 92	3.1	20
209	Onset, time course and prediction of spasticity after stroke or traumatic brain injury. <i>Annals of Physical and Rehabilitation Medicine</i> , 2019 , 62, 431-434	3.8	20
208	Physical activity in community-dwelling stroke survivors and a healthy population is not explained by motor function only. <i>PM and R</i> , 2014 , 6, 139-45	2.2	20
207	Improving the Development, Monitoring and Reporting of Stroke Rehabilitation Research: Consensus-Based Core Recommendations from the Stroke Recovery and Rehabilitation Roundtable. <i>Neurorehabilitation and Neural Repair</i> , 2017 , 31, 877-884	4.7	20
206	Can rehabilitation in the home setting reduce the burden of care for the next-of-kin of stroke victims?. <i>Acta Dermato-Venereologica</i> , 2007 , 39, 27-32	2.2	20
205	Circuit training in community-living "younger" men after stroke. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2007 , 16, 122-9	2.8	20
204	Postural control in persons with late effects of polio. <i>Acta Neurologica Scandinavica</i> , 2006 , 113, 55-61	3.8	20
203	Development of fatigue during repeated eccentric-concentric muscle contractions of plantar flexors in patients with stroke. <i>Archives of Physical Medicine and Rehabilitation</i> , 1999 , 80, 1247-52	2.8	20
202	Predictors of Spasticity After Stroke. Current Physical Medicine and Rehabilitation Reports, 2016, 4, 182-	18 <i>5</i> 7	19
201	Fatigue, psychosocial adaptation and quality of life one year after traumatic brain injury and suspected traumatic axonal injury; evaluations of patients and relatives: a pilot study. <i>Journal of Rehabilitation Medicine</i> , 2013 , 45, 771-7	3.4	19
200	Beyond conventional stroke guidelines: setting priorities. <i>Stroke</i> , 2007 , 38, 2185-90	6.7	19
199	Functional and cognitive capacity and health-related quality of life 2 years after day hospital rehabilitation for stroke: a prospective study. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2007 , 16, 208-15	2.8	19
198	A validation study using a modified version of Postural Assessment Scale for Stroke Patients: Postural Stroke Study in Gothenburg (POSTGOT). <i>Journal of NeuroEngineering and Rehabilitation</i> , 2011 , 8, 57	5.3	18
197	Stop using the Ashworth scale for the assessment of spasticity. <i>Journal of Neurology, Neurosurgery</i> and Psychiatry, 2010 , 81, 2	5.5	18

196	Physical activity, ambulation, and motor impairment late after stroke. <i>Stroke Research and Treatment</i> , 2012 , 2012, 818513	1.7	18	
195	Improved survival after non-traumatic subarachnoid haemorrhage with structured care pathways and modern intensive care. <i>Clinical Neurology and Neurosurgery</i> , 2015 , 138, 52-8	2	17	
194	Patients' expectations of coming home with Very Early Supported Discharge and home rehabilitation after stroke - an interview study. <i>BMC Neurology</i> , 2015 , 15, 235	3.1	17	•
193	A kinematic analysis of a haptic handheld stylus in a virtual environment: a study in healthy subjects. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2007 , 4, 13	5.3	17	
192	Virtual rehabilitation after stroke. Studies in Health Technology and Informatics, 2008, 136, 77-82	0.5	17	
191	Muscle strength in patients with acromegaly at diagnosis and during long-term follow-up. <i>European Journal of Endocrinology</i> , 2017 , 177, 217-226	6.5	16	
190	Walking performance and muscle strength in the later stage poststroke: a nonlinear relationship. <i>Archives of Physical Medicine and Rehabilitation</i> , 2013 , 94, 845-50	2.8	16	
189	Life after cardiac arrest: A very long term follow up. <i>Resuscitation</i> , 2015 , 91, 99-103	4	16	
188	Haptic virtual rehabilitation in stroke: transferring research into clinical practice. <i>Physical Therapy Reviews</i> , 2009 , 14, 322-335	0.7	16	
187	GH Is Needed for the Maturation of Muscle Mass and Strength in Adolescents. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 4765-4770	5.6	16	
186	Consequences and coping strategies six years after a subarachnoid hemorrhage - A qualitative study. <i>PLoS ONE</i> , 2017 , 12, e0181006	3.7	16	
185	Physical performance in individuals with late effects of polio. <i>Journal of Rehabilitation Medicine</i> , 1999 , 31, 244-9		16	
184	Kinematic Analysis Using 3D Motion Capture of Drinking Task in People With and Without Upper-extremity Impairments. <i>Journal of Visualized Experiments</i> , 2018 ,	1.6	15	
183	Recovery of kinematic arm function in well-performing people with subacute stroke: a longitudinal cohort study. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2018 , 15, 67	5.3	15	
182	Rasch analysis of the modified version of the postural assessment scale for stroke patients: postural stroke study in Gothenburg (POSTGOT). <i>BMC Neurology</i> , 2014 , 14, 134	3.1	15	
181	THE STANDING HEEL-RISE TEST IN PATIENTS WITH UPPER MOTOR NEURON LESION DUE TO STROKE. <i>Journal of Rehabilitation Medicine</i> , 1998 , 30, 73-80		15	
180	Rehabilitation after stroke using virtual reality, haptics (force feedback) and telemedicine. <i>Studies in Health Technology and Informatics</i> , 2006 , 124, 51-6	0.5	15	
179	2022 Guideline for the Management of Patients With Spontaneous Intracerebral Hemorrhage: A Guideline From the American Heart Association/American Stroke Association <i>Stroke</i> , 2022 , 101161ST	 -R60 <u>7</u> 00	0000000	

178	Upper extremity recovery after ischaemic and haemorrhagic stroke: Part of the SALGOT study. <i>European Stroke Journal</i> , 2016 , 1, 310-319	5.6	14
177	Unlocking the locked in; a need for team approach in rehabilitation of survivors with locked-in syndrome. <i>Acta Neurologica Scandinavica</i> , 2012 , 125, 192-8	3.8	14
176	Applying the Comprehensive International Classification of Functioning, Disability and Health Core Sets for stroke framework to stroke survivors living in the community. <i>Journal of Rehabilitation Medicine</i> , 2013 , 45, 331-40	3.4	14
175	The Swedish version of the Haemophilia Activity List. <i>Haemophilia</i> , 2011 , 17, 662-8	3.3	14
174	Electrophysiologic evaluation of muscle fatigue development and recovery in late polio. <i>Archives of Physical Medicine and Rehabilitation</i> , 2000 , 81, 770-776	2.8	14
173	Persons with Haemophilia in Sweden- Experiences and Strategies in Everyday Life. A Single Centre Study. <i>PLoS ONE</i> , 2015 , 10, e0139690	3.7	14
172	A home-based, self-administered stimulation program to improve selected hand functions of chronic stroke. <i>NeuroRehabilitation</i> , 2003 , 18, 215-25	2	14
171	Predicting Longitudinal Progression in Functional Mobility After Stroke: A Prospective Cohort Study. <i>Stroke</i> , 2020 , 51, 2179-2187	6.7	13
170	Early prediction of physical activity level 1 year after stroke: a longitudinal cohort study. <i>BMJ Open</i> , 2017 , 7, e016369	3	13
169	Return to work after specialized rehabilitation-An explorative longitudinal study in a cohort of severely disabled persons with stroke in seven countries: The Sunnaas International Network stroke study. <i>Brain and Behavior</i> , 2018 , 8, e01055	3.4	13
168	Post polio syndrome: fatigued patients a specific subgroup?. <i>Journal of Rehabilitation Medicine</i> , 2011 , 43, 39-45	3.4	13
167	Agreement in activities of daily living performance after stroke in a postal questionnaire and interview of community-living persons. <i>Acta Neurologica Scandinavica</i> , 2009 , 119, 390-6	3.8	13
166	Electrophysiologic evaluation of muscle fatigue development and recovery in late polio. <i>Archives of Physical Medicine and Rehabilitation</i> , 2000 , 81, 770-776	2.8	13
165	Well-being among survivors of out-of-hospital cardiac arrest: a cross-sectional retrospective study in Sweden. <i>BMJ Open</i> , 2018 , 8, e021729	3	13
164	Muscle performance in an urban population sample of 40- to 79-year-old men and women. <i>Journal of Rehabilitation Medicine</i> , 2000 , 32, 159-67		13
163	Long-term participation 7-8 years after stroke: Experiences of people in working-age. <i>PLoS ONE</i> , 2019 , 14, e0213447	3.7	12
162	Multidimensional fatigue inventory and post-polio syndrome - a Rasch analysis. <i>Health and Quality of Life Outcomes</i> , 2015 , 13, 20	3	12
161	Life situation 5 years after subarachnoid haemorrhage. <i>Acta Neurologica Scandinavica</i> , 2018 , 137, 99-10	4 3.8	12

160	End-point kinematics using virtual reality explaining upper limb impairment and activity capacity in stroke. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2019 , 16, 82	5.3	12
159	Very Early MoCA Can Predict Functional Dependence at 3 Months After Stroke: A Longitudinal, Cohort Study. <i>Frontiers in Neurology</i> , 2019 , 10, 1051	4.1	12
158	Gender differences in care-seeking behavior and healthcare consumption immediately after whiplash trauma. <i>PLoS ONE</i> , 2017 , 12, e0176328	3.7	12
157	Commercial head-mounted display virtual reality for upper extremity rehabilitation in chronic stroke: a single-case design study. <i>Journal of NeuroEngineering and Rehabilitation</i> , 2020 , 17, 154	5.3	12
156	Self-perceived impact of stroke: A longitudinal comparison between one and five years post-stroke. Journal of Rehabilitation Medicine, 2019 , 51, 660-664	3.4	11
155	NIHSS is not enough for cognitive screening in acute stroke: A cross-sectional, retrospective study. <i>Scientific Reports</i> , 2020 , 10, 534	4.9	11
154	Life satisfaction in persons with severe stroke - A longitudinal report from the Sunnaas International Network (SIN) stroke study. <i>European Stroke Journal</i> , 2017 , 2, 154-162	5.6	11
153	The impact of physical function on participation in the first year post-stroke. <i>Acta Neurologica Scandinavica</i> , 2017 , 135, 649-655	3.8	11
152	Cognitive impact of traumatic axonal injury (TAI) and return to work. <i>Brain Injury</i> , 2013 , 27, 521-8	2.1	11
151	Process skill rather than motor skill seems to be a predictor of costs for rehabilitation after a stroke in working age; a longitudinal study with a 1 year follow up post discharge. <i>BMC Health Services Research</i> , 2007 , 7, 209	2.9	11
150	Experiences of returning to work and maintaining work 7 to 8 years after a stroke: a qualitative interview study in Sweden. <i>BMJ Open</i> , 2018 , 8, e021182	3	11
149	Return to work following diagnosis of low-grade glioma: A nationwide matched cohort study. <i>Neurology</i> , 2020 , 95, e856-e866	6.5	10
148	Early clinical predictors of motor function in the upper extremity one month post-stroke. <i>Journal of Rehabilitation Medicine</i> , 2017 , 49, 216-222	3.4	10
147	Factors enhancing activities of daily living after stroke in specialized rehabilitation: an observational multicenter study within the Sunnaas International Network. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2017 , 53, 725-734	4.4	10
146	Work participation among middle-aged persons with cerebral palsy or spina bifidaa longitudinal study. <i>Disability and Health Journal</i> , 2014 , 7, 251-5	4.2	10
145	A cross sectional study of upper extremity strength ten days after a stroke; relationship between patient-reported and objective measures. <i>BMC Neurology</i> , 2015 , 15, 178	3.1	10
144	Awareness and the arguments for and against the International Classification of Functioning, Disability and Health among representatives of disability organisations. <i>International Journal of Environmental Research and Public Health</i> , 2015 , 12, 3293-300	4.6	10
143	Validation of the Comprehensive ICF Core Set for Stroke by exploring the patient's perspective on functioning in everyday life: a qualitative study. <i>International Journal of Rehabilitation Research</i> , 2014 , 37, 302-10	1.8	10

142	Stretch-shortening cycle in patients with upper motor neuron lesions due to stroke. <i>European Journal of Applied Physiology</i> , 1997 , 75, 312-8	3.4	10
141	The Quebec classification and a new Swedish classification for whiplash-associated disorders in relation to life satisfaction in patients at high risk of chronic functional impairment and disability. <i>Journal of Rehabilitation Medicine</i> , 2002 , 34, 114-8	3.4	10
140	Reliability and validity of the Northwick Park Dependency Score (NPDS) Swedish version 6.0. <i>Clinical Rehabilitation</i> , 2005 , 19, 419-25	3.3	10
139	One-year longitudinal study of psychological distress and self-assessed health in survivors of out-of-hospital cardiac arrest. <i>BMJ Open</i> , 2019 , 9, e029756	3	10
138	Experiences using the poststroke checklist in Sweden with a focus on feasibility and relevance: a mixed-method design. <i>BMJ Open</i> , 2019 , 9, e028218	3	10
137	Translation and cultural validation of clinical observational scales - the Fugl-Meyer assessment for post stroke sensorimotor function in Colombian Spanish. <i>Disability and Rehabilitation</i> , 2019 , 41, 2317-23	3 23 1	10
136	Analyzing the modified rankin scale using concepts of the international classification of functioning, disability and health. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2016 , 52, 203-13	4.4	10
135	Hospital comparison of stroke care in Sweden: a register-based study. <i>BMJ Open</i> , 2017 , 7, e015244	3	9
134	Work-related predictors for return to work after stroke. <i>Acta Neurologica Scandinavica</i> , 2019 , 139, 382-3	388	9
133	Improving life after stroke needs global efforts to implement evidence-based physical activity pathways. <i>International Journal of Stroke</i> , 2019 , 14, 457-459	6.3	9
132	Self-Perceived Participation and Autonomy at 1-Year Post Stroke: A Part of the Stroke Arm Longitudinal Study at the University of Gothenburg (SALGOT Study). <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2018 , 27, 1115-1122	2.8	9
131	Responsiveness of a modified version of the postural assessment scale for stroke patients and longitudinal change in postural control after stroke- Postural Stroke Study in Gothenburg (POSTGOT). <i>Journal of NeuroEngineering and Rehabilitation</i> , 2013 , 10, 8	5.3	9
130	Self-Assessed Physical, Cognitive, and Emotional Impact of Stroke at 1 Month: The Importance of Stroke Severity and Participation. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2017 , 26, 57-63	2.8	9
129	Stroke treated at a neurosurgical ward: a cohort study. <i>Acta Neurologica Scandinavica</i> , 2015 , 132, 329-36	53.8	9
128	Specialized stroke rehabilitation services in seven countries: Preliminary results from nine rehabilitation centers. <i>International Journal of Stroke</i> , 2015 , 10, 1236-46	6.3	9
127	Quality of life in Swedish patients with post-polio syndrome with a focus on age and sex. <i>International Journal of Rehabilitation Research</i> , 2014 , 37, 173-9	1.8	9
126	Sick leave after traumatic brain injury. The person or the diagnosiswhich has greater impact?. <i>Scandinavian Journal of Public Health</i> , 2010 , 38, 541-7	3	9
125	The effects of endurance training in persons with a hereditary myosin myopathy. <i>Acta Neurologica Scandinavica</i> , 2004 , 110, 80-6	3.8	9

124	Patterns and predictors of sick leave after Covid-19 and long Covid in a national Swedish cohort. <i>BMC Public Health</i> , 2021 , 21, 1023	4.1	9	
123	Kinematic upper extremity performance in people with near or fully recovered sensorimotor function after stroke. <i>Physiotherapy Theory and Practice</i> , 2019 , 35, 822-832	1.5	9	
122	Feasibility of Cognitive Functions Screened With the Montreal Cognitive Assessment in Determining ADL Dependence Early After Stroke. <i>Frontiers in Neurology</i> , 2018 , 9, 705	4.1	9	
121	Impaired Motor Function in the Affected Arm Predicts Impaired Postural Balance After Stroke: A Cross Sectional Study. <i>Frontiers in Neurology</i> , 2019 , 10, 912	4.1	8	
120	Improving economic evaluations in stroke: A report from the ESO Health Economics Working Group. <i>European Stroke Journal</i> , 2020 , 5, 184-192	5.6	8	
119	Symptoms and consequences of subarachnoid haemorrhage after 7 years. <i>Acta Neurologica Scandinavica</i> , 2019 , 140, 429-434	3.8	8	
118	Processes influencing participation in the daily lives of immigrants living with polio in Sweden: A secondary analysis. <i>Journal of Occupational Science</i> , 2017 , 24, 203-215	1	8	
117	Self-reported activity and functioning in daily life; the perspective of persons with haemophilia living in Sweden. <i>European Journal of Haematology</i> , 2015 , 95, 336-41	3.8	8	
116	Exploration of some personal factors with the International Classification of Functioning, Disability and Health core sets for stroke. <i>Journal of Rehabilitation Medicine</i> , 2013 , 45, 609-15	3.4	8	
115	Changes in ability, perceived difficulty and use of assistive devices in everyday life: a 4-year follow-up study in people with late effects of polio. <i>Acta Neurologica Scandinavica</i> , 2009 , 120, 324-30	3.8	8	
114	Effects of day hospital rehabilitation after stroke. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2006 , 15, 106-13	2.8	8	
113	Induced shift in myosin heavy chain expression in myosin myopathy by endurance training. <i>Journal of Neurology</i> , 2004 , 251, 179-83	5.5	8	
112	Normal concentrations of serum insulin-like growth factor-1 in late polio. <i>Archives of Physical Medicine and Rehabilitation</i> , 1995 , 76, 732-5	2.8	8	
111	Intra- and interrater reliability of Fugl-Meyer Assessment of Lower Extremity early after stroke. <i>Brazilian Journal of Physical Therapy</i> , 2020 , 25, 709-709	3.7	8	
110	Return to work after stroke: A Swedish nationwide registry-based study. <i>Acta Neurologica Scandinavica</i> , 2020 , 141, 56-64	3.8	8	
109	Experiences, needs, and preferences for follow-up after stroke perceived by people with stroke and healthcare professionals: A focus group study. <i>PLoS ONE</i> , 2019 , 14, e0223338	3.7	7	
108	Upper limb kinematics during the first year after stroke: the stroke arm longitudinal study at the University of Gothenburg (SALGOT). <i>Journal of NeuroEngineering and Rehabilitation</i> , 2020 , 17, 76	5.3	7	
107	Histopathological changes in skeletal muscle associated with chronic ischaemia. <i>Apmis</i> , 2016 , 124, 935-9	94.14	7	

106	A comparison of stroke rehabilitation; data from two national cohorts. <i>Acta Neurologica Scandinavica</i> , 2016 , 134, 284-91	3.8	7
105	Gothenburg Very Early Supported Discharge study (GOTVED): a randomised controlled trial investigating anxiety and overall disability in the first year after stroke. <i>BMC Neurology</i> , 2019 , 19, 277	3.1	7
104	Enhancing patient-provider communication for long-term post-stroke spasticity management. <i>Acta Neurologica Scandinavica</i> , 2013 , 128, 305-10	3.8	7
103	Evidence-based position paper on Physical and Rehabilitation Medicine professional practice for persons with stroke. The European PRM position (UEMS PRM Section). <i>European Journal of Physical and Rehabilitation Medicine</i> , 2018 , 54, 957-970	4.4	7
102	The Association of Pre-stroke Psychosis and Post-stroke Levels of Health, Resource Utilization, and Care Process: A Register-Based Study. <i>Frontiers in Neurology</i> , 2018 , 9, 1042	4.1	7
101	Experienced pain after stroke: a cross-sectional 5-year follow-up study. <i>BMC Neurology</i> , 2020 , 20, 4	3.1	6
100	Occupations and means of living in adults with cerebral palsy or meningomyelocele during two decades in Sweden. <i>Scandinavian Journal of Disability Research</i> , 2011 , 13, 21-35	1.3	6
99	The structural properties of the European Brain Injury Questionnaire. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2004 , 13, 122-8	2.8	6
98	Detection and quantitation of ischemic left ventricular dysfunction using a new video intensity technique for regional wall motion evaluation. <i>American Heart Journal</i> , 1990 , 120, 1058-72	4.9	6
97	Characteristics of Patients at First Visit to a Polio Clinic in Sweden. <i>PLoS ONE</i> , 2016 , 11, e0150286	3.7	6
96	Cognitive medicine - a new approach in health care science. <i>BMC Psychiatry</i> , 2018 , 18, 42	4.2	5
95	Increasing symptoms, met and unmet needs in adults with cerebral palsy or meningomyelocele a longitudinal follow-up. <i>Scandinavian Journal of Disability Research</i> , 2013 , 15, 249-263	1.3	5
94	Cognitive screening and behavioural observation of functional ability in patients with multiple episode schizophrenia: an exploratory study. <i>BMJ Open</i> , 2017 , 7, e014783	3	5
93	How can we tell who is aware? Where does the veracity lie?. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2012 , 21, 812-8	2.8	5
92	Experiences of participation in a Swedish society among adults with cerebral palsy or spina bifida: involvement and challenges. <i>Journal of Social Work in Disability and Rehabilitation</i> , 2013 , 12, 256-71		5
91	VISTA-Rehab: a resource for stroke rehabilitation trials. <i>International Journal of Stroke</i> , 2010 , 5, 447-52	6.3	5
90	Evaluation of Usefulness of the Elinor Console for Home-Based Stroke Rehabilitation 2011,		5
89	Rehabilitation problems after acute disseminated encephalomyelitis: four cases. <i>Journal of Rehabilitation Medicine</i> , 2003 , 35, 20-5	3.4	5

(2011-2003)

88	31P MRS evaluation of fatigue in anterior tibial muscle in postpoliomyelitis patients and healthy volunteers. <i>Clinical Physiology and Functional Imaging</i> , 2003 , 23, 190-8	2.4	5
87	Sex Differences in Stroke Care and Outcome 2005-2018: Observations From the Swedish Stroke Register. <i>Stroke</i> , 2021 , 52, 3233-3242	6.7	5
86	"My life after stroke through a camera lens"- A photovoice study on participation in Sweden. <i>PLoS ONE</i> , 2019 , 14, e0222099	3.7	4
85	Update on the EFFECTS study of fluoxetine for stroke recovery: a randomised controlled trial in Sweden. <i>Trials</i> , 2020 , 21, 233	2.8	4
84	WebRehab: a Swedish database for quality control in rehabilitation. <i>Journal of Rehabilitation Medicine</i> , 2014 , 46, 958-62	3.4	4
83	The ICF: International Classification of Functioning, Disability and Health (ICF)a Swiss army knife? Accessibility and disability in a Scandinavian disability magazine (SDM)a quantitative content analysis. <i>Health Policy</i> , 2012 , 108, 67-75	3.2	4
82	Telehealth with 3D games for stroke rehabilitation. <i>International Journal on Disability and Human Development</i> , 2011 , 10,		4
81	One-legged bicycling as an assessment tool for patients with stroke. <i>Acta Neurologica Scandinavica</i> , 2005 , 111, 373-8	3.8	4
80	Measurement and Evaluation of Outcomes in Rehabilitation. <i>Journal of Rehabilitation Medicine</i> , 2005 , 37, 1-2	3.4	4
79	Games on Prescription! Evaluation of the Elinor Console for Home-Based Stroke Rehabilitation. <i>Lecture Notes in Computer Science</i> , 2013 , 49-64	0.9	4
78	Patients' physical activity in stroke units in Latvia and Sweden. <i>Brain and Behavior</i> , 2021 , 11, e02110	3.4	4
77	Levels of physical activity before and after stroke in relation to early cognitive function. <i>Scientific Reports</i> , 2021 , 11, 9078	4.9	4
76	Which clinical and sociodemographic determinants are associated with self-perceived manual ability at one year after stroke?. <i>Disability and Rehabilitation</i> , 2020 , 42, 2279-2286	2.4	4
75	Occupational gaps 5 years after stroke. <i>Brain and Behavior</i> , 2019 , 9, e01234	3.4	3
74	Easily Conducted Tests During the First Week Post-stroke Can Aid the Prediction of Arm Functioning at 6 Months. <i>Frontiers in Neurology</i> , 2019 , 10, 1371	4.1	3
73	Long-term outcome after reperfusion-treated stroke. <i>Journal of Rehabilitation Medicine</i> , 2017 , 49, 316-	33.14	3
72	The extent of using mobility assistive devices can partly explain fatigue among persons with late effects of polio - a retrospective registry study in Sweden. <i>BMC Neurology</i> , 2016 , 16, 230	3.1	3
71	Functioning of the upper extremity in persons with late polio. <i>European Journal of Neurology</i> , 2011 , 18, 354-358	6	3

70	SiPP (Stroke in Pregnancy and Postpartum): A prospective, observational, international, multicentre study on pathophysiological mechanisms, clinical profile, management and outcome of cerebrovascular diseases in pregnant and postpartum women. <i>European Stroke Journal</i> , 2020 , 5, 193-20	5.6 03	3
69	Changes in walking ability, intellectual disability, and epilepsy in adults with cerebral palsy over 50 years: a population-based follow-up study. <i>Developmental Medicine and Child Neurology</i> , 2021 , 63, 839-	843 ³	3
68	Impact of Intensive Gait Training With and Without Electromechanical Assistance in the Chronic Phase After Stroke-A Multi-Arm Randomized Controlled Trial With a 6 and 12 Months Follow Up. <i>Frontiers in Neuroscience</i> , 2021 , 15, 660726	5.1	3
67	Living in Latvia after stroke: the association between functional, social and personal factors and the level of self-perceived disability-a cross-sectional study. <i>BMJ Open</i> , 2016 , 6, e010327	3	3
66	Examination of the relevance of the ICF cores set for stroke by comparing with the Stroke Impact Scale. <i>Disability and Rehabilitation</i> , 2019 , 41, 508-513	2.4	3
65	Self-perceived mobility in immigrants in Sweden living with the late effects of polio. <i>Disability and Rehabilitation</i> , 2020 , 42, 3203-3208	2.4	3
64	An 8-year longitudinal study of muscle strength, muscle fiber size, and dynamic electromyogram in individuals with late polio 1998 , 21, 1428		3
63	Pre-stroke physical activity in relation to post-stroke outcomes - linked to the International Classification of Functioning, Disability and Health (ICF): A scoping review <i>Journal of Rehabilitation Medicine</i> , 2021 ,	3.4	3
62	Locked-in syndrome in Sweden, an explorative study of persons who underwent rehabilitation: a cohort study. <i>BMJ Open</i> , 2019 , 9, e023185	3	2
61	Relationship between self-reported and objectively measured manual ability varies during the first year post-stroke. <i>Scientific Reports</i> , 2020 , 10, 5093	4.9	2
60	Working capacity after a subarachnoid haemorrhage: A six-year follow-up. <i>Journal of Rehabilitation Medicine</i> , 2017 , 49, 738-743	3.4	2
59	Decline in cognitive function due to diffuse axonal injury does not necessarily imply a corresponding decline in ability to perform activities. <i>Disability and Rehabilitation</i> , 2016 , 38, 1006-15	2.4	2
58	Very early cognitive screening and return to work after stroke. <i>Topics in Stroke Rehabilitation</i> , 2019 , 26, 602-607	2.6	2
57	Muscle performance and fatigue in compensated chronic liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2019 , 54, 925-933	2.4	2
56	"The car is my extra legs" - Experiences of outdoor mobility amongst immigrants in Sweden with late effects of polio. <i>PLoS ONE</i> , 2019 , 14, e0224685	3.7	2
55	Letter by Murphy et al regarding article, "Fugl-Meyer assessment of sensorimotor function after stroke: standardized training procedure for clinical practice and clinical trials". <i>Stroke</i> , 2011 , 42, e402	6.7	2
54	A Comparison between Two Instruments for Assessing Dependency in Daily Activities: Agreement of the Northwick Park Dependency Score with the Functional Independence Measure. <i>Rehabilitation Research and Practice</i> , 2012 , 2012, 769513	1.2	2
53	Evidence-based medicine in physical and rehabilitation medicine: is this evidence-based rehabilitation?. <i>Journal of Rehabilitation Medicine</i> , 2008 , 40, 689-90	3.4	2

(2020-2006)

52	Passive stiffness of the plantar flexion in relation to age and functional performance. <i>Isokinetics and Exercise Science</i> , 2006 , 14, 25-31	0.6	2
51	Sitting up and transferring to a chair: two functional tests for patients with stroke. <i>Journal of Rehabilitation Medicine</i> , 2003 , 35, 180-3	3.4	2
50	Ischemic heart disease and regional left ventricular wall motion: a study comparing radial, centerline and a video intensity based slope technique. <i>International Journal of Cardiovascular Imaging</i> , 1990 , 6, 85-96		2
49	Very Early Cognitive Screening and Self-Reported Feeling of Fatigue Three Months After Stroke. <i>Frontiers in Human Neuroscience</i> , 2021 , 15, 742105	3.3	2
48	Barriers to cognitive screening in acute stroke units. Scientific Reports, 2021, 11, 19621	4.9	2
47	Representativity and co-morbidity: Two factors of importance when reporting health status among survivors of cardiac arrest. <i>Resuscitation</i> , 2016 , 101, 44-9	4	2
46	Return to work predicts perceived participation and autonomy by individuals with stroke. <i>Disability and Rehabilitation</i> , 2020 , 42, 3673-3678	2.4	2
45	Association of prestroke medicine use and health outcomes after ischaemic stroke in Sweden: a registry-based cohort study. <i>BMJ Open</i> , 2020 , 10, e036159	3	2
44	Recovery of arm function during acute to chronic stage of stroke quantified by kinematics. <i>Journal of Rehabilitation Medicine</i> , 2021 , 53, jrm00171	3.4	2
43	Self-reported activity of Swedish persons with haemophilia: Change over 2.5 years. <i>Journal of Rehabilitation Medicine</i> , 2018 , 50, 643-651	3.4	2
42	Effects of Fluoxetine on Outcomes at 12 Months After Acute Stroke: Results From EFFECTS, a Randomized Controlled Trial. <i>Stroke</i> , 2021 , 52, 3082-3087	6.7	2
41	Level of physical activity is positively correlated with perceived impact on life 12 months after stroke: A cross-sectional study. <i>Journal of Rehabilitation Medicine</i> , 2020 , 52, jrm00056	3.4	1
40	Reflecting the World Report on Disability: a report from Sweden. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2014 , 93, S42-6	2.6	1
39	Re: 'Potential effectiveness of three different treatment approaches to improve minimal to moderate arm and hand function after stroke-a pilot randomized controlled clinical trial'. <i>Clinical Rehabilitation</i> , 2012 , 26, 758-9; author reply 759-60	3.3	1
38	Commentary on gait training in patients with spinal cord injury. <i>Journal of Rehabilitation Medicine</i> , 2010 , 42, 527	3.4	1
37	A very long-term longitudinal follow-up of persons with late effects of polio. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2020 , 56, 155-159	4.4	1
36	Recurrent sick leave after COVID-19: investigating the first wave of the pandemic in a comprehensive Swedish registry-based study. <i>BMC Public Health</i> , 2021 , 21, 1914	4.1	1
35	Self-perceived postural balance correlates with postural balance and anxiety during the first year after stroke: a part of the randomized controlled GOTVED study. <i>BMC Neurology</i> , 2020 , 20, 410	3.1	1

34	Differences in self-perceived general health, pain, and depression 1 to 5 years post-stroke related to work status at 1 year. <i>Scientific Reports</i> , 2020 , 10, 13251	4.9	1
33	Health insurance utilisation after ischaemic stroke in Sweden: a retrospective cohort study in a system of universal healthcare and social insurance. <i>BMJ Open</i> , 2021 , 11, e043826	3	1
32	Influence of mobility restrictions on post-stroke pain. Brain and Behavior, 2021, 11, e02092	3.4	1
31	Physical inactivity after stroke: Incidence and early predictors based on 190 individuals in a 1-year follow-up of the Fall Study of Gothenburg. <i>Journal of Rehabilitation Medicine</i> , 2021 , 53, jrm00224	3.4	1
30	A prospective cohort study on longitudinal trajectories of cognitive function after stroke. <i>Scientific Reports</i> , 2021 , 11, 17271	4.9	1
29	Twelve-Month Outcomes of the AFFINITY Trial of Fluoxetine for Functional Recovery After Acute Stroke: AFFINITY Trial Steering Committee on Behalf of the AFFINITY Trial Collaboration. <i>Stroke</i> , 2021 , 52, 2502-2509	6.7	1
28	A register-based study on associations between pre-stroke physical activity and cognition early after stroke (part of PAPSIGOT) <i>Scientific Reports</i> , 2022 , 12, 5779	4.9	1
27	Urdu translation and cross-cultural validation of the Fugl-Meyer assessment in people with stroke. <i>Disability and Rehabilitation</i> , 2021 , 1-6	2.4	1
26	Diastolic regional wall motion asynchrony in patients with hypertension. <i>International Journal of Cardiovascular Imaging</i> , 1992 , 8, 231-42		0
25	Dependency in Activities of Daily Living During the First Year After Stroke. <i>Frontiers in Neurology</i> , 2021 , 12, 736684	4.1	O
24	Prestroke physical activity and outcomes after intracerebral haemorrhage in comparison to ischaemic stroke: protocol for a matched cohort study (part of PAPSIGOT). <i>BMJ Open</i> , 2021 , 11, e05306	57 ³	0
23	A register-based study comparing planned rehabilitation following acute stroke in 2011 and 2017. <i>Scientific Reports</i> , 2021 , 11, 23001	4.9	O
22	Fatigue after acquired brain injury impacts health-related quality of life: an exploratory cohort study. <i>Scientific Reports</i> , 2021 , 11, 22153	4.9	0
21	Quantifying an Upper Extremity Everyday Task With 3D Kinematic Analysis in People With Spinal Cord Injury and Non-disabled Controls. <i>Frontiers in Neurology</i> , 2021 , 12, 755790	4.1	O
20	Health Conditions in Adults With Cerebral Palsy: The Association With CP Subtype and Severity of Impairments. <i>Frontiers in Neurology</i> , 2021 , 12, 732939	4.1	0
19	Levels of physical activity in acute stroke patients treated at a stroke unit: A prospective, observational study. <i>Journal of Rehabilitation Medicine</i> , 2020 , 52, jrm00041	3.4	O
18	Respiratory complications during initial rehabilitation and survival following spinal cord injury in Sweden: a retrospective study. <i>Spinal Cord</i> , 2021 , 59, 659-664	2.7	0
17	Predictors for self-reported feeling of depression three months after stroke: A longitudinal cohort study. <i>Journal of Rehabilitation Medicine</i> , 2021 , 53, jrm00170	3.4	O

LIST OF PUBLICATIONS

16	Associations between upper extremity functioning and kinematics in people with spinal cord injury. Journal of NeuroEngineering and Rehabilitation, 2021, 18, 147	5.3	О
15	Stroke-related health problems and associated actions identified with the post-stroke checklist among nursing home residents <i>BMC Cardiovascular Disorders</i> , 2022 , 22, 50	2.3	O
14	Development of a Digital Tool for People With a Long-Term Condition Using Stroke as a Case Example: Participatory Design Approach. <i>JMIR Human Factors</i> , 2022 , 9, e35478	2.5	O
13	A risk stratification tool for prehospital triage of patients exposed to a whiplash trauma. <i>PLoS ONE</i> , 2019 , 14, e0216694	3.7	
12	Neurologische Rehabilitation [Wohin geht die Reise?. Neuroreha, 2019 , 11, 20-24	0.2	
11	Longitudinal changes of self-perceived manual ability the first year after stroke: a cohort study. <i>BMC Neurology</i> , 2020 , 20, 181	3.1	
10	Olle HI Lectureship 2019: The changing world of stroke rehabilitation. <i>Journal of Rehabilitation Medicine</i> , 2020 , 52, jrm00053	3.4	
9	Commentary on "Strategies for rehabilitation professionals to move evidence-based knowledge into practice: a systematic review". <i>Journal of Rehabilitation Medicine</i> , 2009 , 41, 1033	3.4	
8	Transforming self-reported outcomes from a stroke register to the modified Rankin Scale: a cross-sectional, explorative study. <i>Scientific Reports</i> , 2020 , 10, 17215	4.9	
7	Development of a short-form Swedish version of the Montreal Cognitive Assessment (s-MoCA-SWE): protocol for a cross-sectional study. <i>BMJ Open</i> , 2021 , 11, e049035	3	
6	Clinical interpretation and cutoff scores for manual ability measured by the ABILHAND questionnaire in people with stroke. <i>Topics in Stroke Rehabilitation</i> , 2021 , 1-11	2.6	
5	Pre-stroke Physical Inactivity and Stroke Severity in Male and Female Patients <i>Frontiers in Neurology</i> , 2022 , 13, 831773	4.1	
4	Experiences, needs, and preferences for follow-up after stroke perceived by people with stroke and healthcare professionals: A focus group study 2019 , 14, e0223338		
3	Experiences, needs, and preferences for follow-up after stroke perceived by people with stroke and healthcare professionals: A focus group study 2019 , 14, e0223338		
2	Experiences, needs, and preferences for follow-up after stroke perceived by people with stroke and healthcare professionals: A focus group study 2019 , 14, e0223338		
1	Experiences, needs, and preferences for follow-up after stroke perceived by people with stroke and healthcare professionals: A focus group study 2019 , 14, e0223338		