Eibert R Heerdink

List of Publications by Citations

Source: https://exaly.com/author-pdf/5242004/eibert-r-heerdink-publications-by-citations.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

5,360 62 38 220 h-index g-index citations papers 6,098 5.42 233 3.4 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
220	Patient adherence to medical treatment: a review of reviews. <i>BMC Health Services Research</i> , 2007 , 7, 55	2.9	362
219	NSAIDs associated with increased risk of congestive heart failure in elderly patients taking diuretics. <i>Archives of Internal Medicine</i> , 1998 , 158, 1108-12		199
218	Association of risk of abnormal bleeding with degree of serotonin reuptake inhibition by antidepressants. <i>Archives of Internal Medicine</i> , 2004 , 164, 2367-70		173
217	Effect of a pharmacist-led intervention on diuretic compliance in heart failure patients: a randomized controlled study. <i>Journal of Cardiac Failure</i> , 2003 , 9, 404-11	3.3	148
216	Association of nonsteroidal anti-inflammatory drugs with first occurrence of heart failure and with relapsing heart failure: the Rotterdam Study. <i>Archives of Internal Medicine</i> , 2002 , 162, 265-70		146
215	Rate and determinants of 10-year persistence with antihypertensive drugs. <i>Journal of Hypertension</i> , 2005 , 23, 2101-7	1.9	141
214	Indications for antidepressant drug prescribing in general practice in the Netherlands. <i>Journal of Affective Disorders</i> , 2007 , 98, 109-15	6.6	126
213	Unpredictable bias when using the missing indicator method or complete case analysis for missing confounder values: an empirical example. <i>Journal of Clinical Epidemiology</i> , 2010 , 63, 728-36	5.7	100
212	Effectiveness of interventions by community pharmacists to improve patient adherence to chronic medication: a systematic review. <i>Annals of Pharmacotherapy</i> , 2005 , 39, 319-28	2.9	93
211	Depressive symptoms in subjects with diagnosed and undiagnosed type 2 diabetes. <i>Psychosomatic Medicine</i> , 2007 , 69, 300-5	3.7	86
210	Initiation of antidepressant therapy: do patients follow the GPR prescription?. <i>British Journal of General Practice</i> , 2009 , 59, 81-7	1.6	85
209	Incidence and determinants of long-term use of antidepressants. <i>European Journal of Clinical Pharmacology</i> , 2004 , 60, 57-61	2.8	84
208	Detecting drug-drug interactions using a database for spontaneous adverse drug reactions: an example with diuretics and non-steroidal anti-inflammatory drugs. <i>European Journal of Clinical Pharmacology</i> , 2000 , 56, 733-8	2.8	84
207	Application of the ATC/DDD methodology to monitor antibiotic drug use. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 1998 , 17, 20-4	5.3	77
206	The association between antidepressant use and disturbances in glucose homeostasis: evidence from spontaneous reports. <i>European Journal of Clinical Pharmacology</i> , 2008 , 64, 531-8	2.8	72
205	Construction of drug treatment episodes from drug-dispensing histories is influenced by the gap length. <i>Journal of Clinical Epidemiology</i> , 2010 , 63, 422-7	5.7	69
204	Generic substitution of antihypertensive drugs: does it affect adherence?. <i>Annals of Pharmacotherapy</i> , 2006 , 40, 15-20	2.9	68

(2011-2009)

203	Risk of cerebrovascular events in elderly users of antipsychotics. <i>Journal of Psychopharmacology</i> , 2009 , 23, 909-14	4.6	65	
202	Impact of coaching by community pharmacists on drug attitude of depressive primary care patients and acceptability to patients; a randomized controlled trial. <i>European Neuropsychopharmacology</i> , 2003 , 13, 1-9	1.2	64	
201	Polymorphisms of the LEP- and LEPR gene and obesity in patients using antipsychotic medication. <i>Journal of Clinical Psychopharmacology</i> , 2009 , 29, 21-5	1.7	61	
200	Pharmacotherapy for the treatment of aggressive behavior in general adult psychiatry: A systematic review. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 1013-24	4.6	60	
199	The prevalence and incidence of medicinal cannabis on prescription in The Netherlands. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1575-80	2.8	59	
198	Furthering patient adherence: a position paper of the international expert forum on patient adherence based on an internet forum discussion. <i>BMC Health Services Research</i> , 2008 , 8, 47	2.9	58	
197	Risk of hip/femur fractures in patients using antipsychotics. <i>Bone</i> , 2005 , 37, 864-70	4.7	58	
196	Is aspirin useful in patients on lithium? A pharmacoepidemiological study related to bipolar disorder. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2010 , 82, 9-14	2.8	56	
195	Patient risk profiles and practice variation in nonadherence to antidepressants, antihypertensives and oral hypoglycemics. <i>BMC Health Services Research</i> , 2007 , 7, 51	2.9	53	
194	Lithium use and the risk of fractures. <i>Bone</i> , 2007 , 40, 1252-8	4.7	53	
193	A pharmacy-based coaching program to improve adherence to antidepressant treatment among primary care patients. <i>Psychiatric Services</i> , 2005 , 56, 487-9	3.3	51	
192	Changes in prescribed drug doses after market introduction. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 447-53	2.6	49	
191	Psychotropic drug use in intellectually disabled group-home residents with behavioural problems. <i>Pharmacopsychiatry</i> , 2002 , 35, 19-23	2	45	
190	Discontinuation symptoms in users of selective serotonin reuptake inhibitors in clinical practice: tapering versus abrupt discontinuation. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 303-7	2.8	43	
189	Antidepressant use and off-label prescribing in children and adolescents in Dutch general practice (2001-2005). <i>Pharmacoepidemiology and Drug Safety</i> , 2007 , 16, 1054-62	2.6	42	
188	No increased incidence of diabetes in antidepressant users. <i>International Clinical Psychopharmacology</i> , 2007 , 22, 382-6	2.2	42	
187	Refill persistence with chronic medication assessed from a pharmacy database was influenced by method of calculation. <i>Journal of Clinical Epidemiology</i> , 2006 , 59, 11-7	5.7	42	
186	Use of serotonergic drugs and the risk of bleeding. <i>Clinical Pharmacology and Therapeutics</i> , 2011 , 89, 89-96	6.1	41	

185	A prospective follow-up study of 5669 users of lansoprazole in daily practice. <i>Alimentary Pharmacology and Therapeutics</i> , 1997 , 11, 887-97	6.1	41
184	Proposal of Standardization to Assess Adherence With Medication Records: Methodology Matters. <i>Annals of Pharmacotherapy</i> , 2016 , 50, 360-8	2.9	41
183	Information on drug use in the elderly: a comparison of pharmacy, general-practitioner and patient data. <i>International Journal of Clinical Pharmacy</i> , 1995 , 17, 20-4		39
182	Adverse events and the relation with quality of life in adults with intellectual disability and challenging behaviour using psychotropic drugs. <i>Research in Developmental Disabilities</i> , 2016 , 49-50, 13	3-2 1 7	36
181	The association between compliance with antihypertensive drugs and modification of antihypertensive drug regimen. <i>Journal of Hypertension</i> , 2004 , 22, 1831-7	1.9	36
180	Haloperidol dose when used as active comparator in randomized controlled trials with atypical antipsychotics in schizophrenia: comparison with officially recommended doses. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 897-903	4.6	36
179	Trends in incidence and characteristics of children, adolescents, and adults initiating immediate- or extended-release methylphenidate or atomoxetine in the Netherlands during 2001-2006. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2010 , 20, 55-61	2.9	35
178	Increased psychotropic drug consumption by children in the Netherlands during 1995-2001 is caused by increased use of methylphenidate by boys. <i>European Journal of Clinical Pharmacology</i> , 2004 , 60, 377-9	2.8	35
177	The decision to continue or discontinue treatment: experiences and beliefs of users of selective serotonin-reuptake inhibitors in the initial monthsa qualitative study. <i>Research in Social and Administrative Pharmacy</i> , 2011 , 7, 134-50	2.9	34
176	Refill compliance in type 2 diabetes mellitus: a predictor of switching to insulin therapy?. <i>Pharmacoepidemiology and Drug Safety</i> , 2003 , 12, 121-7	2.6	33
175	Ineffectiveness and adverse events of nitrofurantoin in women with urinary tract infection and renal impairment in primary care. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1701-7	2.8	32
174	Characteristics and reasons associated with nonacceptance of selective serotonin-reuptake inhibitor treatment. <i>Annals of Pharmacotherapy</i> , 2008 , 42, 218-25	2.9	32
173	Less discontinuation of ADHD drug use since the availability of long-acting ADHD medication in children, adolescents and adults under the age of 45 years in the Netherlands. <i>ADHD Attention Deficit and Hyperactivity Disorders</i> , 2010 , 2, 213-20	3.1	31
172	Determinants of multiple psychotropic drug use in patients with mild intellectual disabilities or borderline intellectual functioning and psychiatric or behavioral disorders. <i>General Hospital Psychiatry</i> , 2001 , 23, 345-9	5.6	31
171	Variability between nursing homes in prevalence of antipsychotic use in patients with dementia. <i>International Psychogeriatrics</i> , 2014 , 26, 363-71	3.4	30
170	Association between ADHD drug use and injuries among children and adolescents. <i>European Child and Adolescent Psychiatry</i> , 2014 , 23, 95-102	5.5	30
169	Polymorphisms of the LEP, LEPR and HTR2C gene: obesity and BMI change in patients using antipsychotic medication in a naturalistic setting. <i>Pharmacogenomics</i> , 2011 , 12, 919-23	2.6	30
168	Evaluation of patientsRexperiences with antidepressants reported by means of a medicine reporting system. <i>European Journal of Clinical Pharmacology</i> , 2007 , 63, 1193-9	2.8	30

(2008-2005)

167	Reasons for switching between antipsychotics in daily clinical practice. <i>Pharmacopsychiatry</i> , 2005 , 38, 122-4	2	30	
166	Antipsychotic drug use and the risk of venous thromboembolism in elderly patients. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 526-30	1.7	29	
165	Duration of antidepressant drug treatment and its influence on risk of relapse/recurrence: immortal and neglected time bias. <i>American Journal of Epidemiology</i> , 2009 , 170, 280-5	3.8	28	
164	An algorithm to identify antidepressant users with a diagnosis of depression from prescription data. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 7-15	2.6	28	
163	Prevalence of patients using drugs metabolized by cytochrome P450 2D6 in different populations: a cross-sectional study. <i>Annals of Pharmacotherapy</i> , 2007 , 41, 408-13	2.9	28	
162	Amiodarone-induced thyroid dysfunction associated with cumulative dose. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 601-6	2.6	28	
161	Evaluation of risk profiles for gastrointestinal and cardiovascular adverse effects in nonselective NSAID and COX-2 inhibitor users: a cohort study using pharmacy dispensing data in The Netherlands. <i>Drug Safety</i> , 2008 , 31, 143-58	5.1	27	
160	Structured Medication Review to Improve Pharmacotherapy in People with Intellectual Disability and Behavioural Problems. <i>Journal of Applied Research in Intellectual Disabilities</i> , 2016 , 29, 346-55	2.2	26	
159	Quantifying antibiotic use in paediatrics: a proposal for neonatal DDDs. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2010 , 29, 1301-3	5.3	26	
158	Effects of NSAIDs on the incidence of hospitalisations for renal dysfunction in users of ACE inhibitors. <i>Drug Safety</i> , 2003 , 26, 983-9	5.1	26	
157	Is new drug prescribing in primary care specialist induced?. BMC Health Services Research, 2009, 9, 6	2.9	24	
156	Antidepressant use before and after initiation of diabetes mellitus treatment. <i>Diabetologia</i> , 2009 , 52, 425-32	10.3	24	
155	Drug-drug interactions as a determinant of elevated lithium serum levels in daily clinical practice. <i>Bipolar Disorders</i> , 2005 , 7, 274-80	3.8	24	
154	Effects of Telephone Counseling Intervention by Pharmacists (TelCIP) on Medication Adherence; Results of a Cluster Randomized Trial. <i>Frontiers in Pharmacology</i> , 2016 , 7, 269	5.6	24	
153	Prevalence and characteristics of psychotropic drug use in institutionalized children and adolescents with mild intellectual disability. <i>Research in Developmental Disabilities</i> , 2013 , 34, 3159-67	2.7	23	
152	PatientsRperceptions of information received at the start of selective serotonin-reuptake inhibitor treatment: implications for community pharmacy. <i>Annals of Pharmacotherapy</i> , 2009 , 43, 642-9	2.9	23	
151	The course of behavioral problems in elderly nursing home patients with dementia when treated with antipsychotics. <i>International Psychogeriatrics</i> , 2009 , 21, 931-40	3.4	23	
150	Influence of antidepressants on glycaemic control in patients with diabetes mellitus. Pharmacoepidemiology and Drug Safety, 2008, 17, 577-86	2.6	23	

149	Adverse events in users of sertraline: results from an observational study in psychiatric practice in The Netherlands. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 655-62	2.6	23
148	Use of sympathomimetic drugs leads to increased risk of hospitalization for arrhythmias in patients with congestive heart failure. <i>Archives of Internal Medicine</i> , 2000 , 160, 2477-80		22
147	Spontaneous lapses in dosing during chronic treatment with selective serotonin reuptake inhibitors. <i>British Journal of Psychiatry</i> , 2001 , 179, 519-22	5.4	22
146	The relation between risk-taking behavior and alcohol use in young adults is different for men and women. <i>Drug and Alcohol Dependence</i> , 2015 , 155, 222-7	4.9	21
145	Determinants of the range of drugs prescribed in general practice: a cross-sectional analysis. <i>BMC Health Services Research</i> , 2007 , 7, 132	2.9	21
144	The association between antidepressant use and hypoglycaemia in diabetic patients: a nested case-control study. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 336-44	2.6	21
143	Potential bias in pharmacoepidemiological studies due to the length of the drug free period: a study on antidepressant drug use in adults in the Netherlands. <i>Pharmacoepidemiology and Drug Safety</i> , 2006 , 15, 338-43	2.6	21
142	Patterns of pharmacotherapy in patients hospitalised for congestive heart failure. <i>European Journal of Heart Failure</i> , 2003 , 5, 195-200	12.3	21
141	Visualizing Pharmacological Activities of Antidepressants: A Novel Approach. <i>The Open Pharmacology Journal</i> , 2008 , 2, 54-62	0.3	21
140	Association between hospitalization and discontinuity of medication therapy used in the community setting in the Netherlands. <i>Annals of Pharmacotherapy</i> , 2008 , 42, 933-9	2.9	20
139	Pharmacogenetics of glucose-lowering drug treatment: a systematic review. <i>Molecular Diagnosis and Therapy</i> , 2007 , 11, 291-302	4.5	20
138	Does the use of an electronic reminder device with or without counseling improve adherence to lipid-lowering treatment? The results of a randomized controlled trial. <i>Frontiers in Pharmacology</i> , 2013 , 4, 69	5.6	19
137	Representation of people with intellectual disabilities in randomised controlled trials on antipsychotic treatment for behavioural problems. <i>Journal of Intellectual Disability Research</i> , 2011 , 55, 650-64	3.2	19
136	Association between LEP and LEPR gene polymorphisms and dyslipidemia in patients using atypical antipsychotic medication. <i>Psychiatric Genetics</i> , 2010 , 20, 311-6	2.9	19
135	Genetic variation in the renin-angiotensin system modifies the beneficial effects of ACE inhibitors on the risk of diabetes mellitus among hypertensives. <i>Journal of Human Hypertension</i> , 2008 , 22, 774-80	2.6	19
134	The effect of pharmacotherapy audit meetings on early new drug prescribing by general practitioners. <i>Annals of Pharmacotherapy</i> , 2007 , 41, 319-24	2.9	19
133	Does the length of the first antidepressant treatment episode influence risk and time to a second episode?. <i>Journal of Clinical Psychopharmacology</i> , 2009 , 29, 69-72	1.7	18
132	Linking community pharmacy dispensing data to prescribing data of general practitioners. <i>BMC Medical Informatics and Decision Making</i> , 2006 , 6, 18	3.6	18

(2005-2006)

131	Between-country variation in the utilization of antihypertensive agents: Guidelines and clinical practice. <i>Journal of Human Hypertension</i> , 2006 , 20, 917-22	2.6	18
130	Reported Adverse Drug Reactions in Children and Adolescents Treated with Antipsychotics. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2019 , 29, 124-132	2.9	17
129	The use of driving impairing medicines: a European survey. <i>European Journal of Clinical Pharmacology</i> , 2009 , 65, 1139-47	2.8	17
128	Instructions for clinical and biomarker monitoring in the Summary of Product Characteristics (SmPC) for psychotropic drugs: Overview and applicability in clinical practice. <i>Journal of Psychopharmacology</i> , 2015 , 29, 1248-54	4.6	16
127	Association between lithium serum level, mood state, and patient-reported adverse drug reactions during long-term lithium treatment: a naturalistic follow-up study. <i>Bipolar Disorders</i> , 2009 , 11, 434-40	3.8	16
126	Combined HTR2C-LEP genotype as a determinant of obesity in patients using antipsychotic medication. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 702-5	1.7	16
125	Initial non-compliance with antihypertensive monotherapy is followed by complete discontinuation of antihypertensive therapy. <i>Pharmacoepidemiology and Drug Safety</i> , 2006 , 15, 587-93	2.6	16
124	The impact of hospitalisation on the initiation and long-term use of benzodiazepines. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 291-5	2.8	16
123	Pharmacogenetic testing to predict antipsychotic-induced weight gain: a systematic review. <i>Pharmacogenomics</i> , 2011 , 12, 1213-27	2.6	15
122	Estimates of statin discontinuation rates are influenced by exposure and outcome definitions. <i>Annals of Pharmacotherapy</i> , 2011 , 45, 576-81	2.9	15
121	Differences in body mass index z-scores and weight status in a Dutch pediatric psychiatric population with and without use of second-generation antipsychotics. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2012 , 22, 166-73	2.9	15
120	Changes in outpatient lithium treatment in the Netherlands during 1996-2005. <i>Journal of Affective Disorders</i> , 2008 , 111, 94-9	6.6	15
119	Seclusion and the use of antipsychotics in hospitalized psychiatric patients. <i>Psychology, Crime and Law</i> , 2005 , 11, 489-495	1.4	15
118	Medicinal Cannabis on Prescription in The Netherlands: Statistics for 2003-2016. <i>Cannabis and Cannabinoid Research</i> , 2018 , 3, 54-55	4.6	15
117	Seasonal patterns of initiating antidepressant therapy in general practice in the Netherlands during 2002-2007. <i>Journal of Affective Disorders</i> , 2010 , 122, 208-12	6.6	14
116	Aggression of psychiatric patients associated with the use of as-needed medication. <i>Pharmacopsychiatry</i> , 2007 , 40, 25-9	2	14
115	Characteristics of diarrhoea in 10,008 users of lansoprazole in daily practice: which co-factors contribute?. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 703-8	2.6	14
114	Psychoactive drug prescribing by Dutch child and adolescent psychiatrists. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2005 , 94, 1484-1487	3.1	14

113	The Effect of Glycopyrrolate on Nocturnal Sialorrhea in Patients Using Clozapine: A Randomized, Crossover, Double-Blind, Placebo-Controlled Trial. <i>Journal of Clinical Psychopharmacology</i> , 2017 , 37, 155	5-161	13
112	No difference in between-country variability in use of newly approved orphan and non- orphan medicinal productsa pilot study. <i>Orphanet Journal of Rare Diseases</i> , 2009 , 4, 27	4.2	13
111	Clinical relevance of patients with epilepsy included in clinical trials. <i>Epilepsia</i> , 2008 , 49, 1479-80	6.4	13
110	The association between exposure to COX-2 inhibitors and schizophrenia deterioration. A nested case-control study. <i>Pharmacopsychiatry</i> , 2007 , 40, 111-5	2	13
109	PatientsRgeneral satisfaction with telephone counseling by pharmacists and effects on satisfaction with information and beliefs about medicines: Results from a cluster randomized trial. <i>Patient Education and Counseling</i> , 2015 , 98, 797-804	3.1	12
108	Lazy sunday afternoons: the negative impact of interruptions in patientsRdaily routine on adherence to oral antidiabetic medication. A multilevel analysis of electronic monitoring data. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1599-606	2.8	12
107	Low treatment adherence with antipsychotics is associated with relapse in psychotic disorders within six months after discharge. <i>Pharmacopsychiatry</i> , 2010 , 43, 221-4	2	12
106	PatientsRperceptions and illness severity at start of antidepressant treatment in general practice. <i>International Journal of Pharmacy Practice</i> , 2010 , 18, 217-25	1.7	12
105	The impact of environmental temperature on lithium serum levels. <i>Bipolar Disorders</i> , 2007 , 9, 603-8	3.8	12
104	Should rosuvastatin be withdrawn from the market?. <i>Lancet, The</i> , 2004 , 364, 1577; author reply 1578-9	40	12
103	Long-term persistence and adherence with non-vitamin K oral anticoagulants in patients with atrial fibrillation and their associations with stroke risk. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2021 , 7, f72-f80	6.4	12
102	Rates and determinants of reinitiating antihypertensive therapy after prolonged stoppage: a population-based study. <i>Journal of Hypertension</i> , 2007 , 25, 689-97	1.9	11
101	Suicide Behavior Before and After the Start with Antidepressants: A High Persistent Risk in the First Month of Treatment Among the Young. <i>International Journal of Neuropsychopharmacology</i> , 2015 , 19,	5.8	10
100	Experiences with medical cannabis in the treatment of veterans with PTSD: Results from a focus group discussion. <i>European Neuropsychopharmacology</i> , 2020 , 36, 244-254	1.2	10
99	Movement Disorders in Adults With Intellectual Disability and Behavioral Problems Associated With Use of Antipsychotics. <i>Journal of Clinical Psychopharmacology</i> , 2016 , 36, 308-13	1.7	10
98	Diabetes and Glucose Dysregulation and Transition to Delirium in ICU Patients. <i>Critical Care Medicine</i> , 2018 , 46, 1444-1449	1.4	10
97	Effects of a TELephone Counselling Intervention by Pharmacist (TelCIP) on medication adherence, patient beliefs and satisfaction with information for patients starting treatment: study protocol for a cluster randomized controlled trial. <i>BMC Health Services Research</i> , 2014 , 14, 219	2.9	10
96	Methylphenidate: use in daily practice. <i>International Journal of Clinical Pharmacy</i> , 2005 , 27, 197-201		10

(2006-2001)

95	Prescribing patterns in patients using new antidepressants. <i>British Journal of Clinical Pharmacology</i> , 2001 , 51, 181-3	3.8	10
94	Differences in ADHD medication usage patterns in children and adolescents from different cultural backgrounds in the Netherlands. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015 , 50, 1153-62	4.5	9
93	Attitudes towards medication use in a general population of adolescents. <i>European Journal of Pediatrics</i> , 2014 , 173, 483-8	4.1	9
92	Variable access to clopidogrel in a harmonized EU market. <i>Value in Health</i> , 2008 , 11, 989-95	3.3	9
91	Urine osmolality, cyclic AMP and aquaporin-2 in urine of patients under lithium treatment in response to water loading followed by vasopressin administration. <i>European Journal of Pharmacology</i> , 2007 , 566, 50-7	5.3	9
90	Need for medicine-based evidence in pharmacotherapy. <i>British Journal of Psychiatry</i> , 2004 , 184, 452	5.4	9
89	Evaluation of clarity of presentation and applicability of monitoring instructions for patients using lithium in clinical practice guidelines for treatment of bipolar disorder. <i>Bipolar Disorders</i> , 2018 , 20, 708-708-708	7 3 8	9
88	Monitoring of patients treated with lithium for bipolar disorder: an international survey. <i>International Journal of Bipolar Disorders</i> , 2018 , 6, 12	5.4	8
87	The association between patient-reported drug taking and gaps and overlaps in antidepressant drug dispensing. <i>Annals of Pharmacotherapy</i> , 2010 , 44, 1755-61	2.9	8
86	Patient characteristics associated with prescribing of a newly introduced drug: the case of rofecoxib. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 157-9	2.8	8
85	The trade-off between cardiovascular and gastrointestinal effects of rofecoxib. <i>Pharmacoepidemiology and Drug Safety</i> , 2005 , 14, 437-41	2.6	8
84	Glucose variability during delirium in diabetic and non-diabetic intensive care unit patients: A prospective cohort study. <i>PLoS ONE</i> , 2018 , 13, e0205637	3.7	8
83	Effects of a Proactive Interdisciplinary Self-Management (PRISMA) program on medication adherence in patients with type 2 diabetes in primary care: a randomized controlled trial. <i>Patient Preference and Adherence</i> , 2019 , 13, 749-759	2.4	7
82	Safety of clozapine use during pregnancy: Analysis of international pharmacovigilance data. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 725-735	2.6	7
81	Somatic health care utilization by patients treated for substance use disorders. <i>Drug and Alcohol Dependence</i> , 2017 , 178, 277-284	4.9	7
80	Risk of acute coronary syndrome in elderly users of antipsychotic drugs: a nested case-control study. <i>Heart</i> , 2012 , 98, 1166-71	5.1	7
79	Impact analysis of the discontinuation of reimbursement: the case of oral contraceptives. <i>Contraception</i> , 2008 , 78, 399-404	2.5	7
78	Which pharmacists contribute to high-level pharmacotherapy audit meetings with general practitioners?. <i>Annals of Pharmacotherapy</i> , 2006 , 40, 1640-6	2.9	7

77	antihypertensive drug classes: Simple Additive Weighting (SAW) and Technique for Order Preference by Similarity to an Ideal Solution (TOPSIS). <i>American Journal of Cardiovascular Drugs</i> , 2006 , 6, 251-8	4	7
76	Benzodiazepine pathways in the chronically ill. <i>Pharmacoepidemiology and Drug Safety</i> , 1999 , 8, 325-30	2.6	7
75	Antipsychotic drug use associated with urinary tract infections in older women. <i>Maturitas</i> , 2017 , 98, 46-	5 9	6
74	Medication self-management support for people with diabetes and low health literacy: A needs assessment. <i>PLoS ONE</i> , 2020 , 15, e0232022	3.7	6
73	Association Between Urinary Tract Infections and Antipsychotic Drug Use in Older Adults. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 296-301	1.7	6
72	Less medication switching after initial start with atypical antipsychotics. <i>European Neuropsychopharmacology</i> , 2004 , 14, 1-5	1.2	6
71	Short-acting parenteral antipsychotics drive choice for classical versus atypical agents. <i>European Journal of Clinical Pharmacology</i> , 2003 , 58, 757-60	2.8	6
70	Determinants of headache in lansoprazole users in The Netherlands: results from a nested case-control study. <i>Drug Safety</i> , 2002 , 25, 287-95	5.1	6
69	Is sexual dysfunctioning a reason for discontinuation in users of selective serotonin reuptake inhibitors?. <i>Journal of Clinical Psychopharmacology</i> , 2002 , 22, 96-8	1.7	6
68	Waste-minimising measures to achieve sustainable supply and use of medication. <i>Sustainable Chemistry and Pharmacy</i> , 2021 , 20, 100400	3.9	6
67	Associations Between Personality Traits and Adherence to Antidepressants Assessed Through Self-Report, Electronic Monitoring, and Pharmacy Dispensing Data: A Pilot Study. <i>Journal of Clinical Psychopharmacology</i> , 2016 , 36, 465-71	1.7	5
66	The association between prescription change frequency, chronic disease score and hospital admissions: a case control study. <i>BMC Pharmacology & Doctor</i> 2013, 14, 39	2.6	5
65	Transitions from General Practitioner to Psychiatrist Care (or vice versa) during a first antidepressant treatment episode. <i>Pharmacopsychiatry</i> , 2010 , 43, 179-83	2	5
64	Trials assessing parmacotherapeutic management of aggression in psychiatric patients: comparability with clinical practice. <i>Pharmacopsychiatry</i> , 2010 , 43, 205-9	2	5
63	Evaluation of available treatment guidelines for the management of lithium intoxication. <i>Therapeutic Drug Monitoring</i> , 2009 , 31, 247-60	3.2	5
62	Influence of antidepressant use on glycemic control in patients with diabetes mellitus: an open-label comparative study. <i>Journal of Clinical Psychopharmacology</i> , 2009 , 29, 405-8	1.7	5
61	Impaired glucose homeostasis after imipramine intake in a diabetic patient. <i>Journal of Clinical Psychopharmacology</i> , 2005 , 25, 621-3	1.7	5
60	Channelling new antidepressants to problem patients may be factor in fatal toxicity. <i>BMJ, The</i> , 2003 , 326, 600	5.9	5

59	Development of eHOME, a Mobile Instrument for Reporting, Monitoring, and Consulting Drug-Related Problems in Home Care: Human-Centered Design Study. <i>JMIR Human Factors</i> , 2018 , 5, e10) ^{2.5}	5
58	Compliance with the guidelines for laboratory monitoring of patients treated with lithium: A retrospective follow-up study among ambulatory patients in the Netherlands. <i>Bipolar Disorders</i> , 2019 , 21, 419-427	3.8	5
57	Monitoring of Metabolic, Cardiac, and Endocrine Indicators in Youth Treated With Antipsychotics as Reported by Health Care Professionals. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 489-493	1.7	5
56	Dispensing of psychotropic medication among 400,000 immigrants in The Netherlands. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017 , 52, 963-977	4.5	4
55	Unbound Fraction of Clozapine Significantly Decreases with Elevated Plasma Concentrations of the Inflammatory Acute-Phase Protein Alpha-1-Acid Glycoprotein. <i>Clinical Pharmacokinetics</i> , 2019 , 58, 1069-	-1075	4
54	Risk of Hospitalization for Hypoglycemia in Older Patients with Diabetes Using Antipsychotic Drugs. <i>American Journal of Geriatric Psychiatry</i> , 2015 , 23, 1144-53	6.5	4
53	Guidance by physicians and pharmacists during antidepressant therapy: patientsRneeds and suggestions for improvement. <i>BMC Psychiatry</i> , 2017 , 17, 388	4.2	4
52	Severe Cognitive Impairment Associated With a High Free But Therapeutic Total Concentration of Valproic Acid Due to Hypoalbuminemia in an Older Patient With Bipolar Disorder. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 265-268	1.7	4
51	Discontinuation of somatic medication during psychiatric hospitalization. <i>Annals of Pharmacotherapy</i> , 2014 , 48, 1415-24	2.9	4
50	Fluorescence of neutrophil granulocytes as a biomarker for clozapine use. <i>European Neuropsychopharmacology</i> , 2013 , 23, 1408-13	1.2	4
49	The association between serotonin 2C receptor polymorphisms and weight gain and eating behavior in patients using mirtazapine: a prospective follow-up study. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 207-9	1.7	4
48	Determinants of psychotropic drug usage in a general intensive care unit. <i>General Hospital Psychiatry</i> , 1998 , 20, 371-6	5.6	4
47	The association between concomitant use of serotonergic antidepressants and lithium-induced polyuria. A multicenter medical chart review study. <i>Pharmacopsychiatry</i> , 2008 , 41, 129-33	2	4
46	Determinants of compliance with methylphenidate therapy in children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2006 , 95, 1674-6	3.1	4
45	Safety review in 10,008 users of lansoprazole in daily practice. <i>Pharmacoepidemiology and Drug Safety</i> , 2000 , 9, 383-91	2.6	4
44	Prevalence of Psychotropic Medication Use Among Dutch Military Personnel Between 2003 and 2012 and Its Comparison to the Dutch General Population. <i>Military Medicine</i> , 2017 , 182, e1584-e1588	1.3	3
43	The impact of ethnic density on dispensing of antipsychotic and antidepressant medication among immigrants in the Netherlands. <i>Social Science and Medicine</i> , 2018 , 211, 87-94	5.1	3
42	The association between CYP2D6 genotype and switching antipsychotic medication to clozapine. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1927-32	2.8	3

41	Discontinuation of anticoagulant care during admission to a psychiatric hospital. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1025-9	2.8	3
40	Clozapine re-exposure after dilated cardiomyopathy. <i>BMJ Case Reports</i> , 2017 , 2017,	0.9	3
39	Prevalence of medication use for somatic disease in institutionalized psychiatric patients. <i>Pharmacopsychiatry</i> , 2013 , 46, 274-80	2	3
38	Influence of therapeutic complexity on medication adherence in the Netherlands. <i>Archives of Internal Medicine</i> , 2011 , 171, 864-5; author reply 865		3
37	Prescriber adoption of newly approved selective COX-2 inhibitors. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 1168-74	2.6	3
36	Changes in the defined daily dose; CYP2D6/CYP3A metabolism as an indicator for dose-setting problems. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 243-6	2.8	3
35	Long-term therapy with spironolactone. International Journal of Clinical Pharmacy, 2001, 23, 132-4		3
34	Effect of reimbursement restriction policy on the use of benzodiazepines in the Netherlands: an interrupted time series analysis. <i>BMJ Open</i> , 2019 , 9, e029148	3	3
33	Effects of the Proactive interdisciplinary self-management (PRISMA) program on self-reported and clinical outcomes in type 2 diabetes: a pragmatic randomized controlled trial. <i>BMC Endocrine Disorders</i> , 2019 , 19, 139	3.3	3
32	Discontinuation Rates of Antidepressant Use by Dutch Soldiers. <i>Military Medicine</i> , 2019 , 184, 868-874	1.3	2
31	Medication Discontinuation in Patients After Discharge From a Psychiatric Hospital. <i>Annals of Pharmacotherapy</i> , 2015 , 49, 1085-95	2.9	2
30	Comparing episodes of antidepressants use with intermittent episodes of no use: A higher relative risk of suicide attempts but not of suicide at young age. <i>Journal of Psychopharmacology</i> , 2016 , 30, 1000-	. 4 .6	2
29	Antidepressant use in pregnancy: knowledge transfer and translation of research findings. <i>Journal of Evaluation in Clinical Practice</i> , 2015 , 21, 579-83	2.5	2
28	Association between HTR2C polymorphisms and obesity in patients without antipsychotic drugs. <i>Journal of Clinical Psychopharmacology</i> , 2012 , 32, 715-40	1.7	2
27	Early identification of long-term poor adherence in ambulatory patients. <i>Annals of Pharmacotherapy</i> , 2006 , 40, 2277-8	2.9	2
26	Effects of the Proactive Interdisciplinary Self-Management (PRISMA) Program on Online Care Platform Usage in Patients with Type 2 Diabetes in Primary Care: A Randomized Controlled Trial. <i>Journal of Diabetes Research</i> , 2020 , 2020, 5013142	3.9	2
25	Medication self-management: Considerations and decisions by older people living at home. <i>Research in Social and Administrative Pharmacy</i> , 2020 , 18, 2410-2410	2.9	2
24	Predicting rehospitalization in patients treated with antipsychotics: a prospective observational study. <i>Therapeutic Advances in Psychopharmacology</i> , 2018 , 8, 213-229	4.9	2

23	Patient monitoring: the hidden costs of treatment with antipsychotics. <i>British Journal of Clinical Pharmacology</i> , 2016 , 82, 570-2	3.8	1
22	Clozapine-induced hypersalivation: the association between quantification, perceived burden and treatment satisfaction reported by patients. <i>Therapeutic Advances in Psychopharmacology</i> , 2017 , 7, 20	9-248	1
21	Association between concomitant psychiatric drug use, and patients Roeliefs about and persistence with chronic cardiovascular medication. <i>International Journal of Clinical Practice</i> , 2015 , 69, 328-35	2.9	1
20	Lithium: farmacologie in de psychiatrie. <i>Neuropraxis</i> , 2008 , 12, 147-153	Ο	1
19	Factors associated with non-response in proton pump inhibitor users: a study of lansoprazole therapy. <i>International Journal of Clinical Pharmacy</i> , 2001 , 23, 107-10		1
18	Continued use of the proton-pump inhibitor lansoprazole following Helicobacter pylori eradication. <i>European Journal of Clinical Pharmacology</i> , 2000 , 55, 773-6	2.8	1
17	Associations Between Characteristics of Web-Based Diabetes News and Readers Sentiments: Observational Study in the Netherlands. <i>Journal of Medical Internet Research</i> , 2019 , 21, e14554	7.6	1
16	The effect of non-adherence to antipsychotic treatment on rehospitalization in patients with psychotic disorders. <i>Therapeutic Advances in Psychopharmacology</i> , 2021 , 11, 20451253211027449	4.9	1
15	Monitoring of Adverse Drug Reaction-Related Parameters in Children and Adolescents Treated With Antipsychotic Drugs in Psychiatric Outpatient Clinics. <i>Frontiers in Psychiatry</i> , 2021 , 12, 640377	5	1
14	Predicting Unsuccessful Clozapine Treatment After First Use in Adult Patients With Psychotic Disorders. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 604-608	1.7	1
13	Home Care PatientsRExperiences with Home Care NursesRSupport in Medication Adherence. <i>Patient Preference and Adherence</i> , 2021 , 15, 1929-1940	2.4	1
12	Psychotropic medication use in former ICU patients with mental health problems: A prospective observational follow-up study. <i>Journal of Critical Care</i> , 2020 , 59, 112-117	4	О
11	The impact of health literacy on beliefs about medication in a Dutch medication-using population. <i>European Journal of Clinical Pharmacology</i> , 2021 , 77, 1219-1224	2.8	O
10	Clarity and applicability of adverse drug reaction-related monitoring instructions in clinical practice guidelines for children and adolescents treated with antipsychotic drugs: a review of six clinical practice guidelines <i>BMJ Open</i> , 2022 , 12, e058940	3	O
9	Pharmacological Management of Aggressive Behavior in Psychiatric Patients 2017, 1-15		
8	The association between lifting an administrative restriction on antidepressant dispensing and treatment patterns in Iceland. <i>Health Policy</i> , 2013 , 111, 193-9	3.2	
7	Nonmedical Use of Attention-Deficit/Hyperactivity Disorder Medication Among Secondary School Students in The Netherlands. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2015 , 25, 649-52	2.9	
6	Combined HTR2C-LEP and HTR2C-LEPR genotypes as a determinant for obesity in patients without antipsychotic drugs. <i>Journal of Clinical Psychopharmacology</i> , 2014 , 34, 266-8	1.7	

5	Unexpected lower prevalence of depression in patients with diabetes: selection bias in a waiting room population. <i>Epidemiology</i> , 2009 , 20, 419-23	3.1
4	Gardarsdottir et al. Respond to "Assessing Treatment Effects With Observational Analyses". <i>American Journal of Epidemiology</i> , 2009 , 170, 288-288	3.8
3	Use of antidepressants in pregnancy does not increase risk of postpartum haemorrhage. <i>Evidence-Based Mental Health</i> , 2008 , 11, 125	11.1
2	Correlated measures in longitudinal analysis of daily drug use patterns in a general intensive care unit. <i>Pharmacoepidemiology and Drug Safety</i> , 2004 , 13, 395-8	2.6
1	They promised this ten years ago. Effects of diabetes news characteristics on patientsRperceptions and attitudes towards medical innovations and therapy adherence. <i>PLoS ONE</i> , 2021 , 16, e0255587	3.7