Eibert R Heerdink

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/5242004/eibert-r-heerdink-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

5,360 62 38 220 h-index g-index citations papers 6,098 5.42 233 3.4 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
220	Clarity and applicability of adverse drug reaction-related monitoring instructions in clinical practice guidelines for children and adolescents treated with antipsychotic drugs: a review of six clinical practice guidelines <i>BMJ Open</i> , 2022 , 12, e058940	3	O
219	Waste-minimising measures to achieve sustainable supply and use of medication. <i>Sustainable Chemistry and Pharmacy</i> , 2021 , 20, 100400	3.9	6
218	Long-term persistence and adherence with non-vitamin K oral anticoagulants in patients with atrial fibrillation and their associations with stroke risk. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2021 , 7, f72-f80	6.4	12
217	The effect of non-adherence to antipsychotic treatment on rehospitalization in patients with psychotic disorders. <i>Therapeutic Advances in Psychopharmacology</i> , 2021 , 11, 20451253211027449	4.9	1
216	Monitoring of Adverse Drug Reaction-Related Parameters in Children and Adolescents Treated With Antipsychotic Drugs in Psychiatric Outpatient Clinics. <i>Frontiers in Psychiatry</i> , 2021 , 12, 640377	5	1
215	The impact of health literacy on beliefs about medication in a Dutch medication-using population. <i>European Journal of Clinical Pharmacology</i> , 2021 , 77, 1219-1224	2.8	О
214	They promised this ten years ago. Effects of diabetes news characteristics on patientsRperceptions and attitudes towards medical innovations and therapy adherence. <i>PLoS ONE</i> , 2021 , 16, e0255587	3.7	
213	Home Care PatientsRExperiences with Home Care NursesRSupport in Medication Adherence. <i>Patient Preference and Adherence</i> , 2021 , 15, 1929-1940	2.4	1
212	Safety of clozapine use during pregnancy: Analysis of international pharmacovigilance data. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 725-735	2.6	7
211	Experiences with medical cannabis in the treatment of veterans with PTSD: Results from a focus group discussion. <i>European Neuropsychopharmacology</i> , 2020 , 36, 244-254	1.2	10
210	Psychotropic medication use in former ICU patients with mental health problems: A prospective observational follow-up study. <i>Journal of Critical Care</i> , 2020 , 59, 112-117	4	O
209	Medication self-management support for people with diabetes and low health literacy: A needs assessment. <i>PLoS ONE</i> , 2020 , 15, e0232022	3.7	6
208	Effects of the Proactive Interdisciplinary Self-Management (PRISMA) Program on Online Care Platform Usage in Patients with Type 2 Diabetes in Primary Care: A Randomized Controlled Trial. <i>Journal of Diabetes Research</i> , 2020 , 2020, 5013142	3.9	2
207	Medication self-management: Considerations and decisions by older people living at home. <i>Research in Social and Administrative Pharmacy</i> , 2020 , 18, 2410-2410	2.9	2
206	Reported Adverse Drug Reactions in Children and Adolescents Treated with Antipsychotics. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2019 , 29, 124-132	2.9	17
205	Effects of a Proactive Interdisciplinary Self-Management (PRISMA) program on medication adherence in patients with type 2 diabetes in primary care: a randomized controlled trial. <i>Patient Preference and Adherence</i> , 2019 , 13, 749-759	2.4	7
204	Unbound Fraction of Clozapine Significantly Decreases with Elevated Plasma Concentrations of the Inflammatory Acute-Phase Protein Alpha-1-Acid Glycoprotein. <i>Clinical Pharmacokinetics</i> , 2019 , 58, 1069-	1075	4

203	Discontinuation Rates of Antidepressant Use by Dutch Soldiers. <i>Military Medicine</i> , 2019 , 184, 868-874	1.3	2
202	Associations Between Characteristics of Web-Based Diabetes News and Readers Rentiments: Observational Study in the Netherlands. <i>Journal of Medical Internet Research</i> , 2019 , 21, e14554	7.6	1
201	Effect of reimbursement restriction policy on the use of benzodiazepines in the Netherlands: an interrupted time series analysis. <i>BMJ Open</i> , 2019 , 9, e029148	3	3
200	Effects of the Proactive interdisciplinary self-management (PRISMA) program on self-reported and clinical outcomes in type 2 diabetes: a pragmatic randomized controlled trial. <i>BMC Endocrine Disorders</i> , 2019 , 19, 139	3.3	3
199	Compliance with the guidelines for laboratory monitoring of patients treated with lithium: A retrospective follow-up study among ambulatory patients in the Netherlands. <i>Bipolar Disorders</i> , 2019 , 21, 419-427	3.8	5
198	Severe Cognitive Impairment Associated With a High Free But Therapeutic Total Concentration of Valproic Acid Due to Hypoalbuminemia in an Older Patient With Bipolar Disorder. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 265-268	1.7	4
197	Monitoring of patients treated with lithium for bipolar disorder: an international survey. <i>International Journal of Bipolar Disorders</i> , 2018 , 6, 12	5.4	8
196	Diabetes and Glucose Dysregulation and Transition to Delirium in ICU Patients. <i>Critical Care Medicine</i> , 2018 , 46, 1444-1449	1.4	10
195	Association Between Urinary Tract Infections and Antipsychotic Drug Use in Older Adults. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 296-301	1.7	6
194	The impact of ethnic density on dispensing of antipsychotic and antidepressant medication among immigrants in the Netherlands. <i>Social Science and Medicine</i> , 2018 , 211, 87-94	5.1	3
193	Development of eHOME, a Mobile Instrument for Reporting, Monitoring, and Consulting Drug-Related Problems in Home Care: Human-Centered Design Study. <i>JMIR Human Factors</i> , 2018 , 5, e1	0 ^{2.5}	5
192	Monitoring of Metabolic, Cardiac, and Endocrine Indicators in Youth Treated With Antipsychotics as Reported by Health Care Professionals. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 489-493	1.7	5
191	Predicting rehospitalization in patients treated with antipsychotics: a prospective observational study. <i>Therapeutic Advances in Psychopharmacology</i> , 2018 , 8, 213-229	4.9	2
190	Predicting Unsuccessful Clozapine Treatment After First Use in Adult Patients With Psychotic Disorders. <i>Journal of Clinical Psychopharmacology</i> , 2018 , 38, 604-608	1.7	1
189	Glucose variability during delirium in diabetic and non-diabetic intensive care unit patients: A prospective cohort study. <i>PLoS ONE</i> , 2018 , 13, e0205637	3.7	8
188	Evaluation of clarity of presentation and applicability of monitoring instructions for patients using lithium in clinical practice guidelines for treatment of bipolar disorder. <i>Bipolar Disorders</i> , 2018 , 20, 708-	7 2 8	9
187	Medicinal Cannabis on Prescription in The Netherlands: Statistics for 2003-2016. <i>Cannabis and Cannabinoid Research</i> , 2018 , 3, 54-55	4.6	15
186	Antipsychotic drug use associated with urinary tract infections in older women. <i>Maturitas</i> , 2017 , 98, 46-	-5 6)	6

185	The Effect of Glycopyrrolate on Nocturnal Sialorrhea in Patients Using Clozapine: A Randomized, Crossover, Double-Blind, Placebo-Controlled Trial. <i>Journal of Clinical Psychopharmacology</i> , 2017 , 37, 155	5 -1 761	13
184	Dispensing of psychotropic medication among 400,000 immigrants in The Netherlands. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017 , 52, 963-977	4.5	4
183	Guidance by physicians and pharmacists during antidepressant therapy: patientsRneeds and suggestions for improvement. <i>BMC Psychiatry</i> , 2017 , 17, 388	4.2	4
182	Prevalence of Psychotropic Medication Use Among Dutch Military Personnel Between 2003 and 2012 and Its Comparison to the Dutch General Population. <i>Military Medicine</i> , 2017 , 182, e1584-e1588	1.3	3
181	Pharmacological Management of Aggressive Behavior in Psychiatric Patients 2017, 1-15		
180	Clozapine-induced hypersalivation: the association between quantification, perceived burden and treatment satisfaction reported by patients. <i>Therapeutic Advances in Psychopharmacology</i> , 2017 , 7, 209-	248	1
179	Somatic health care utilization by patients treated for substance use disorders. <i>Drug and Alcohol Dependence</i> , 2017 , 178, 277-284	4.9	7
178	Clozapine re-exposure after dilated cardiomyopathy. <i>BMJ Case Reports</i> , 2017 , 2017,	0.9	3
177	Patient monitoring: the hidden costs of treatment with antipsychotics. <i>British Journal of Clinical Pharmacology</i> , 2016 , 82, 570-2	3.8	1
176	Movement Disorders in Adults With Intellectual Disability and Behavioral Problems Associated With Use of Antipsychotics. <i>Journal of Clinical Psychopharmacology</i> , 2016 , 36, 308-13	1.7	10
175	Associations Between Personality Traits and Adherence to Antidepressants Assessed Through Self-Report, Electronic Monitoring, and Pharmacy Dispensing Data: A Pilot Study. <i>Journal of Clinical Psychopharmacology</i> , 2016 , 36, 465-71	1.7	5
174	Comparing episodes of antidepressants use with intermittent episodes of no use: A higher relative risk of suicide attempts but not of suicide at young age. <i>Journal of Psychopharmacology</i> , 2016 , 30, 1000-	4 .6	2
173	Structured Medication Review to Improve Pharmacotherapy in People with Intellectual Disability and Behavioural Problems. <i>Journal of Applied Research in Intellectual Disabilities</i> , 2016 , 29, 346-55	2.2	26
172	Adverse events and the relation with quality of life in adults with intellectual disability and challenging behaviour using psychotropic drugs. <i>Research in Developmental Disabilities</i> , 2016 , 49-50, 13-	217	36
171	Effects of Telephone Counseling Intervention by Pharmacists (TelCIP) on Medication Adherence; Results of a Cluster Randomized Trial. <i>Frontiers in Pharmacology</i> , 2016 , 7, 269	5.6	24
170	Proposal of Standardization to Assess Adherence With Medication Records: Methodology Matters. <i>Annals of Pharmacotherapy</i> , 2016 , 50, 360-8	2.9	41
169	Suicide Behavior Before and After the Start with Antidepressants: A High Persistent Risk in the First Month of Treatment Among the Young. <i>International Journal of Neuropsychopharmacology</i> , 2015 , 19,	5.8	10
168	PatientsRgeneral satisfaction with telephone counseling by pharmacists and effects on satisfaction with information and beliefs about medicines: Results from a cluster randomized trial. <i>Patient Education and Counseling</i> 2015 , 98, 797-804	3.1	12

(2013-2015)

167	Medication Discontinuation in Patients After Discharge From a Psychiatric Hospital. <i>Annals of Pharmacotherapy</i> , 2015 , 49, 1085-95	2.9	2	
166	Differences in ADHD medication usage patterns in children and adolescents from different cultural backgrounds in the Netherlands. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015 , 50, 1153-62	4.5	9	
165	Risk of Hospitalization for Hypoglycemia in Older Patients with Diabetes Using Antipsychotic Drugs. <i>American Journal of Geriatric Psychiatry</i> , 2015 , 23, 1144-53	6.5	4	
164	The relation between risk-taking behavior and alcohol use in young adults is different for men and women. <i>Drug and Alcohol Dependence</i> , 2015 , 155, 222-7	4.9	21	
163	Instructions for clinical and biomarker monitoring in the Summary of Product Characteristics (SmPC) for psychotropic drugs: Overview and applicability in clinical practice. <i>Journal of Psychopharmacology</i> , 2015 , 29, 1248-54	4.6	16	
162	Antidepressant use in pregnancy: knowledge transfer and translation of research findings. <i>Journal of Evaluation in Clinical Practice</i> , 2015 , 21, 579-83	2.5	2	
161	Association between concomitant psychiatric drug use, and patientsRbeliefs about and persistence with chronic cardiovascular medication. <i>International Journal of Clinical Practice</i> , 2015 , 69, 328-35	2.9	1	
160	Nonmedical Use of Attention-Deficit/Hyperactivity Disorder Medication Among Secondary School Students in The Netherlands. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2015 , 25, 649-52	2.9		
159	Discontinuation of somatic medication during psychiatric hospitalization. <i>Annals of Pharmacotherapy</i> , 2014 , 48, 1415-24	2.9	4	
158	Attitudes towards medication use in a general population of adolescents. <i>European Journal of Pediatrics</i> , 2014 , 173, 483-8	4.1	9	
157	Effects of a TELephone Counselling Intervention by Pharmacist (TelCIP) on medication adherence, patient beliefs and satisfaction with information for patients starting treatment: study protocol for a cluster randomized controlled trial. <i>BMC Health Services Research</i> , 2014 , 14, 219	2.9	10	
156	Combined HTR2C-LEP and HTR2C-LEPR genotypes as a determinant for obesity in patients without antipsychotic drugs. <i>Journal of Clinical Psychopharmacology</i> , 2014 , 34, 266-8	1.7		
155	Variability between nursing homes in prevalence of antipsychotic use in patients with dementia. <i>International Psychogeriatrics</i> , 2014 , 26, 363-71	3.4	30	
154	Association between ADHD drug use and injuries among children and adolescents. <i>European Child and Adolescent Psychiatry</i> , 2014 , 23, 95-102	5.5	30	
153	The association between CYP2D6 genotype and switching antipsychotic medication to clozapine. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1927-32	2.8	3	
152	The prevalence and incidence of medicinal cannabis on prescription in The Netherlands. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1575-80	2.8	59	
151	Lazy sunday afternoons: the negative impact of interruptions in patientsRdaily routine on adherence to oral antidiabetic medication. A multilevel analysis of electronic monitoring data. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1599-606	2.8	12	
150	Discontinuation of anticoagulant care during admission to a psychiatric hospital. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1025-9	2.8	3	

149	The association between prescription change frequency, chronic disease score and hospital admissions: a case control study. <i>BMC Pharmacology & Doctor Study</i> , 2013 , 14, 39	2.6	5
148	Fluorescence of neutrophil granulocytes as a biomarker for clozapine use. <i>European Neuropsychopharmacology</i> , 2013 , 23, 1408-13	1.2	4
147	The association between lifting an administrative restriction on antidepressant dispensing and treatment patterns in Iceland. <i>Health Policy</i> , 2013 , 111, 193-9	3.2	
146	Prevalence and characteristics of psychotropic drug use in institutionalized children and adolescents with mild intellectual disability. <i>Research in Developmental Disabilities</i> , 2013 , 34, 3159-67	2.7	23
145	Ineffectiveness and adverse events of nitrofurantoin in women with urinary tract infection and renal impairment in primary care. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1701-7	2.8	32
144	Prevalence of medication use for somatic disease in institutionalized psychiatric patients. <i>Pharmacopsychiatry</i> , 2013 , 46, 274-80	2	3
143	Does the use of an electronic reminder device with or without counseling improve adherence to lipid-lowering treatment? The results of a randomized controlled trial. <i>Frontiers in Pharmacology</i> , 2013 , 4, 69	5.6	19
142	Risk of acute coronary syndrome in elderly users of antipsychotic drugs: a nested case-control study. <i>Heart</i> , 2012 , 98, 1166-71	5.1	7
141	Differences in body mass index z-scores and weight status in a Dutch pediatric psychiatric population with and without use of second-generation antipsychotics. <i>Journal of Child and Adolescent Psychopharmacology</i> , 2012 , 22, 166-73	2.9	15
140	Association between HTR2C polymorphisms and obesity in patients without antipsychotic drugs. Journal of Clinical Psychopharmacology, 2012 , 32, 715-40	1.7	2
139	Influence of therapeutic complexity on medication adherence in the Netherlands. <i>Archives of Internal Medicine</i> , 2011 , 171, 864-5; author reply 865		3
138	Use of serotonergic drugs and the risk of bleeding. <i>Clinical Pharmacology and Therapeutics</i> , 2011 , 89, 89-96	6.1	41
137	Representation of people with intellectual disabilities in randomised controlled trials on antipsychotic treatment for behavioural problems. <i>Journal of Intellectual Disability Research</i> , 2011 , 55, 650-64	3.2	19
136	The decision to continue or discontinue treatment: experiences and beliefs of users of selective serotonin-reuptake inhibitors in the initial monthsa qualitative study. <i>Research in Social and Administrative Pharmacy</i> , 2011 , 7, 134-50	2.9	34
135	Pharmacogenetic testing to predict antipsychotic-induced weight gain: a systematic review. <i>Pharmacogenomics</i> , 2011 , 12, 1213-27	2.6	15
134	Polymorphisms of the LEP, LEPR and HTR2C gene: obesity and BMI change in patients using antipsychotic medication in a naturalistic setting. <i>Pharmacogenomics</i> , 2011 , 12, 919-23	2.6	30
133	Estimates of statin discontinuation rates are influenced by exposure and outcome definitions. <i>Annals of Pharmacotherapy</i> , 2011 , 45, 576-81	2.9	15
132	The association between patient-reported drug taking and gaps and overlaps in antidepressant drug dispensing. <i>Annals of Pharmacotherapy</i> , 2010 , 44, 1755-61	2.9	8

(2009-2010)

131	of Child and Adolescent Psychopharmacology, 2010 , 20, 55-61	2.9	35
130	Transitions from General Practitioner to Psychiatrist Care (or vice versa) during a first antidepressant treatment episode. <i>Pharmacopsychiatry</i> , 2010 , 43, 179-83	2	5
129	Low treatment adherence with antipsychotics is associated with relapse in psychotic disorders within six months after discharge. <i>Pharmacopsychiatry</i> , 2010 , 43, 221-4	2	12
128	Trials assessing parmacotherapeutic management of aggression in psychiatric patients: comparability with clinical practice. <i>Pharmacopsychiatry</i> , 2010 , 43, 205-9	2	5
127	Construction of drug treatment episodes from drug-dispensing histories is influenced by the gap length. <i>Journal of Clinical Epidemiology</i> , 2010 , 63, 422-7	5.7	69
126	Unpredictable bias when using the missing indicator method or complete case analysis for missing confounder values: an empirical example. <i>Journal of Clinical Epidemiology</i> , 2010 , 63, 728-36	5.7	100
125	Is aspirin useful in patients on lithium? A pharmacoepidemiological study related to bipolar disorder. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2010 , 82, 9-14	2.8	56
124	PatientsRperceptions and illness severity at start of antidepressant treatment in general practice. <i>International Journal of Pharmacy Practice</i> , 2010 , 18, 217-25	1.7	12
123	Antipsychotic drug use and the risk of venous thromboembolism in elderly patients. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 526-30	1.7	29
122	Association between LEP and LEPR gene polymorphisms and dyslipidemia in patients using atypical antipsychotic medication. <i>Psychiatric Genetics</i> , 2010 , 20, 311-6	2.9	19
121	Combined HTR2C-LEP genotype as a determinant of obesity in patients using antipsychotic medication. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 702-5	1.7	16
120	The association between serotonin 2C receptor polymorphisms and weight gain and eating behavior in patients using mirtazapine: a prospective follow-up study. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 207-9	1.7	4
119	Quantifying antibiotic use in paediatrics: a proposal for neonatal DDDs. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2010 , 29, 1301-3	5.3	26
118	Less discontinuation of ADHD drug use since the availability of long-acting ADHD medication in children, adolescents and adults under the age of 45 years in the Netherlands. <i>ADHD Attention Deficit and Hyperactivity Disorders</i> , 2010 , 2, 213-20	3.1	31
117	Seasonal patterns of initiating antidepressant therapy in general practice in the Netherlands during 2002-2007. <i>Journal of Affective Disorders</i> , 2010 , 122, 208-12	6.6	14
116	Risk of cerebrovascular events in elderly users of antipsychotics. <i>Journal of Psychopharmacology</i> , 2009 , 23, 909-14	4.6	65
115	Unexpected lower prevalence of depression in patients with diabetes: selection bias in a waiting room population. <i>Epidemiology</i> , 2009 , 20, 419-23	3.1	
114	Gardarsdottir et al. Respond to "Assessing Treatment Effects With Observational Analyses". <i>American Journal of Epidemiology</i> , 2009 , 170, 288-288	3.8	

113	Duration of antidepressant drug treatment and its influence on risk of relapse/recurrence: immortal and neglected time bias. <i>American Journal of Epidemiology</i> , 2009 , 170, 280-5	3.8	28
112	PatientsRperceptions of information received at the start of selective serotonin-reuptake inhibitor treatment: implications for community pharmacy. <i>Annals of Pharmacotherapy</i> , 2009 , 43, 642-9	2.9	23
111	Is new drug prescribing in primary care specialist induced?. BMC Health Services Research, 2009, 9, 6	2.9	24
110	Antidepressant use before and after initiation of diabetes mellitus treatment. <i>Diabetologia</i> , 2009 , 52, 425-32	10.3	24
109	The use of driving impairing medicines: a European survey. <i>European Journal of Clinical Pharmacology</i> , 2009 , 65, 1139-47	2.8	17
108	An algorithm to identify antidepressant users with a diagnosis of depression from prescription data. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 7-15	2.6	28
107	Association between lithium serum level, mood state, and patient-reported adverse drug reactions during long-term lithium treatment: a naturalistic follow-up study. <i>Bipolar Disorders</i> , 2009 , 11, 434-40	3.8	16
106	No difference in between-country variability in use of newly approved orphan and non- orphan medicinal productsa pilot study. <i>Orphanet Journal of Rare Diseases</i> , 2009 , 4, 27	4.2	13
105	Evaluation of available treatment guidelines for the management of lithium intoxication. <i>Therapeutic Drug Monitoring</i> , 2009 , 31, 247-60	3.2	5
104	Does the length of the first antidepressant treatment episode influence risk and time to a second episode?. <i>Journal of Clinical Psychopharmacology</i> , 2009 , 29, 69-72	1.7	18
103	Polymorphisms of the LEP- and LEPR gene and obesity in patients using antipsychotic medication. Journal of Clinical Psychopharmacology, 2009 , 29, 21-5	1.7	61
102	Influence of antidepressant use on glycemic control in patients with diabetes mellitus: an open-label comparative study. <i>Journal of Clinical Psychopharmacology</i> , 2009 , 29, 405-8	1.7	5
101	Initiation of antidepressant therapy: do patients follow the GPR prescription?. <i>British Journal of General Practice</i> , 2009 , 59, 81-7	1.6	85
100	The course of behavioral problems in elderly nursing home patients with dementia when treated with antipsychotics. <i>International Psychogeriatrics</i> , 2009 , 21, 931-40	3.4	23
99	Variable access to clopidogrel in a harmonized EU market. Value in Health, 2008, 11, 989-95	3.3	9
98	Clinical relevance of patients with epilepsy included in clinical trials. <i>Epilepsia</i> , 2008 , 49, 1479-80	6.4	13
97	Furthering patient adherence: a position paper of the international expert forum on patient adherence based on an internet forum discussion. <i>BMC Health Services Research</i> , 2008 , 8, 47	2.9	58
96	Changes in outpatient lithium treatment in the Netherlands during 1996-2005. <i>Journal of Affective Disorders</i> , 2008 , 111, 94-9	6.6	15

(2007-2008)

95	NSAID and COX-2 inhibitor users: a cohort study using pharmacy dispensing data in The Netherlands. <i>Drug Safety</i> , 2008 , 31, 143-58	5.1	27
94	Genetic variation in the renin-angiotensin system modifies the beneficial effects of ACE inhibitors on the risk of diabetes mellitus among hypertensives. <i>Journal of Human Hypertension</i> , 2008 , 22, 774-80	2.6	19
93	Impact analysis of the discontinuation of reimbursement: the case of oral contraceptives. <i>Contraception</i> , 2008 , 78, 399-404	2.5	7
92	Characteristics and reasons associated with nonacceptance of selective serotonin-reuptake inhibitor treatment. <i>Annals of Pharmacotherapy</i> , 2008 , 42, 218-25	2.9	32
91	Use of antidepressants in pregnancy does not increase risk of postpartum haemorrhage. <i>Evidence-Based Mental Health</i> , 2008 , 11, 125	11.1	
90	Association between hospitalization and discontinuity of medication therapy used in the community setting in the Netherlands. <i>Annals of Pharmacotherapy</i> , 2008 , 42, 933-9	2.9	20
89	The association between concomitant use of serotonergic antidepressants and lithium-induced polyuria. A multicenter medical chart review study. <i>Pharmacopsychiatry</i> , 2008 , 41, 129-33	2	4
88	The association between antidepressant use and disturbances in glucose homeostasis: evidence from spontaneous reports. <i>European Journal of Clinical Pharmacology</i> , 2008 , 64, 531-8	2.8	72
87	The association between antidepressant use and hypoglycaemia in diabetic patients: a nested case-control study. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 336-44	2.6	21
86	Influence of antidepressants on glycaemic control in patients with diabetes mellitus. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 577-86	2.6	23
85	Prescriber adoption of newly approved selective COX-2 inhibitors. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 1168-74	2.6	3
84	Lithium: farmacologie in de psychiatrie. <i>Neuropraxis</i> , 2008 , 12, 147-153	О	1
83	Visualizing Pharmacological Activities of Antidepressants: A Novel Approach. <i>The Open Pharmacology Journal</i> , 2008 , 2, 54-62	0.3	21
82	Prevalence of patients using drugs metabolized by cytochrome P450 2D6 in different populations: a cross-sectional study. <i>Annals of Pharmacotherapy</i> , 2007 , 41, 408-13	2.9	28
81	Pharmacogenetics of glucose-lowering drug treatment: a systematic review. <i>Molecular Diagnosis and Therapy</i> , 2007 , 11, 291-302	4.5	20
80	Antidepressant use and off-label prescribing in children and adolescents in Dutch general practice (2001-2005). <i>Pharmacoepidemiology and Drug Safety</i> , 2007 , 16, 1054-62	2.6	42
79	The impact of environmental temperature on lithium serum levels. <i>Bipolar Disorders</i> , 2007 , 9, 603-8	3.8	12
78	Determinants of the range of drugs prescribed in general practice: a cross-sectional analysis. <i>BMC Health Services Research</i> , 2007 , 7, 132	2.9	21

77	Patient risk profiles and practice variation in nonadherence to antidepressants, antihypertensives and oral hypoglycemics. <i>BMC Health Services Research</i> , 2007 , 7, 51	2.9	53
76	Patient adherence to medical treatment: a review of reviews. <i>BMC Health Services Research</i> , 2007 , 7, 55	2.9	362
75	Indications for antidepressant drug prescribing in general practice in the Netherlands. <i>Journal of Affective Disorders</i> , 2007 , 98, 109-15	6.6	126
74	Urine osmolality, cyclic AMP and aquaporin-2 in urine of patients under lithium treatment in response to water loading followed by vasopressin administration. <i>European Journal of Pharmacology</i> , 2007 , 566, 50-7	5.3	9
73	Evaluation of patientsRexperiences with antidepressants reported by means of a medicine reporting system. <i>European Journal of Clinical Pharmacology</i> , 2007 , 63, 1193-9	2.8	30
72	The association between exposure to COX-2 inhibitors and schizophrenia deterioration. A nested case-control study. <i>Pharmacopsychiatry</i> , 2007 , 40, 111-5	2	13
71	Aggression of psychiatric patients associated with the use of as-needed medication. <i>Pharmacopsychiatry</i> , 2007 , 40, 25-9	2	14
70	The effect of pharmacotherapy audit meetings on early new drug prescribing by general practitioners. <i>Annals of Pharmacotherapy</i> , 2007 , 41, 319-24	2.9	19
69	Rates and determinants of reinitiating antihypertensive therapy after prolonged stoppage: a population-based study. <i>Journal of Hypertension</i> , 2007 , 25, 689-97	1.9	11
68	No increased incidence of diabetes in antidepressant users. <i>International Clinical Psychopharmacology</i> , 2007 , 22, 382-6	2.2	42
67	Depressive symptoms in subjects with diagnosed and undiagnosed type 2 diabetes. <i>Psychosomatic Medicine</i> , 2007 , 69, 300-5	3.7	86
66	Lithium use and the risk of fractures. <i>Bone</i> , 2007 , 40, 1252-8	4.7	53
65	Determinants of compliance with methylphenidate therapy in children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2006 , 95, 1674-6	3.1	4
64	Linking community pharmacy dispensing data to prescribing data of general practitioners. <i>BMC Medical Informatics and Decision Making</i> , 2006 , 6, 18	3.6	18
63	Generic substitution of antihypertensive drugs: does it affect adherence?. <i>Annals of Pharmacotherapy</i> , 2006 , 40, 15-20	2.9	68
62	Which pharmacists contribute to high-level pharmacotherapy audit meetings with general practitioners?. <i>Annals of Pharmacotherapy</i> , 2006 , 40, 1640-6	2.9	7
61	Early identification of long-term poor adherence in ambulatory patients. <i>Annals of Pharmacotherapy</i> , 2006 , 40, 2277-8	2.9	2
60	A comparison of two multiple-characteristic decision-making models for the comparison of antihypertensive drug classes: Simple Additive Weighting (SAW) and Technique for Order Preference by Similarity to an Ideal Solution (TOPSIS). <i>American Journal of Cardiovascular Drugs</i> ,	4	7

(2005-2006)

59	Refill persistence with chronic medication assessed from a pharmacy database was influenced by method of calculation. <i>Journal of Clinical Epidemiology</i> , 2006 , 59, 11-7	5.7	42
58	Potential bias in pharmacoepidemiological studies due to the length of the drug free period: a study on antidepressant drug use in adults in the Netherlands. <i>Pharmacoepidemiology and Drug Safety</i> , 2006 , 15, 338-43	2.6	21
57	Initial non-compliance with antihypertensive monotherapy is followed by complete discontinuation of antihypertensive therapy. <i>Pharmacoepidemiology and Drug Safety</i> , 2006 , 15, 587-93	2.6	16
56	Between-country variation in the utilization of antihypertensive agents: Guidelines and clinical practice. <i>Journal of Human Hypertension</i> , 2006 , 20, 917-22	2.6	18
55	Haloperidol dose when used as active comparator in randomized controlled trials with atypical antipsychotics in schizophrenia: comparison with officially recommended doses. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 897-903	4.6	36
54	Pharmacotherapy for the treatment of aggressive behavior in general adult psychiatry: A systematic review. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 1013-24	4.6	60
53	Effectiveness of interventions by community pharmacists to improve patient adherence to chronic medication: a systematic review. <i>Annals of Pharmacotherapy</i> , 2005 , 39, 319-28	2.9	93
52	Risk of hip/femur fractures in patients using antipsychotics. <i>Bone</i> , 2005 , 37, 864-70	4.7	58
51	Rate and determinants of 10-year persistence with antihypertensive drugs. <i>Journal of Hypertension</i> , 2005 , 23, 2101-7	1.9	141
50	Impaired glucose homeostasis after imipramine intake in a diabetic patient. <i>Journal of Clinical Psychopharmacology</i> , 2005 , 25, 621-3	1.7	5
49	Drug-drug interactions as a determinant of elevated lithium serum levels in daily clinical practice. <i>Bipolar Disorders</i> , 2005 , 7, 274-80	3.8	24
48	Patient characteristics associated with prescribing of a newly introduced drug: the case of rofecoxib. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 157-9	2.8	8
47	Changes in the defined daily dose; CYP2D6/CYP3A metabolism as an indicator for dose-setting problems. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 243-6	2.8	3
46	The impact of hospitalisation on the initiation and long-term use of benzodiazepines. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 291-5	2.8	16
45	Discontinuation symptoms in users of selective serotonin reuptake inhibitors in clinical practice: tapering versus abrupt discontinuation. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 303-7	2.8	43
44	Methylphenidate: use in daily practice. International Journal of Clinical Pharmacy, 2005, 27, 197-201		10
43	The trade-off between cardiovascular and gastrointestinal effects of rofecoxib. <i>Pharmacoepidemiology and Drug Safety</i> , 2005 , 14, 437-41	2.6	8
42	A pharmacy-based coaching program to improve adherence to antidepressant treatment among primary care patients. <i>Psychiatric Services</i> , 2005 , 56, 487-9	3.3	51

41	Reasons for switching between antipsychotics in daily clinical practice. <i>Pharmacopsychiatry</i> , 2005 , 38, 122-4	2	30
40	Seclusion and the use of antipsychotics in hospitalized psychiatric patients. <i>Psychology, Crime and Law</i> , 2005 , 11, 489-495	1.4	15
39	Psychoactive drug prescribing by Dutch child and adolescent psychiatrists. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2005 , 94, 1484-1487	3.1	14
38	Need for medicine-based evidence in pharmacotherapy. <i>British Journal of Psychiatry</i> , 2004 , 184, 452	5.4	9
37	Incidence and determinants of long-term use of antidepressants. <i>European Journal of Clinical Pharmacology</i> , 2004 , 60, 57-61	2.8	84
36	Increased psychotropic drug consumption by children in the Netherlands during 1995-2001 is caused by increased use of methylphenidate by boys. <i>European Journal of Clinical Pharmacology</i> , 2004 , 60, 377-9	2.8	35
35	Correlated measures in longitudinal analysis of daily drug use patterns in a general intensive care unit. <i>Pharmacoepidemiology and Drug Safety</i> , 2004 , 13, 395-8	2.6	
34	Less medication switching after initial start with atypical antipsychotics. <i>European Neuropsychopharmacology</i> , 2004 , 14, 1-5	1.2	6
33	Should rosuvastatin be withdrawn from the market?. <i>Lancet, The</i> , 2004 , 364, 1577; author reply 1578-9	40	12
32	The association between compliance with antihypertensive drugs and modification of antihypertensive drug regimen. <i>Journal of Hypertension</i> , 2004 , 22, 1831-7	1.9	36
31	Association of risk of abnormal bleeding with degree of serotonin reuptake inhibition by antidepressants. <i>Archives of Internal Medicine</i> , 2004 , 164, 2367-70		173
30	Patterns of pharmacotherapy in patients hospitalised for congestive heart failure. <i>European Journal of Heart Failure</i> , 2003 , 5, 195-200	12.3	21
29	Short-acting parenteral antipsychotics drive choice for classical versus atypical agents. <i>European Journal of Clinical Pharmacology</i> , 2003 , 58, 757-60	2.8	6
28	Refill compliance in type 2 diabetes mellitus: a predictor of switching to insulin therapy?. <i>Pharmacoepidemiology and Drug Safety</i> , 2003 , 12, 121-7	2.6	33
27	Effects of NSAIDs on the incidence of hospitalisations for renal dysfunction in users of ACE inhibitors. <i>Drug Safety</i> , 2003 , 26, 983-9	5.1	26
26	Effect of a pharmacist-led intervention on diuretic compliance in heart failure patients: a randomized controlled study. <i>Journal of Cardiac Failure</i> , 2003 , 9, 404-11	3.3	148
25	Impact of coaching by community pharmacists on drug attitude of depressive primary care patients and acceptability to patients; a randomized controlled trial. <i>European Neuropsychopharmacology</i> , 2003 , 13, 1-9	1.2	64
24	Channelling new antidepressants to problem patients may be factor in fatal toxicity. <i>BMJ, The</i> , 2003 , 326, 600	5.9	5

23	Amiodarone-induced thyroid dysfunction associated with cumulative dose. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 601-6	2.6	28
22	Changes in prescribed drug doses after market introduction. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 447-53	2.6	49
21	Adverse events in users of sertraline: results from an observational study in psychiatric practice in The Netherlands. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 655-62	2.6	23
20	Characteristics of diarrhoea in 10,008 users of lansoprazole in daily practice: which co-factors contribute?. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 703-8	2.6	14
19	Psychotropic drug use in intellectually disabled group-home residents with behavioural problems. <i>Pharmacopsychiatry</i> , 2002 , 35, 19-23	2	45
18	Association of nonsteroidal anti-inflammatory drugs with first occurrence of heart failure and with relapsing heart failure: the Rotterdam Study. <i>Archives of Internal Medicine</i> , 2002 , 162, 265-70		146
17	Determinants of headache in lansoprazole users in The Netherlands: results from a nested case-control study. <i>Drug Safety</i> , 2002 , 25, 287-95	5.1	6
16	Is sexual dysfunctioning a reason for discontinuation in users of selective serotonin reuptake inhibitors?. <i>Journal of Clinical Psychopharmacology</i> , 2002 , 22, 96-8	1.7	6
15	Factors associated with non-response in proton pump inhibitor users: a study of lansoprazole therapy. <i>International Journal of Clinical Pharmacy</i> , 2001 , 23, 107-10		1
14	Long-term therapy with spironolactone. International Journal of Clinical Pharmacy, 2001, 23, 132-4		3
13	Prescribing patterns in patients using new antidepressants. <i>British Journal of Clinical Pharmacology</i> , 2001 , 51, 181-3	3.8	10
12	Determinants of multiple psychotropic drug use in patients with mild intellectual disabilities or borderline intellectual functioning and psychiatric or behavioral disorders. <i>General Hospital Psychiatry</i> , 2001 , 23, 345-9	5.6	31
11	Spontaneous lapses in dosing during chronic treatment with selective serotonin reuptake inhibitors. <i>British Journal of Psychiatry</i> , 2001 , 179, 519-22	5.4	22
10	Use of sympathomimetic drugs leads to increased risk of hospitalization for arrhythmias in patients with congestive heart failure. <i>Archives of Internal Medicine</i> , 2000 , 160, 2477-80		22
9	Safety review in 10,008 users of lansoprazole in daily practice. <i>Pharmacoepidemiology and Drug Safety</i> , 2000 , 9, 383-91	2.6	4
8	Detecting drug-drug interactions using a database for spontaneous adverse drug reactions: an example with diuretics and non-steroidal anti-inflammatory drugs. <i>European Journal of Clinical Pharmacology</i> , 2000 , 56, 733-8	2.8	84
7	Continued use of the proton-pump inhibitor lansoprazole following Helicobacter pylori eradication. <i>European Journal of Clinical Pharmacology</i> , 2000 , 55, 773-6	2.8	1
6	Benzodiazepine pathways in the chronically ill. <i>Pharmacoepidemiology and Drug Safety</i> , 1999 , 8, 325-30	2.6	7

5	Determinants of psychotropic drug usage in a general intensive care unit. <i>General Hospital Psychiatry</i> , 1998 , 20, 371-6	5.6	4
4	Application of the ATC/DDD methodology to monitor antibiotic drug use. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 1998 , 17, 20-4	5.3	77
3	NSAIDs associated with increased risk of congestive heart failure in elderly patients taking diuretics. <i>Archives of Internal Medicine</i> , 1998 , 158, 1108-12		199
2	A prospective follow-up study of 5669 users of lansoprazole in daily practice. <i>Alimentary Pharmacology and Therapeutics</i> , 1997 , 11, 887-97	6.1	41
1	Information on drug use in the elderly: a comparison of pharmacy, general-practitioner and patient data. <i>International Journal of Clinical Pharmacy</i> , 1995 , 17, 20-4		39