

Kim Khaw

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5243574/kim-khaw-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

30
papers

1,513
citations

15
h-index

30
g-index

30
ext. papers

1,761
ext. citations

3.2
avg, IF

4.24
L-index

#	Paper	IF	Citations
30	Performance of a closed-loop feedback computer-controlled infusion system for maintaining blood pressure during spinal anaesthesia for caesarean section: a randomized controlled comparison of norepinephrine versus phenylephrine. <i>Journal of Clinical Monitoring and Computing</i> , 2017 , 31, 617-623	2	31
29	Dose-Response Curves for Intrathecal Bupivacaine, Levobupivacaine, and Ropivacaine Given for Labor Analgesia in Nulliparous Women. <i>Regional Anesthesia and Pain Medicine</i> , 2017 , 42, 788-792	3.4	5
28	The effect of head up tilting on bioreactance cardiac output and stroke volume readings using suprasternal transcutaneous Doppler as a control in healthy young adults. <i>Journal of Clinical Monitoring and Computing</i> , 2016 , 30, 519-26	2	13
27	Randomized double-blinded comparison of norepinephrine and phenylephrine for maintenance of blood pressure during spinal anesthesia for cesarean delivery. <i>Anesthesiology</i> , 2015 , 122, 736-45	4.3	138
26	Randomized Double-blinded Comparison of Norepinephrine and Phenylephrine for Maintenance of Blood Pressure During Spinal Anesthesia for Cesarean Delivery. <i>Survey of Anesthesiology</i> , 2015 , 59, 267-268		
25	Synergistic interaction between fentanyl and bupivacaine given intrathecally for labor analgesia. <i>Anesthesiology</i> , 2014 , 120, 1126-36	4.3	36
24	The force applied to successfully turn a foetus during reattempts of external cephalic version is substantially reduced when performed under spinal analgesia. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2012 , 25, 719-22	2	15
23	Determination and comparison of graded dose-response curves for epidural bupivacaine and ropivacaine for analgesia in laboring nulliparous women. <i>Anesthesiology</i> , 2010 , 113, 445-53	4.3	23
22	Ephedrine and Phenylephrine Use during Cesarean Delivery. <i>Anesthesiology</i> , 2010 , 112, 1290-1294	4.3	1
21	Placental transfer and fetal metabolic effects of phenylephrine and ephedrine during spinal anesthesia for cesarean delivery. <i>Anesthesiology</i> , 2009 , 111, 506-12	4.3	189
20	Low-dose spinal anesthesia with low-dose phenylephrine infusions for cesarean delivery: better but not necessarily best. <i>Anesthesiology</i> , 2009 , 111, 210-1; author reply 213-6	4.3	1
19	A randomized double-blinded comparison of phenylephrine and ephedrine infusion combinations to maintain blood pressure during spinal anesthesia for cesarean delivery: the effects on fetal acid-base status and hemodynamic control. <i>Anesthesia and Analgesia</i> , 2008 , 107, 1295-302	3.9	148
18	A prospective comparison of vasopressor requirement and hemodynamic changes during spinal anesthesia for cesarean delivery in patients with multiple gestation versus singleton pregnancy. <i>Anesthesia and Analgesia</i> , 2007 , 104, 407-11	3.9	11
17	HYPOTENSION DURING SPINAL ANAESTHESIA FOR CAESAREAN SECTION: IMPLICATIONS, DETECTION PREVENTION AND TREATMENT. <i>Fetal and Maternal Medicine Review</i> , 2006 , 17, 157-183		15
16	Maternal and neonatal effects of remifentanyl at induction of general anesthesia for cesarean delivery: a randomized, double-blind, controlled trial. <i>Anesthesiology</i> , 2006 , 104, 14-20	4.3	127
15	Vasopressors in obstetrics: what should we be using?. <i>Current Opinion in Anaesthesiology</i> , 2006 , 19, 238-43		86
14	Prevention of hypotension during spinal anesthesia for cesarean delivery: an effective technique using combination phenylephrine infusion and crystalloid cohydration. <i>Anesthesiology</i> , 2005 , 103, 744-50	4.3	237

13	Prophylactic phenylephrine infusion for preventing hypotension during spinal anesthesia for cesarean delivery. <i>Anesthesia and Analgesia</i> , 2004 , 98, 815-21, table of contents	3.9	110
12	Spinal ropivacaine for cesarean delivery: a comparison of hyperbaric and plain solutions. <i>Anesthesia and Analgesia</i> , 2002 , 94, 680-5; table of contents	3.9	76
11	High failure rate of small doses of ropivacaine when used alone for spinal anesthesia for cesarean delivery. <i>Regional Anesthesia and Pain Medicine</i> , 2002 , 27, 533;author reply 534-5	3.4	
10	High failure rate of small doses of ropivacaine when used alone for spinal anesthesia for cesarean delivery. <i>Regional Anesthesia and Pain Medicine</i> , 2002 , 27, 533-533	3.4	
9	The limitations of ropivacaine with epinephrine as an epidural test dose in parturients. <i>Anesthesia and Analgesia</i> , 2001 , 92, 1529-31	3.9	9
8	Metaraminol infusion for maintenance of arterial blood pressure during spinal anesthesia for cesarean delivery: the effect of a crystalloid bolus. <i>Anesthesia and Analgesia</i> , 2001 , 93, 703-8	3.9	31
7	A dose-response study of prophylactic intravenous ephedrine for the prevention of hypotension during spinal anesthesia for cesarean delivery. <i>Anesthesia and Analgesia</i> , 2000 , 90, 1390-5	3.9	138
6	Perioperative care in left ventricular volume reduction. <i>Journal of Cardiac Surgery</i> , 1999 , 14, 136-40	1.3	0
5	Ministernotomy for aortic valve replacement in a patient with osteogenesis imperfecta. <i>Annals of Thoracic Surgery</i> , 1999 , 67, 1171-3	2.7	10
4	Routine intraoperative angiography improves the early patency of coronary grafts performed on the beating heart. <i>Chest</i> , 1999 , 115, 987-90	5.3	25
3	Upper T mini-sternotomy for aortic valve operations. <i>Chest</i> , 1998 , 114, 291-4	5.3	18
2	Postoperative analgesic requirement after cesarean section: a comparison of anesthetic induction with ketamine or thiopental. <i>Anesthesia and Analgesia</i> , 1997 , 85, 1294-8	3.9	10
1	Patient-controlled epidural analgesia after caesarean section using meperidine. <i>Canadian Journal of Anaesthesia</i> , 1997 , 44, 702-6	3	10