peng jing Zhu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5249601/publications.pdf Version: 2024-02-01

| | | 687363 | 713466 |
|----------|----------------|--------------|----------------|
| 22 | 1,111 | 13 | 21 |
| papers | citations | h-index | g-index |
| | | | |
| | | | |
| | | | |
| 22 | 22 | 22 | 946 |
| all docs | docs citations | times ranked | citing authors |
| | | | |

PENC LINC 7HU

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Chiral derivatives of covalent organic framework TpBD (NH ₂) ₂ used as stationary phases in gas chromatography. Chirality, 2022, 34, 462-472. | 2.6 | 7 |
| 2 | Enantioselective resolutions by highâ€performance liquid choromatography using chiral inorganic mesoporous silica. Separation Science Plus, 2021, 4, 77-85. | 0.6 | 3 |
| 3 | A [3Â+Â6] prismatic homochiral organic cage used as stationary phase for gas chromatography. Microchemical Journal, 2021, 170, 106650. | 4.5 | 7 |
| 4 | Chiral metalâ€organic cages used as stationary phase for enantioseparations in capillary electrochromatography. Electrophoresis, 2020, 41, 104-111. | 2.4 | 19 |
| 5 | Recent advances of application of porous molecular cages for enantioselective recognition and separation. Journal of Separation Science, 2020, 43, 134-149. | 2.5 | 55 |
| 6 | Gas chromatographic separation of enantiomers on novel chiral stationary phases. TrAC - Trends in Analytical Chemistry, 2020, 124, 115808. | 11.4 | 72 |
| 7 | Enantiomeric Separation on a Homochiral Porous Organic Cage-Based Chiral Stationary Phase by Gas Chromatography. Chromatographia, 2020, 83, 703-713. | 1.3 | 11 |
| 8 | Recent development trends for chiral stationary phases based on chitosan derivatives, cyclofructan derivatives and chiral porous materials in high performance liquid chromatography. Journal of Separation Science, 2019, 42, 6-20. | 2.5 | 63 |
| 9 | A novel chiral inorganic mesoporous silica used as a stationary phase in GC. Chirality, 2019, 31, 1053-1059. | 2.6 | 17 |
| 10 | Chiral Inorganic mesoporous materials used as the stationary phase in GC. Separation Science Plus, 2019, 2, 432-439. | 0.6 | 9 |
| 11 | An Enantioselective Potentiometric Sensor for 2-Amino-1-Butanol Based on Chiral Porous Organic Cage CC3-R. Molecules, 2019, 24, 420. | 3.8 | 9 |
| 12 | Enantioseparations by Gas Chromatography Using Porous Organic Cages as Stationary Phase. Methods in Molecular Biology, 2019, 1985, 45-55. | 0.9 | 2 |
| 13 | A highly ordered chiral inorganic mesoporous material used as stationary phase for high-resolution gas chromatographic separations. Journal of Chromatography A, 2018, 1557, 99-106. | 3.7 | 22 |
| 14 | Homochiral porous organic cage used as stationary phase for open tubular capillary electrochromatography. Analytica Chimica Acta, 2018, 999, 169-175. | 5.4 | 64 |
| 15 | Homochiral Metal–Organic Cage for Gas Chromatographic Separations. Analytical Chemistry, 2018, 90, 9182-9188. | 6.5 | 59 |
| 16 | A chiral porous organic cage for molecular recognition using gas chromatography. Analytica Chimica Acta, 2016, 903, 156-163. | 5.4 | 60 |
| 17 | Homochiral Porous Organic Cage with High Selectivity for the Separation of Racemates in Gas Chromatography. Analytical Chemistry, 2015, 87, 7817-7824. | 6.5 | 121 |
| 18 | Highly selective separation of enantiomers using a chiral porous organic cage. Journal of Chromatography A, 2015, 1426, 174-182. | 3.7 | 60 |

PENG JING ZHU

| # | Article | IF | CITATIONS |
|----|--|-------------|-----------|
| 19 | Chiral metal–organic framework used as stationary phases for capillary electrochromatography. Analytica Chimica Acta, 2014, 830, 49-55. | 5.4 | 85 |
| 20 | Novel Inorganic Mesoporous Material with Chiral Nematic Structure Derived from Nanocrystalline Cellulose for High-Resolution Gas Chromatographic Separations. Analytical Chemistry, 2014, 86, 9595-9602. | 6.5 | 72 |
| 21 | Chiral Metal–Organic Frameworks for High-Resolution Gas Chromatographic Separations. Journal of the American Chemical Society, 2011, 133, 11892-11895. | 13.7 | 293 |
| 22 | Separation of enantiomers by openâ€tubular capillary electrochromatography using (R)â€1,1′â€biâ€2â€naph derivatives as chiral stationary phases. Separation Science Plus, 0, , . | thol 0.6 | 1 |