Jeffrey Lipman

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/5251153/jeffrey-lipman-publications-by-year.pdf

Version: 2024-04-03

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

24,496 76 137 539 h-index g-index citations papers 6.2 605 29,295 7.04 avg, IF L-index ext. citations ext. papers

| # | Paper | IF | Citations |
|-----|--|------|-----------|
| 539 | Pharmacodynamics of ceftriaxone for the treatment of methicillin-susceptible Staphylococcus aureus: Is it a viable treatment option?. <i>International Journal of Antimicrobial Agents</i> , 2022 , 106537 | 14.3 | O |
| 538 | Innovation in microsampling for therapeutic drug monitoring of gentamicin in neonates: a proof-of-concept study <i>International Journal of Antimicrobial Agents</i> , 2022 , 59, 106513 | 14.3 | |
| 537 | Evaluation of low-volume plasma sampling for the analysis of meropenem in clinical samples <i>Analytical and Bioanalytical Chemistry</i> , 2022 , 1 | 4.4 | O |
| 536 | Epidemiology and Age-Related Mortality in Critically Ill Patients With Intra-Abdominal Infection or Sepsis: An International Cohort Study <i>International Journal of Antimicrobial Agents</i> , 2022 , 106591 | 14.3 | 0 |
| 535 | Multicenter Population Pharmacokinetic Study of Unbound Ceftriaxone in Critically Ill Patients <i>Antimicrobial Agents and Chemotherapy</i> , 2022 , e0218921 | 5.9 | 1 |
| 534 | A validated LC-MS/MS method for the simultaneous quantification of the novel combination antibiotic, ceftolozane-tazobactam, in plasma (total and unbound), CSF, urine and renal replacement therapy effluent: application to pilot pharmacokinetic studies. <i>Clinical Chemistry and Laboratory Medicine</i> , 2021, 59, 921-933 | 5.9 | 2 |
| 533 | A personalised approach to antibiotic pharmacokinetics and pharmacodynamics in critically ill patients. <i>Anaesthesia, Critical Care & amp; Pain Medicine</i> , 2021 , 40, 100970 | 3 | 5 |
| 532 | Early Sequential Microcirculation Assessment In Shocked Patients as a Predictor of Outcome: A Prospective Observational Cohort Study. <i>Shock</i> , 2021 , 55, 581-586 | 3.4 | 4 |
| 531 | Re-expansion pulmonary oedema with Takotsubo cardiomyopathy: a rare complication of giant hepatic cyst drainage. <i>ANZ Journal of Surgery</i> , 2021 , 91, 2524-2527 | 1 | O |
| 530 | Genomic surveillance, characterization and intervention of a polymicrobial multidrug-resistant outbreak in critical care. <i>Microbial Genomics</i> , 2021 , 7, | 4.4 | 6 |
| 529 | Evaluation of Meropenem-Ciprofloxacin Combination Dosage Regimens for the Pharmacokinetics of Critically Ill Patients With Augmented Renal Clearance. <i>Clinical Pharmacology and Therapeutics</i> , 2021 , 109, 1104-1115 | 6.1 | 5 |
| 528 | An international survey on aminoglycoside practices in critically ill patients: the AMINO III study. <i>Annals of Intensive Care</i> , 2021 , 11, 49 | 8.9 | 5 |
| 527 | Impact of the Epithelial Lining Fluid Milieu on Amikacin Pharmacodynamics Against Pseudomonas aeruginosa. <i>Drugs in R and D</i> , 2021 , 21, 203-215 | 3.4 | 1 |
| 526 | Lorazepam challenge test for catatonia. EMA - Emergency Medicine Australasia, 2021, 33, 763-764 | 1.5 | 1 |
| 525 | Development and validation of a UHPLC-MS/MS method to measure cefotaxime and metabolite desacetylcefotaxime in blood plasma: a pilot study suitable for capillary microsampling in critically ill children. <i>Analytical and Bioanalytical Chemistry</i> , 2021 , 413, 4483-4491 | 4.4 | O |
| 524 | Influence of the Obesity Phenotype on the Adequacy of Antibiotic Prophylaxis with Cefoxitin for Obese Patients Undergoing Bariatric Surgery: Lessons Learnt and Future Considerations. <i>European Journal of Drug Metabolism and Pharmacokinetics</i> , 2021 , 46, 479-485 | 2.7 | 1 |
| 523 | A Multicenter Observational Study Evaluating Outcomes Associated With Antibiotic Combination Versus Monotherapy in Patients With Septic Shock 2021 , 3, e0383 | | |

| 522 | Antimicrobial Lessons From a Large Observational Cohort on Intra-abdominal Infections in Intensive Care Units. <i>Drugs</i> , 2021 , 81, 1065-1078 | 12.1 | 3 |
|-----|---|------|----|
| 521 | Microsampling to support pharmacokinetic clinical studies in pediatrics. <i>Pediatric Research</i> , 2021 , | 3.2 | 1 |
| 520 | Prospective Cohort Study of Micafungin Population Pharmacokinetic Analysis in Plasma and Peritoneal Fluid in Septic Patients with Intra-abdominal Infections. <i>Antimicrobial Agents and Chemotherapy</i> , 2021 , 65, e0230720 | 5.9 | 0 |
| 519 | Pharmacodynamics of once- versus twice-daily dosing of nebulized amikacin in an in vitro Hollow-Fiber Infection Model against 3 clinical isolates of Pseudomonas aeruginosa. <i>Diagnostic Microbiology and Infectious Disease</i> , 2021 , 100, 115329 | 2.9 | 1 |
| 518 | Optimising Treatment Outcomes for Children and Adults Through Rapid Genome Sequencing of Sepsis Pathogens. A Study Protocol for a Prospective, Multi-Centre Trial (DIRECT). <i>Frontiers in Cellular and Infection Microbiology</i> , 2021 , 11, 667680 | 5.9 | 1 |
| 517 | The Effect of Renal Replacement Therapy and Antibiotic Dose on Antibiotic Concentrations in Critically Ill Patients: Data From the Multinational Sampling Antibiotics in Renal Replacement Therapy Study. <i>Clinical Infectious Diseases</i> , 2021 , 72, 1369-1378 | 11.6 | 37 |
| 516 | Ceftriaxone dosing in patients admitted from the emergency department with sepsis. <i>European Journal of Clinical Pharmacology</i> , 2021 , 77, 207-214 | 2.8 | 2 |
| 515 | Population Pharmacokinetics of Levetiracetam in Patients with Traumatic Brain Injury and Subarachnoid Hemorrhage Exhibiting Augmented Renal Clearance. <i>Clinical Pharmacokinetics</i> , 2021 , 60, 655-664 | 6.2 | 6 |
| 514 | A narrative review on antimicrobial therapy in septic shock: updates and controversies. <i>Current Opinion in Anaesthesiology</i> , 2021 , 34, 92-98 | 2.9 | 1 |
| 513 | Ventilation practices in burn patients-an international prospective observational cohort study <i>Burns and Trauma</i> , 2021 , 9, tkab034 | 5.3 | O |
| 512 | A Loading Micafungin Dose in Critically Ill Patients Undergoing Continuous Venovenous Hemofiltration or Continuous Venovenous Hemodiafiltration: A Population Pharmacokinetic Analysis. <i>Therapeutic Drug Monitoring</i> , 2021 , 43, 747-755 | 3.2 | O |
| 511 | Initial antimicrobial management of sepsis. <i>Critical Care</i> , 2021 , 25, 307 | 10.8 | 14 |
| 510 | Meropenem Versus Piperacillin-Tazobactam for Definitive Treatment of Bloodstream Infections Caused by AmpC Lactamase-Producing spp, , , spp, or : A Pilot Multicenter Randomized Controlled Trial (MERINO-2). <i>Open Forum Infectious Diseases</i> , 2021 , 8, ofab387 | 1 | 11 |
| 509 | Pharmacokinetics of fluconazole and ganciclovir as combination antimicrobial chemotherapy on ECMO: a case report. <i>International Journal of Antimicrobial Agents</i> , 2021 , 58, 106431 | 14.3 | O |
| 508 | Global personalization of antibiotic therapy in critically ill patients. <i>Expert Review of Precision Medicine and Drug Development</i> , 2021 , 6, 87-93 | 1.6 | 0 |
| 507 | Population pharmacokinetics of intra-peritoneal gentamicin and the impact of varying dwell times on pharmacodynamic target attainment in patients with acute peritonitis undergoing peritoneal dialysis <i>Antimicrobial Agents and Chemotherapy</i> , 2021 , AAC0167921 | 5.9 | Ο |
| 506 | An international survey of adherence to Surviving Sepsis Campaign Guidelines 2016 regarding fluid resuscitation and vasopressors in the initial management of septic shock <i>Journal of Critical Care</i> , 2021 , | 4 | 1 |
| 505 | Caspofungin Population Pharmacokinetic Analysis in Plasma and Peritoneal Fluid in Septic Patients with Intra-Abdominal Infections: A Prospective Cohort Study <i>Clinical Pharmacokinetics</i> , 2021 , 1 | 6.2 | |

| 504 | Pulmonary infections complicating ARDS. <i>Intensive Care Medicine</i> , 2020 , 46, 2168-2183 | 14.5 | 19 |
|-----|--|--------------|----|
| 503 | An Integrated Dialysis Pharmacometric (IDP) Model to Evaluate the Pharmacokinetics in Patients Undergoing Renal Replacement Therapy. <i>Pharmaceutical Research</i> , 2020 , 37, 96 | 4.5 | 4 |
| 502 | Pharmacokinetics of Sulfamethoxazole and Trimethoprim During Venovenous Extracorporeal Membrane Oxygenation: A Case Report. <i>Pharmacotherapy</i> , 2020 , 40, 713-717 | 5.8 | 1 |
| 501 | Comparative Plasma Pharmacokinetics of Ceftriaxone and Ertapenem in Normoalbuminemia, Hypoalbuminemia, and Albumin Replacement in a Sheep Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2020 , 64, | 5.9 | 2 |
| 500 | Antimicrobial de-escalation in the critically ill patient and assessment of clinical cure: the DIANA study. <i>Intensive Care Medicine</i> , 2020 , 46, 1404-1417 | 14.5 | 18 |
| 499 | Pharmacokinetic and pharmacodynamic considerations in antimicrobial therapy for sepsis. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2020 , 16, 415-430 | 5.5 | 7 |
| 498 | Development of Population and Bayesian Models for Applied Use in Patients Receiving Cefepime. <i>Clinical Pharmacokinetics</i> , 2020 , 59, 1027-1036 | 6.2 | 5 |
| 497 | Conventional Pig as Animal Model for Human Renal Drug Excretion Processes: Unravelling the Porcine Renal Function by Use of a Cocktail of Exogenous Markers. <i>Frontiers in Pharmacology</i> , 2020 , 11, 883 | 5.6 | 6 |
| 496 | Pharmacodynamic evaluation of intermittent versus extended and continuous infusions of piperacillin/tazobactam in a hollow-fibre infection model against Klebsiella pneumoniae. <i>Journal of Antimicrobial Chemotherapy</i> , 2020 , 75, 2633-2640 | 5.1 | 3 |
| 495 | Prophylactic Cefazolin Dosing in Women With Body Mass Index >35 kg[m-2 Undergoing Cesarean Delivery: A Pharmacokinetic Study of Plasma and Interstitial Fluid. <i>Anesthesia and Analgesia</i> , 2020 , 131, 199-207 | 3.9 | 6 |
| 494 | Integrating multiple genomic technologies to investigate an outbreak of carbapenemase-producing Enterobacter hormaechei. <i>Nature Communications</i> , 2020 , 11, 466 | 17.4 | 14 |
| 493 | Antimicrobial-associated harm in critical care: a narrative review. <i>Intensive Care Medicine</i> , 2020 , 46, 225- | 28 45 | 36 |
| 492 | Emerging Treatment Options for Infections by Multidrug-Resistant Gram-Positive Microorganisms. <i>Microorganisms</i> , 2020 , 8, | 4.9 | 17 |
| 491 | Pharmacokinetic/pharmacodynamic adequacy of polymyxin B against extensively drug-resistant Gram-negative bacteria in critically ill, general ward and cystic fibrosis patient populations. <i>International Journal of Antimicrobial Agents</i> , 2020 , 55, 105943 | 14.3 | 6 |
| 490 | On-Site Therapeutic Drug Monitoring. <i>Trends in Biotechnology</i> , 2020 , 38, 1262-1277 | 15.1 | 55 |
| 489 | There is no fire without smoke! Pathophysiology and treatment of inhalational injury in burns: A narrative review. <i>Anaesthesia and Intensive Care</i> , 2020 , 48, 114-122 | 1.1 | 1 |
| 488 | International facilitators and barriers to family engagement in the ICU: Results of a qualitative analysis. <i>Journal of Critical Care</i> , 2020 , 58, 72-77 | 4 | 11 |
| 487 | Lactam pharmacodynamics in Gram-negative bloodstream infections in the critically ill. <i>Journal of Antimicrobial Chemotherapy</i> , 2020 , 75, 429-433 | 5.1 | 25 |

| 486 | Ventilator-Associated Tracheobronchitis: To Treat or Not to Treat?. Antibiotics, 2020, 9, | 4.9 | 2 |
|-----|--|------|----|
| 485 | Optimization of flucloxacillin dosing regimens in critically ill patients using population pharmacokinetic modelling of total and unbound concentrations. <i>Journal of Antimicrobial Chemotherapy</i> , 2020 , 75, 2641-2649 | 5.1 | 8 |
| 484 | Elactam antibiotic versus combined Elactam antibiotics and single daily dosing regimens of aminoglycosides for treating serious infections: A meta-analysis. <i>International Journal of Antimicrobial Agents</i> , 2020 , 55, 105839 | 14.3 | 14 |
| 483 | Development and validation of LC-MS/MS methods to measure tobramycin and lincomycin in plasma, microdialysis fluid and urine: application to a pilot pharmacokinetic research study. <i>Clinical Chemistry and Laboratory Medicine</i> , 2020 , 58, 274-284 | 5.9 | 3 |
| 482 | Cerebrospinal Fluid Penetration of Ceftolozane-Tazobactam in Critically Ill Patients with an Indwelling External Ventricular Drain. <i>Antimicrobial Agents and Chemotherapy</i> , 2020 , 65, | 5.9 | 6 |
| 481 | A Systematic Review of Studies Reporting Antibiotic Pharmacokinetic Data in the Cerebrospinal Fluid of Critically Ill Patients with Uninflamed Meninges. <i>Antimicrobial Agents and Chemotherapy</i> , 2020 , 65, | 5.9 | 4 |
| 480 | Pharmacodynamic Evaluation of Plasma and Epithelial Lining Fluid Exposures of Amikacin against Pseudomonas aeruginosa in a Dynamic Hollow-Fiber Infection Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2020 , 64, | 5.9 | 2 |
| 479 | Vancomycin Serum Concentration after 48 h of Administration: A 3-Years Survey in an Intensive Care Unit. <i>Antibiotics</i> , 2020 , 9, | 4.9 | 2 |
| 478 | Infection, Sepsis and the Inflammatory Response: Mechanisms and Therapy. <i>Frontiers in Medicine</i> , 2020 , 7, 588863 | 4.9 | 4 |
| 477 | Prolonged Versus Intermittent Infusion of £Lactam Antibiotics: A Systematic Review and Meta-Regression of Bacterial Killing in Preclinical Infection Models. <i>Clinical Pharmacokinetics</i> , 2020 , 59, 1237-1250 | 6.2 | 7 |
| 476 | Ticarcillin and piperacillin adsorption on to polyethersulfone haemodiafilter membranes in an ex-vivo circuit. <i>International Journal of Antimicrobial Agents</i> , 2020 , 56, 106058 | 14.3 | 1 |
| 475 | When not to start antibiotics: avoiding antibiotic overuse in the intensive care unit. <i>Clinical Microbiology and Infection</i> , 2020 , 26, 35-40 | 9.5 | 19 |
| 474 | Pharmacokinetics of Total and Unbound Cefazolin during Veno-Arterial Extracorporeal Membrane Oxygenation: A Case Report. <i>Chemotherapy</i> , 2019 , 64, 115-118 | 3.2 | 2 |
| 473 | Lefamulin. Comment on: "Novel Antibiotics for Multidrug-Resistant Gram-Positive Microorganisms., 2019, , 270". <i>Microorganisms</i> , 2019 , 7, | 4.9 | 8 |
| 472 | Saturable elimination of piperacillin in critically ill patients: implications for continuous infusion. <i>International Journal of Antimicrobial Agents</i> , 2019 , 54, 741-749 | 14.3 | 8 |
| 471 | The relationship between ventilator-associated pneumonia and chronic obstructive pulmonary disease: what is the current evidence?. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2019 , 38, 637-647 | 5.3 | 6 |
| 470 | Population pharmacokinetics of total and unbound concentrations of intravenous posaconazole in adult critically ill patients. <i>Critical Care</i> , 2019 , 23, 205 | 10.8 | 15 |
| 469 | Antibiotic exposure at the site of infection: principles and assessment of tissue penetration. <i>Expert Review of Clinical Pharmacology</i> , 2019 , 12, 623-634 | 3.8 | 17 |

| 468 | Evaluation of pharmacokinetic/pharmacodynamic and clinical outcomes with 6-hourly empiric piperacillin-tazobactam dosing in hematological malignancy patients with febrile neutropenia. Journal of Infection and Chemotherapy, 2019 , 25, 503-508 | 2.2 | 5 |
|-----------------|---|------|----|
| 467 | Analysis of capillary microsamples obtained from a skin-prick to measure vancomycin concentrations as a valid alternative to conventional sampling: A bridging study. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2019 , 169, 288-292 | 3.5 | 4 |
| 466 | Early target attainment of continuous infusion piperacillin/tazobactam and meropenem in critically ill patients: A prospective observational study. <i>Journal of Critical Care</i> , 2019 , 52, 75-79 | 4 | 17 |
| 465 | In-vitro adsorption and sieving coefficient of ticarcillin-clavulanate during continuous haemofiltration. <i>International Journal of Antimicrobial Agents</i> , 2019 , 54, 261-264 | 14.3 | 6 |
| 464 | Challenges in the management of septic shock: a narrative review. <i>Intensive Care Medicine</i> , 2019 , 45, 420-433 | 14.5 | 22 |
| 463 | A Robust Statistical Approach to Analyse Population Pharmacokinetic Data in Critically Ill Patients Receiving Renal Replacement Therapy. <i>Clinical Pharmacokinetics</i> , 2019 , 58, 263-270 | 6.2 | |
| 462 | Novel Antibiotics for Multidrug-Resistant Gram-Positive Microorganisms. <i>Microorganisms</i> , 2019 , 7, | 4.9 | 37 |
| 461 | What Antibiotic Exposures Are Required to Suppress the Emergence of Resistance for Gram-Negative Bacteria? A Systematic Review. <i>Clinical Pharmacokinetics</i> , 2019 , 58, 1407-1443 | 6.2 | 47 |
| 460 | A Multicenter Trial of Vena Cava Filters in Severely Injured Patients. <i>New England Journal of Medicine</i> , 2019 , 381, 328-337 | 59.2 | 69 |
| 459 | Population pharmacokinetics of ticarcillin in critically ill patients receiving extended daily diafiltration. <i>International Journal of Antimicrobial Agents</i> , 2019 , 54, 351-355 | 14.3 | 1 |
| 458 | Population Pharmacokinetics of Unbound Ceftolozane and Tazobactam in Critically Ill Patients without Renal Dysfunction. <i>Antimicrobial Agents and Chemotherapy</i> , 2019 , 63, | 5.9 | 25 |
| 457 | Epidemiology of intra-abdominal infection and sepsis in critically ill patients: "AbSeS", a multinational observational cohort study and ESICM Trials Group Project. <i>Intensive Care Medicine</i> , 2019 , 45, 1703-1717 | 14.5 | 40 |
| 456 | Defining optimal dosing of ciprofloxacin in patients with septic shock. <i>Journal of Antimicrobial Chemotherapy</i> , 2019 , 74, 1662-1669 | 5.1 | 11 |
| 455 | 1573. Population Pharmacokinetic Analyses for Cefepime in Adult and Pediatric Patients. <i>Open Forum Infectious Diseases</i> , 2019 , 6, S574-S575 | 1 | 78 |
| 454 | Amoxicillin-Clavulanate Dosing in the Intensive Care Unit: The Additive Effect of Renal Replacement Therapy in a Patient with Normal Kidney Function. <i>Chemotherapy</i> , 2019 , 64, 173-176 | 3.2 | 3 |
| 453 | Meropenem-Tobramycin Combination Regimens Combat Carbapenem-Resistant Pseudomonas aeruginosa in the Hollow-Fiber Infection Model Simulating Augmented Renal Clearance in Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2019 , 64, | 5.9 | 7 |
| 452 | A validated LC-MSMS method for the simultaneous quantification of meropenem and vaborbactam in human plasma and renal replacement therapy effluent and its application to a pharmacokinetic study. <i>Analytical and Bioanalytical Chemistry</i> , 2019 , 411, 7831-7840 | 4.4 | 7 |
| 45 ¹ | A Population Pharmacokinetic Model-Guided Evaluation of Ceftolozane-Tazobactam Dosing in Critically Ill Patients Undergoing Continuous Venovenous Hemodiafiltration. <i>Antimicrobial Agents and Chemotherapy</i> , 2019 , 64. | 5.9 | 18 |

| 450 | Pharmacokinetics of Enteric-Coated Mycophenolate Sodium in Lupus Nephritis (POEMSLUN). <i>Therapeutic Drug Monitoring</i> , 2019 , 41, 703-713 | 3.2 | 4 |
|-----|---|------------------|----|
| 449 | Population pharmacokinetics and evaluation of the predictive performance of pharmacokinetic models in critically ill patients receiving continuous infusion meropenem: a comparison of eight pharmacokinetic models. <i>Journal of Antimicrobial Chemotherapy</i> , 2019 , 74, 432-441 | 5.1 | 32 |
| 448 | Intrapulmonary pharmacokinetics of antibiotics used to treat nosocomial pneumonia caused by Gram-negative bacilli: A systematic review. <i>International Journal of Antimicrobial Agents</i> , 2019 , 53, 234-24 | 44 .3 | 25 |
| 447 | Infections by multidrug-resistant Gram-negative Bacteria: WhatIs new in our arsenal and whatIs in the pipeline?. <i>International Journal of Antimicrobial Agents</i> , 2019 , 53, 211-224 | 14.3 | 46 |
| 446 | Plasma and interstitial fluid population pharmacokinetics of vancomycin in critically ill patients with sepsis. <i>International Journal of Antimicrobial Agents</i> , 2019 , 53, 137-142 | 14.3 | 7 |
| 445 | A protocol for a phase 3 multicentre randomised controlled trial of continuous versus intermittent Elactam antibiotic infusion in critically ill patients with sepsis: BLING III. <i>Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine</i> , 2019 , 21, 63-68 | 2.8 | 9 |
| 444 | Survey of critical care practice in Australian and New Zealand burn referral centres. <i>Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine</i> , 2019 , 21, 303-304 | 2.8 | 1 |
| 443 | Pharmacokinetics of piperacillin in critically ill patients with acute kidney injury receiving sustained low-efficiency diafiltration. <i>Journal of Antimicrobial Chemotherapy</i> , 2018 , 73, 1647-1650 | 5.1 | 12 |
| 442 | Optimization and Evaluation of Piperacillin-Tobramycin Combination Dosage Regimens against Pseudomonas aeruginosa for Patients with Altered Pharmacokinetics via the Hollow-Fiber Infection Model and Mechanism-Based Modeling. <i>Antimicrobial Agents and Chemotherapy</i> , 2018 , 62, | 5.9 | 11 |
| 441 | Utility of community-acquired pneumonia severity scores in guiding disposition from the emergency department: Intensive care or short-stay unit?. <i>EMA - Emergency Medicine Australasia</i> , 2018 , 30, 538-546 | 1.5 | 4 |
| 440 | Pharmacokinetics of Intravenous Posaconazole in Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2018 , 62, | 5.9 | 12 |
| 439 | Re: Is a randomized trial of a short course of aminoglycoside added to Elactam antibiotics for empirical treatment in critically ill patients with sepsis justified?L Pharmacokinetic/pharmacodynamic considerations. Clinical Microbiology and Infection, 2018, 24, 666-667. | 9.5 7 | 1 |
| 438 | Antimicrobial resistance and antibiotic stewardship programs in the ICU: insistence and persistence in the fight against resistance. A position statement from ESICM/ESCMID/WAAAR round table on multi-drug resistance. <i>Intensive Care Medicine</i> , 2018 , 44, 189-196 | 14.5 | 65 |
| 437 | Differences in suppression of regrowth and resistance despite similar initial bacterial killing for meropenem and piperacillin/tazobactam against Pseudomonas aeruginosa and Escherichia coli. <i>Diagnostic Microbiology and Infectious Disease</i> , 2018 , 91, 69-76 | 2.9 | 3 |
| 436 | Individualising Therapy to Minimize Bacterial Multidrug Resistance. <i>Drugs</i> , 2018 , 78, 621-641 | 12.1 | 32 |
| 435 | Clinical application of microsampling versus conventional sampling techniques in the quantitative bioanalysis of antibiotics: a systematic review. <i>Bioanalysis</i> , 2018 , 10, 407-423 | 2.1 | 20 |
| 434 | Population pharmacokinetics of vancomycin in critically ill patients receiving prolonged intermittent renal replacement therapy. <i>International Journal of Antimicrobial Agents</i> , 2018 , 52, 151-157 | 14.3 | 13 |
| 433 | Maximally effective dosing regimens of meropenem in patients with septic shock. <i>Journal of Antimicrobial Chemotherapy</i> , 2018 , 73, 191-198 | 5.1 | 29 |

| 432 | A UHPLC-MS/MS method for the simultaneous determination of piperacillin and tazobactam in plasma (total and unbound), urine and renal replacement therapy effluent. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018 , 148, 324-333 | 3.5 | 15 |
|-----|---|------|-----|
| 431 | The effects of major burn related pathophysiological changes on the pharmacokinetics and pharmacodynamics of drug use: An appraisal utilizing antibiotics. <i>Advanced Drug Delivery Reviews</i> , 2018 , 123, 65-74 | 18.5 | 27 |
| 430 | Antibiotic Distribution into Cerebrospinal Fluid: Can Dosing Safely Account for Drug and Disease Factors in the Treatment of Ventriculostomy-Associated Infections?. <i>Clinical Pharmacokinetics</i> , 2018 , 57, 439-454 | 6.2 | 14 |
| 429 | Characteristics, treatment and outcomes for all emergency department patients fulfilling criteria for septic shock: a prospective observational study. <i>European Journal of Emergency Medicine</i> , 2018 , 25, 97-104 | 2.3 | 7 |
| 428 | A pharmacokinetic case study of intravenous posaconazole in a critically ill patient with hypoalbuminaemia receiving continuous venovenous haemodiafiltration. <i>International Journal of Antimicrobial Agents</i> , 2018 , 52, 506-509 | 14.3 | 5 |
| 427 | Therapeutic drug monitoring of Elactam antibiotics in the critically ill: direct measurement of unbound drug concentrations to achieve appropriate drug exposures. <i>Journal of Antimicrobial Chemotherapy</i> , 2018 , 73, 3087-3094 | 5.1 | 63 |
| 426 | Characterization of Effects of Renal Replacement Therapy Modalities and Settings on Pharmacokinetics of Meropenem and Vaborbactam. <i>Antimicrobial Agents and Chemotherapy</i> , 2018 , 62, | 5.9 | 18 |
| 425 | New paradigm for rapid achievement of appropriate therapy in special populations: coupling antibiotic dose optimization rapid microbiological methods. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2018 , 14, 693-708 | 5.5 | 8 |
| 424 | Optimised dosing of vancomycin in critically ill Indigenous Australian patients with severe sepsis. <i>Anaesthesia and Intensive Care</i> , 2018 , 46, 374-380 | 1.1 | 7 |
| 423 | Prevalence of augmented renal clearance and performance of glomerular filtration estimates in Indigenous Australian patients requiring intensive care admission. <i>Anaesthesia and Intensive Care</i> , 2018 , 46, 42-50 | 1.1 | 8 |
| 422 | A comparison between raw and predicted mortality in a paediatric intensive care unit in South Africa. <i>BMC Research Notes</i> , 2018 , 11, 829 | 2.3 | 2 |
| 421 | The Statistical Curriculum Within Randomized Controlled Trials in Critical Illness. <i>Critical Care Medicine</i> , 2018 , 46, 1985-1990 | 1.4 | 2 |
| 420 | The authors reply. Critical Care Medicine, 2018, 46, e725-e726 | 1.4 | |
| 419 | Strategies to reduce inappropriate laboratory blood test orders in intensive care are effective and safe: a before-and-after quality improvement study. <i>Anaesthesia and Intensive Care</i> , 2018 , 46, 313-320 | 1.1 | 8 |
| 418 | A research pathway for the study of the delivery and disposition of nebulised antibiotics: an incremental approach from in vitro to large animal models. <i>Intensive Care Medicine Experimental</i> , 2018 , 6, 17 | 3.7 | 4 |
| 417 | Sepsis in Intensive Care Unit Patients: Worldwide Data From the Intensive Care over Nations Audit. <i>Open Forum Infectious Diseases</i> , 2018 , 5, ofy313 | 1 | 120 |
| 416 | Patient and family engagement in the ICU: Report from the task force of the World Federation of Societies of Intensive and Critical Care Medicine. <i>Journal of Critical Care</i> , 2018 , 48, 251-256 | 4 | 40 |
| 415 | Population pharmacokinetics of intravenous paracetamol in critically ill patients with traumatic brain injury. <i>Journal of Critical Care</i> , 2018 , 47, 15-20 | 4 | 3 |

(2017-2017)

| 414 | Augmented Renal Clearance in Traumatic Brain Injury: A Single-Center Observational Study of Atrial Natriuretic Peptide, Cardiac Output, and Creatinine Clearance. <i>Journal of Neurotrauma</i> , 2017 , 34, 137-144 | 5.4 | 33 |
|-----|---|--------|----|
| 413 | Population Pharmacokinetics of Piperacillin in Nonobese, Obese, and Morbidly Obese Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2017 , 61, | 5.9 | 33 |
| 412 | The intensive care medicine research agenda on multidrug-resistant bacteria, antibiotics, and stewardship. <i>Intensive Care Medicine</i> , 2017 , 43, 1187-1197 | 14.5 | 66 |
| 411 | Pyroglutamate (5-oxoproline) measured with hydrophilic interaction chromatography (HILIC) tandem mass spectrometry in acutely ill patients. <i>Clinica Chimica Acta</i> , 2017 , 466, 72-77 | 6.2 | 2 |
| 410 | Substantial Impact of Altered Pharmacokinetics in Critically Ill Patients on the Antibacterial Effects of Meropenem Evaluated via the Dynamic Hollow-Fiber Infection Model. <i>Antimicrobial Agents and Chemotherapy</i> , 2017 , 61, | 5.9 | 20 |
| 409 | Association between augmented renal clearance and clinical outcomes in patients receiving Elactam antibiotic therapy by continuous or intermittent infusion: a nested cohort study of the BLING-II randomised, placebo-controlled, clinical trial. <i>International Journal of Antimicrobial Agents</i> , | 14.3 | 56 |
| 408 | Dosing antibiotic prophylaxis during cardiopulmonary bypass-a higher level of complexity? A structured review. <i>International Journal of Antimicrobial Agents</i> , 2017 , 49, 395-402 | 14.3 | 13 |
| 407 | "Salvage treatment" for infections by extensively- and pan-drug-resistant pathogens is common and often sub-optimal. <i>Intensive Care Medicine</i> , 2017 , 43, 1164-1166 | 14.5 | 6 |
| 406 | Factors influencing psychological, social and health outcomes after major burn injuries in adults: cohort study protocol. <i>BMJ Open</i> , 2017 , 7, e017545 | 3 | 13 |
| 405 | Comparative total and unbound pharmacokinetics of cefazolin administered by bolus versus continuous infusion in patients undergoing major surgery: a randomized controlled trial. <i>British Journal of Anaesthesia</i> , 2017 , 118, 876-882 | 5.4 | 10 |
| 404 | Population Pharmacokinetics of Tigecycline in Critically Ill Patients with Severe Infections. <i>Antimicrobial Agents and Chemotherapy</i> , 2017 , 61, | 5.9 | 30 |
| 403 | Antibiotic dosing for multidrug-resistant pathogen pneumonia. <i>Current Opinion in Infectious Diseases</i> , 2017 , 30, 231-239 | 5.4 | 6 |
| 402 | Right Dose, Right Now: Customized Drug Dosing in the Critically Ill. Critical Care Medicine, 2017, 45, 331 | -3:346 | 45 |
| 401 | WhatIs new in pharmacokinetics of antimicrobials in AKI and RRT?. <i>Intensive Care Medicine</i> , 2017 , 43, 904-906 | 14.5 | 5 |
| 400 | Impact of Elactam antibiotic therapeutic drug monitoring on dose adjustments in critically ill patients undergoing continuous renal replacement therapy. <i>International Journal of Antimicrobial Agents</i> , 2017 , 49, 589-594 | 14.3 | 34 |
| 399 | Caspofungin Population Pharmacokinetics in Critically Ill Patients Undergoing Continuous Veno-Venous Haemofiltration or Haemodiafiltration. <i>Clinical Pharmacokinetics</i> , 2017 , 56, 1057-1068 | 6.2 | 25 |
| 398 | An LC-MS/MS method to determine vancomycin in plasma (total and unbound), urine and renal replacement therapy effluent. <i>Bioanalysis</i> , 2017 , 9, 911-924 | 2.1 | 11 |
| 397 | Detailed assessment of benefits and risks of retrievable inferior vena cava filters on patients with complicated injuries: the da Vinci multicentre randomised controlled trial study protocol. <i>BMJ Open</i> , 2017 , 7, e016747 | 3 | 9 |

| 396 | Using Population Pharmacokinetic Modeling and Monte Carlo Simulations To Determine whether Standard Doses of Piperacillin in Piperacillin-Tazobactam Regimens Are Adequate for the Management of Febrile Neutropenia. <i>Antimicrobial Agents and Chemotherapy</i> , 2017 , 61, | 5.9 | 13 |
|-----|---|---------|-----|
| 395 | A validated UHPLC-MS/MS method for the measurement of riluzole in plasma and myocardial tissue samples. <i>Biomedical Chromatography</i> , 2017 , 31, e4030 | 1.7 | 1 |
| 394 | Impact of renal replacement modalities on the clearance of piperacillin-tazobactam administered via continuous infusion in critically ill patients. <i>International Journal of Antimicrobial Agents</i> , 2017 , 50, 227-231 | 14.3 | 14 |
| 393 | Hemoglobin concentrations and RBC transfusion thresholds in patients with acute brain injury: an international survey. <i>Critical Care</i> , 2017 , 21, 159 | 10.8 | 24 |
| 392 | Proposed primary endpoints for use in clinical trials that compare treatment options for bloodstream infection in adults: a consensus definition. <i>Clinical Microbiology and Infection</i> , 2017 , 23, 533 | 3-25-41 | 36 |
| 391 | Pharmacodynamics of Aerosolized Fosfomycin and Amikacin against Resistant Clinical Isolates of Pseudomonas aeruginosa and Klebsiella pneumoniae in a Hollow-Fiber Infection Model: Experimental Basis for Combination Therapy. <i>Antimicrobial Agents and Chemotherapy</i> , 2017 , 61, | 5.9 | 16 |
| 390 | Does Furosemide Increase Oxidative Stress in Acute Kidney Injury?. <i>Antioxidants and Redox Signaling</i> , 2017 , 26, 221-226 | 8.4 | 16 |
| 389 | Interpretation of Care Guidelines for Obese Women in Labor: Intergroup Language and Social Identity. <i>Journal of Language and Social Psychology</i> , 2017 , 36, 388-414 | 1.2 | 1 |
| 388 | Systemic Inflammatory Response Syndrome, Quick Sequential Organ Function Assessment, and Organ Dysfunction: Insights From a Prospective Database of ED Patients With Infection. <i>Chest</i> , 2017 , 151, 586-596 | 5.3 | 178 |
| 387 | Population Pharmacokinetics of Doripenem in Critically Ill Patients with Sepsis in a Malaysian Intensive Care Unit. <i>Antimicrobial Agents and Chemotherapy</i> , 2016 , 60, 206-14 | 5.9 | 11 |
| 386 | Understanding PK/PD. Intensive Care Medicine, 2016, 42, 1797-1800 | 14.5 | 38 |
| 385 | Stability of Antibiotics for Intraperitoneal Administration in Extraneal 7.5% Icodextrin Peritoneal Dialysis Bags (STAB Study). <i>Peritoneal Dialysis International</i> , 2016 , 36, 421-6 | 2.8 | 8 |
| 384 | Uncertainty in Antibiotic Dosing in Critically Ill Neonate and Pediatric Patients: Can Microsampling Provide the Answers?. <i>Clinical Therapeutics</i> , 2016 , 38, 1961-75 | 3.5 | 19 |
| 383 | Integrated pharmacokinetic-pharmacodynamic modelling to evaluate antimicrobial prophylaxis in abdominal surgery. <i>Journal of Antimicrobial Chemotherapy</i> , 2016 , 71, 2902-8 | 5.1 | 7 |
| 382 | Pharmacokinetics of Piperacillin in Critically Ill Australian Indigenous Patients with Severe Sepsis. <i>Antimicrobial Agents and Chemotherapy</i> , 2016 , 60, 7402-7406 | 5.9 | 13 |
| 381 | Total and unbound ceftriaxone pharmacokinetics in critically ill Australian Indigenous patients with severe sepsis. <i>International Journal of Antimicrobial Agents</i> , 2016 , 48, 748-752 | 14.3 | 17 |
| 380 | Determination of Cefalothin and Cefazolin in Human Plasma, Urine and Peritoneal Dialysate by UHPLC-MS/MS: application to a pilot pharmacokinetic study in humans. <i>Biomedical Chromatography</i> , 2016 , 30, 872-9 | 1.7 | 16 |
| 379 | We need to optimize piperacillin-tazobactam dosing in critically ill patients-but how?. <i>Critical Care</i> , 2016 , 20, 163 | 10.8 | 6 |

| 378 | "Not out of the woods"a wifels perspective: bedside communication. <i>Intensive Care Medicine</i> , 2016 , 42, 446-447 | 14.5 | 4 |
|-----|--|------|-----|
| 377 | Effect of different renal function on antibacterial effects of piperacillin against Pseudomonas aeruginosa evaluated via the hollow-fibre infection model and mechanism-based modelling. <i>Journal of Antimicrobial Chemotherapy</i> , 2016 , 71, 2509-20 | 5.1 | 26 |
| 376 | CAMERA2 - combination antibiotic therapy for methicillin-resistant Staphylococcus aureus infection: study protocol for a randomised controlled trial. <i>Trials</i> , 2016 , 17, 170 | 2.8 | 45 |
| 375 | Influence of Renal Replacement Modalities on Amikacin Population Pharmacokinetics in Critically Ill Patients on Continuous Renal Replacement Therapy. <i>Antimicrobial Agents and Chemotherapy</i> , 2016 , 60, 4901-9 | 5.9 | 19 |
| 374 | Severity Scores in Emergency Department Patients With Presumed Infection: A Prospective Validation Study. <i>Critical Care Medicine</i> , 2016 , 44, 539-47 | 1.4 | 36 |
| 373 | Beta-Lactam Infusion in Severe Sepsis (BLISS): a prospective, two-centre, open-labelled randomised controlled trial of continuous versus intermittent beta-lactam infusion in critically ill patients with severe sepsis. <i>Intensive Care Medicine</i> , 2016 , 42, 1535-1545 | 14.5 | 172 |
| 372 | Pharmacokinetics of Intraperitoneal Cefalothin and Cefazolin in Patients Being Treated for Peritoneal Dialysis-Associated Peritonitis. <i>Peritoneal Dialysis International</i> , 2016 , 36, 415-20 | 2.8 | 5 |
| 371 | What is the effect of obesity on piperacillin and meropenem trough concentrations in critically ill patients?. <i>Journal of Antimicrobial Chemotherapy</i> , 2016 , 71, 696-702 | 5.1 | 24 |
| 370 | In Vivo Microdialysis To Determine Subcutaneous Interstitial Fluid Penetration and Pharmacokinetics of Fluconazole in Intensive Care Unit Patients with Sepsis. <i>Antimicrobial Agents and Chemotherapy</i> , 2016 , 60, 827-32 | 5.9 | 13 |
| 369 | Continuous versus Intermittent Lactam Infusion in Severe Sepsis. A Meta-analysis of Individual Patient Data from Randomized Trials. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2016 , 194, 681-91 | 10.2 | 192 |
| 368 | Effect of obesity on the pharmacokinetics of antimicrobials in critically ill patients: A structured review. <i>International Journal of Antimicrobial Agents</i> , 2016 , 47, 259-68 | 14.3 | 63 |
| 367 | The use and risks of antibiotics in critically ill patients. Expert Opinion on Drug Safety, 2016, 15, 667-78 | 4.1 | 16 |
| 366 | SaMpling Antibiotics in Renal Replacement Therapy (SMARRT): an observational pharmacokinetic study in critically ill patients. <i>BMC Infectious Diseases</i> , 2016 , 16, 103 | 4 | 11 |
| 365 | Comparison of equal doses of continuous venovenous haemofiltration and haemodiafiltration on ciprofloxacin population pharmacokinetics in critically ill patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2016 , 71, 1643-50 | 5.1 | 12 |
| 364 | A Systematic Review of the Definitions, Determinants, and Clinical Outcomes of Antimicrobial De-escalation in the Intensive Care Unit. <i>Clinical Infectious Diseases</i> , 2016 , 62, 1009-1017 | 11.6 | 111 |
| 363 | Population pharmacokinetics of linezolid in critically ill patients on renal replacement therapy: comparison of equal doses in continuous venovenous haemofiltration and continuous venovenous haemodiafiltration. <i>Journal of Antimicrobial Chemotherapy</i> , 2016 , 71, 464-70 | 5.1 | 23 |
| 362 | Is prolonged infusion of piperacillin/tazobactam and meropenem in critically ill patients associated with improved pharmacokinetic/pharmacodynamic and patient outcomes? An observation from the Defining Antibiotic Levels in Intensive care unit patients (DALI) cohort. <i>Journal of Antimicrobial</i> | 5.1 | 104 |
| 361 | Chemotherapy, 2016 , 71, 196-207 Anaesthetic management of obese parturients: what is the evidence supporting practice guidelines?. Anaesthesia and Intensive Care, 2016 , 44, 552-9 | 1.1 | 2 |

| 360 | Bugs, genes, and the intensive care unit. <i>Journal of Thoracic Disease</i> , 2016 , 8, E788-90 | 2.6 | 1 |
|-----|---|-----------|--------------|
| 359 | AnaesthetistsLexperiences with the early labour epidural recommendation for obese parturients: a qualitative study. <i>Anaesthesia and Intensive Care</i> , 2016 , 44, 620-7 | 1.1 | 3 |
| 358 | EPHA4-FC TREATMENT REDUCES ISCHEMIA/REPERFUSION-INDUCED INTESTINAL INJURY BY INHIBITING VASCULAR PERMEABILITY. <i>Shock</i> , 2016 , 45, 184-91 | 3.4 | 10 |
| 357 | The Value of E-Learning for the Prevention of Healthcare-Associated Infections. <i>Infection Control and Hospital Epidemiology</i> , 2016 , 37, 1052-9 | 2 | 13 |
| 356 | Therapeutic drug monitoring of anti-infective agents in critically ill patients. <i>Expert Review of Clinical Pharmacology</i> , 2016 , 9, 961-79 | 3.8 | 77 |
| 355 | Is inhaled prophylactic heparin useful for prevention and Management of Pneumonia in ventilated ICU patients?: The IPHIVAP investigators of the Australian and New Zealand Intensive Care Society Clinical Trials Group. <i>Journal of Critical Care</i> , 2016 , 34, 95-102 | 4 | 15 |
| 354 | New Regimen for Continuous Infusion of Vancomycin in Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2016 , 60, 4750-6 | 5.9 | 35 |
| 353 | Effect of Obesity on the Population Pharmacokinetics of Meropenem in Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2016 , 60, 4577-84 | 5.9 | 27 |
| 352 | Determinants of Urinary Output Response to IV Furosemide in Acute Kidney Injury: A Pharmacokinetic/Pharmacodynamic Study. <i>Critical Care Medicine</i> , 2016 , 44, e923-9 | 1.4 | 15 |
| 351 | Is there a role for microsampling in antibiotic pharmacokinetic studies?. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2016 , 12, 601-14 | 5.5 | 16 |
| 350 | Should Elactam antibiotics be administered by continuous infusion in critically ill patients? A survey of Australia and New Zealand intensive care unit doctors and pharmacists. <i>International Journal of Antimicrobial Agents</i> , 2016 , 47, 436-8 | 14.3 | 13 |
| 349 | Effect of Obesity on the Population Pharmacokinetics of Fluconazole in Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2016 , 60, 6550-6557 | 5.9 | 26 |
| 348 | Problematic Dichotomization of Risk for Intensive Care Unit (ICU)-Acquired Invasive Candidiasis: Results Using a Risk-Predictive Model to Categorize 3 Levels of Risk From a Multicenter Prospective Cohort of Australian ICU Patients. <i>Clinical Infectious Diseases</i> , 2016 , 63, 1463-1469 | 11.6 | 29 |
| 347 | Effect of time on recovery of plasma microsamples for the quantitative determination of vancomycin. <i>Bioanalysis</i> , 2016 , 8, 2235-2242 | 2.1 | 25 |
| 346 | Sepsis: frontiers in diagnosis, resuscitation and antibiotic therapy. <i>Intensive Care Medicine</i> , 2016 , 42, 19 | 958-41.96 | 59 81 |
| 345 | Optimising meropenem dosing in critically ill Australian Indigenous patients with severe sepsis. <i>International Journal of Antimicrobial Agents</i> , 2016 , 48, 542-546 | 14.3 | 23 |
| 344 | Use of neostigmine for acute colonic pseudo-obstruction in a patient receiving dexmedetomidine. Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine, 2016 , 18, 59-61 | 2.8 | 1 |
| 343 | Is high-dose Elactam therapy associated with excessive drug toxicity in critically ill patients?. <i>Minerva Anestesiologica</i> , 2016 , 82, 957-65 | 1.9 | 19 |

(2015-2015)

| 342 | Candida colonization as a risk marker for invasive candidiasis in mixed medical-surgical intensive care units: development and evaluation of a simple, standard protocol. <i>Journal of Clinical Microbiology</i> , 2015 , 53, 1324-30 | 9.7 | 32 |
|---|---|--|----------------|
| 341 | From the intensive care bed: we need to hear and listen. <i>Intensive Care Medicine</i> , 2015 , 41, 921-2 | 14.5 | 1 |
| 340 | A meta-analysis of ventriculostomy-associated cerebrospinal fluid infections. <i>BMC Infectious Diseases</i> , 2015 , 15, 3 | 4 | 70 |
| 339 | Pharmacokinetic variability and exposures of fluconazole, anidulafungin, and caspofungin in intensive care unit patients: Data from multinational Defining Antibiotic Levels in Intensive care unit (DALI) patients Study. <i>Critical Care</i> , 2015 , 19, 33 | 10.8 | 83 |
| 338 | Pharmacokinetics of piperacillin in critically ill patients receiving continuous venovenous haemofiltration: A randomised controlled trial of continuous infusion versus intermittent bolus administration. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 39-44 | 14.3 | 23 |
| 337 | Influence of sustained low-efficiency diafiltration (SLED-f) on interstitial fluid concentrations of fluconazole in a critically ill patient: Use of microdialysis. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 121-4 | 14.3 | 5 |
| 336 | A Simulation Study Reveals Lack of Pharmacokinetic/Pharmacodynamic Target Attainment in De-escalated Antibiotic Therapy in Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2015 , 59, 4689-94 | 5.9 | 16 |
| 335 | A validated method for the quantification of fosfomycin on dried plasma spots by HPLC-MS/MS: application to a pilot pharmacokinetic study in humans. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2015 , 115, 509-14 | 3.5 | 18 |
| 334 | A Multicenter Randomized Trial of Continuous versus Intermittent Lactam Infusion in Severe Sepsis. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2015 , 192, 1298-305 | 10.2 | 147 |
| | | | |
| 333 | The authors reply. Critical Care Medicine, 2015, 43, e101-2 | 1.4 | |
| 333 332 | The authors reply. <i>Critical Care Medicine</i> , 2015 , 43, e101-2 Vancomycin-associated nephrotoxicity: A meta-analysis of administration by continuous versus intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 249-53 | 1.4 | 46 |
| | Vancomycin-associated nephrotoxicity: A meta-analysis of administration by continuous versus | | 46 |
| 332 | Vancomycin-associated nephrotoxicity: A meta-analysis of administration by continuous versus intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 249-53 Therapeutic drug monitoring of the Elactam antibiotics: what is the evidence and which patients | 14.3 | |
| 33 ² 33 ¹ | Vancomycin-associated nephrotoxicity: A meta-analysis of administration by continuous versus intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 249-53 Therapeutic drug monitoring of the Elactam antibiotics: what is the evidence and which patients should we be using it for?. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 3178-83 The ADMIN-ICU survey: a survey on antimicrobial dosing and monitoring in ICUs. <i>Journal of</i> | 14.3 5.1 | 88 |
| 332 331 330 | Vancomycin-associated nephrotoxicity: A meta-analysis of administration by continuous versus intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 249-53 Therapeutic drug monitoring of the Elactam antibiotics: what is the evidence and which patients should we be using it for?. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 3178-83 The ADMIN-ICU survey: a survey on antimicrobial dosing and monitoring in ICUs. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 2671-7 | 14.3 5.1 5.1 | 88 |
| 33 ² 33 ¹ 33 ⁰ | Vancomycin-associated nephrotoxicity: A meta-analysis of administration by continuous versus intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 249-53 Therapeutic drug monitoring of the Elactam antibiotics: what is the evidence and which patients should we be using it for?. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 3178-83 The ADMIN-ICU survey: a survey on antimicrobial dosing and monitoring in ICUs. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 2671-7 The authors reply. <i>Critical Care Medicine</i> , 2015 , 43, e154-5 Meropenem versus piperacillin-tazobactam for definitive treatment of bloodstream infections due to ceftriaxone non-susceptible Escherichia coli and Klebsiella spp (the MERINO trial): study protocol | 14.35.15.11.4 | 88 |
| 33 ² 33 ¹ 33 ⁰ 32 ⁹ 32 ⁸ | Vancomycin-associated nephrotoxicity: A meta-analysis of administration by continuous versus intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 249-53 Therapeutic drug monitoring of the Elactam antibiotics: what is the evidence and which patients should we be using it for?. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 3178-83 The ADMIN-ICU survey: a survey on antimicrobial dosing and monitoring in ICUs. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 2671-7 The authors reply. <i>Critical Care Medicine</i> , 2015 , 43, e154-5 Meropenem versus piperacillin-tazobactam for definitive treatment of bloodstream infections due to ceftriaxone non-susceptible Escherichia coli and Klebsiella spp (the MERINO trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2015 , 16, 24 Even high-dose extended infusions may not yield desired concentrations of Elactams: the value of | 14.3 5.1 5.1 1.4 2.8 | 88 82 46 |

| 324 | Albumin concentration significantly impacts on free teicoplanin plasma concentrations in non-critically ill patients with chronic bone sepsis. <i>International Journal of Antimicrobial Agents</i> , 2015 , 45, 647-51 | 14.3 | 13 |
|-----|--|---------------|-----|
| 323 | An UHPLC-MS/MS method for the simultaneous determination of ampicillin and sulbactam in human plasma and urine. <i>Bioanalysis</i> , 2015 , 7, 2311-2319 | 2.1 | 5 |
| 322 | Antibiotic dose optimization in critically ill patients. <i>Medicina Intensiva</i> , 2015 , 39, 563-72 | 1.2 | 27 |
| 321 | Assays for therapeutic drug monitoring of Elactam antibiotics: A structured review. <i>International Journal of Antimicrobial Agents</i> , 2015 , 46, 367-75 | 14.3 | 65 |
| 320 | Population Pharmacokinetics of Fosfomycin in Critically Ill Patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2015 , 59, 6471-6 | 5.9 | 48 |
| 319 | A simple LC-MS/MS method using HILIC chromatography for the determination of fosfomycin in plasma and urine: application to a pilot pharmacokinetic study in humans. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2015 , 105, 39-45 | 3.5 | 19 |
| 318 | Interethnic differences in pharmacokinetics of antibacterials. Clinical Pharmacokinetics, 2015, 54, 243-6 | 06.2 | 16 |
| 317 | Can we use an ex vivo continuous hemofiltration model to describe the adsorption and elimination of meropenem and piperacillin?. <i>International Journal of Artificial Organs</i> , 2015 , 38, 419-24 | 1.9 | 11 |
| 316 | Pharmacokinetic/pharmacodynamic considerations for the optimization of antimicrobial delivery in the critically ill. <i>Current Opinion in Critical Care</i> , 2015 , 21, 412-20 | 3.5 | 55 |
| 315 | A Molecular Host Response Assay to Discriminate Between Sepsis and Infection-Negative Systemic Inflammation in Critically Ill Patients: Discovery and Validation in Independent Cohorts. <i>PLoS Medicine</i> , 2015 , 12, e1001916 | 11.6 | 111 |
| 314 | The Effect of Paracetamol on Core Body Temperature in Acute Traumatic Brain Injury: A Randomised, Controlled Clinical Trial. <i>PLoS ONE</i> , 2015 , 10, e0144740 | 3.7 | 16 |
| 313 | Factors associated with vancomycin nephrotoxicity in the critically ill. <i>Anaesthesia and Intensive Care</i> , 2015 , 43, 594-9 | 1.1 | 31 |
| 312 | Are standard doses of piperacillin sufficient for critically ill patients with augmented creatinine clearance?. <i>Critical Care</i> , 2015 , 19, 28 | 10.8 | 86 |
| 311 | Subtleties in practical application of prolonged infusion of Elactam antibiotics. <i>International Journal of Antimicrobial Agents</i> , 2015 , 45, 461-3 | 14.3 | 24 |
| 310 | A multicenter study on the effect of continuous hemodiafiltration intensity on antibiotic pharmacokinetics. <i>Critical Care</i> , 2015 , 19, 84 | 10.8 | 86 |
| 309 | Quantitative bioanalytical validation of fosfomycin in human whole blood with volumetric absorptive microsampling. <i>Bioanalysis</i> , 2015 , 7, 2585-95 | 2.1 | 36 |
| 308 | To increase or decrease dosage of antimicrobials in septic patients during continuous renal replacement therapy: the eternal doubt. <i>Current Opinion in Pharmacology</i> , 2015 , 24, 68-78 | 5.1 | 21 |
| 307 | Antibiotic dose optimization in critically ill patients. <i>Medicina Intensiva (English Edition)</i> , 2015 , 39, 563-5 | 7 ð .2 | 1 |

| 306 | Pharmacokinetics of fluconazole in critically ill patients with acute kidney injury receiving sustained low-efficiency diafiltration. <i>International Journal of Antimicrobial Agents</i> , 2015 , 45, 192-5 | 14.3 | 5 |
|-------------|--|------|-----|
| 305 | Pharmacokinetics of meropenem in critically ill patients receiving continuous venovenous haemofiltration: a randomised controlled trial of continuous infusion versus intermittent bolus administration. <i>International Journal of Antimicrobial Agents</i> , 2015 , 45, 41-5 | 14.3 | 33 |
| 304 | Comparison of the accuracy and precision of pharmacokinetic equations to predict free meropenem concentrations in critically ill patients. <i>Antimicrobial Agents and Chemotherapy</i> , 2015 , 59, 1411-7 | 5.9 | 19 |
| 303 | Are interstitial fluid concentrations of meropenem equivalent to plasma concentrations in critically ill patients receiving continuous renal replacement therapy?. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 528-33 | 5.1 | 17 |
| 302 | How can we ensure effective antibiotic dosing in critically ill patients receiving different types of renal replacement therapy?. <i>Diagnostic Microbiology and Infectious Disease</i> , 2015 , 82, 92-103 | 2.9 | 54 |
| 301 | The Durban World Congress Ethics Round Table IV: health care professional end-of-life decision making. <i>Journal of Critical Care</i> , 2015 , 30, 224-30 | 4 | 20 |
| 300 | Plasma and target-site subcutaneous tissue population pharmacokinetics and dosing simulations of cefazolin in post-trauma critically ill patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2015 , 70, 1495-502 | 5.1 | 43 |
| 299 | Seated and semi-recumbent positioning of the ventilated intensive care patient - effect on gas exchange, respiratory mechanics and hemodynamics. <i>Heart and Lung: Journal of Acute and Critical Care</i> , 2014 , 43, 105-11 | 2.6 | 12 |
| 298 | Individualised antibiotic dosing for patients who are critically ill: challenges and potential solutions. <i>Lancet Infectious Diseases, The</i> , 2014 , 14, 498-509 | 25.5 | 534 |
| 297 | Variability in protein binding of teicoplanin and achievement of therapeutic drug monitoring targets in critically ill patients: lessons from the DALI Study. <i>International Journal of Antimicrobial Agents</i> , 2014 , 43, 423-30 | 14.3 | 29 |
| 296 | The influence of acute kidney injury on antimicrobial dosing in critically ill patients: are dose reductions always necessary?. <i>Diagnostic Microbiology and Infectious Disease</i> , 2014 , 79, 77-84 | 2.9 | 43 |
| 295 | A national survey of renal replacement therapy prescribing practice for acute kidney injury in Malaysian intensive care units. <i>Nephrology</i> , 2014 , 19, 507-12 | 2.2 | 10 |
| 294 | Altered pharmacokinetics of piperacillin in febrile neutropenic patients with hematological malignancy. <i>Antimicrobial Agents and Chemotherapy</i> , 2014 , 58, 3533-7 | 5.9 | 30 |
| 293 | An international, multicentre survey of Elactam antibiotic therapeutic drug monitoring practice in intensive care units. <i>Journal of Antimicrobial Chemotherapy</i> , 2014 , 69, 1416-23 | 5.1 | 144 |
| 292 | DALI: defining antibiotic levels in intensive care unit patients: are current flactam antibiotic doses sufficient for critically ill patients?. <i>Clinical Infectious Diseases</i> , 2014 , 58, 1072-83 | 11.6 | 564 |
| 291 | Plasma and cerebrospinal fluid concentrations of linezolid in neurosurgical critically ill patients with proven or suspected central nervous system infections. <i>International Journal of Antimicrobial Agents</i> , 2014 , 44, 409-15 | 14.3 | 34 |
| 2 90 | The Durban World Congress Ethics Round Table Conference Report: I. Differences between withholding and withdrawing life-sustaining treatments. <i>Journal of Critical Care</i> , 2014 , 29, 890-5 | 4 | 28 |
| 289 | Reply to Rhodes et al. <i>Clinical Infectious Diseases</i> , 2014 , 59, 907-8 | 11.6 | 2 |

| 288 | Abdominal infections in the intensive care unit: characteristics, treatment and determinants of outcome. <i>BMC Infectious Diseases</i> , 2014 , 14, 420 | 4 | 64 |
|-----|---|------------------|-----|
| 287 | Risk factors for target non-attainment during empirical treatment with Elactam antibiotics in critically ill patients. <i>Intensive Care Medicine</i> , 2014 , 40, 1340-51 | 14.5 | 112 |
| 286 | The Durban World Congress Ethics Round Table Conference Report: II. Withholding or withdrawing of treatment in elderly patients admitted to the intensive care unit. <i>Journal of Critical Care</i> , 2014 , 29, 896-901 | 4 | 29 |
| 285 | Does contemporary vancomycin dosing achieve therapeutic targets in a heterogeneous clinical cohort of critically ill patients? Data from the multinational DALI study. <i>Critical Care</i> , 2014 , 18, R99 | 10.8 | 65 |
| 284 | Pharmacokinetics of piperacillin and tazobactam in plasma and subcutaneous interstitial fluid in critically ill patients receiving continuous venovenous haemodiafiltration. <i>International Journal of Antimicrobial Agents</i> , 2014 , 43, 343-8 | 14.3 | 32 |
| 283 | Infections and use of antibiotics in patients admitted for severe acute pancreatitis: data from the EPIC II study. <i>Surgical Infections</i> , 2014 , 15, 394-8 | 2 | 19 |
| 282 | Population pharmacokinetics and dose simulation of vancomycin in critically ill patients during high-volume haemofiltration. <i>International Journal of Antimicrobial Agents</i> , 2014 , 44, 163-7 | 14.3 | 21 |
| 281 | Variable ganciclovir concentrations in a critically ill patient receiving continuous renal replacement therapy and plasma exchange?. <i>International Journal of Antimicrobial Agents</i> , 2014 , 43, 572-3 | 14.3 | 5 |
| 280 | Population pharmacokinetics and dosing simulations of cefuroxime in critically ill patients: non-standard dosing approaches are required to achieve therapeutic exposures. <i>Journal of Antimicrobial Chemotherapy</i> , 2014 , 69, 2797-803 | 5.1 | 25 |
| 279 | The Durban World Congress Ethics Round Table Conference Report: III. Withdrawing Mechanical ventilationthe approach should be individualized. <i>Journal of Critical Care</i> , 2014 , 29, 902-7 | 4 | 22 |
| 278 | The authors reply. <i>Critical Care Medicine</i> , 2014 , 42, e675-6 | 1.4 | |
| 277 | The authors reply. <i>Critical Care Medicine</i> , 2014 , 42, e602-3 | 1.4 | |
| 276 | Tracheal cuff pressure monitoring in the ICU: a literature review and survey of current practice in Queensland. <i>Anaesthesia and Intensive Care</i> , 2014 , 42, 761-70 | 1.1 | 14 |
| 275 | Select critically ill patients at risk of augmented renal clearance: experience in a Malaysian intensive care unit. <i>Anaesthesia and Intensive Care</i> , 2014 , 42, 715-22 | 1.1 | 20 |
| 274 | How do we use therapeutic drug monitoring to improve outcomes from severe infections in critically ill patients?. <i>BMC Infectious Diseases</i> , 2014 , 14, 288 | 4 | 67 |
| 273 | Low flucloxacillin concentrations in a patient with central nervous system infection: the need for plasma and cerebrospinal fluid drug monitoring in the ICU. <i>Annals of Pharmacotherapy</i> , 2014 , 48, 1380-4 | 1 ^{2.9} | 14 |
| 272 | Antimicrobial stewardship of Elactams in intensive care units. <i>Expert Review of Anti-Infective Therapy</i> , 2014 , 12, 581-95 | 5.5 | 11 |
| 271 | The effect of pathophysiology on pharmacokinetics in the critically ill patientconcepts appraised by the example of antimicrobial agents. <i>Advanced Drug Delivery Reviews</i> , 2014 , 77, 3-11 | 18.5 | 242 |

(2013-2014)

| 270 | to have high levels of antibiotic resistance compared to those with low levels. <i>BMC Infectious Diseases</i> , 2014 , 14, 513 | 4 | 12 |
|-----|--|------|-----|
| 269 | The combined effects of extracorporeal membrane oxygenation and renal replacement therapy on meropenem pharmacokinetics: a matched cohort study. <i>Critical Care</i> , 2014 , 18, 565 | 10.8 | 66 |
| 268 | Determining the mechanisms underlying augmented renal drug clearance in the critically ill: use of exogenous marker compounds. <i>Critical Care</i> , 2014 , 18, 657 | 10.8 | 50 |
| 267 | Burn resuscitationhourly urine output versus alternative endpoints: a systematic review. <i>Shock</i> , 2014 , 42, 295-306 | 3.4 | 40 |
| 266 | The impact of variation in renal replacement therapy settings on piperacillin, meropenem, and vancomycin drug clearance in the critically ill: an analysis of published literature and dosing regimens*. <i>Critical Care Medicine</i> , 2014 , 42, 1640-50 | 1.4 | 81 |
| 265 | Vancomycin-associated nephrotoxicity in the critically ill: a retrospective multivariate regression analysis*. <i>Critical Care Medicine</i> , 2014 , 42, 2527-36 | 1.4 | 71 |
| 264 | Augmented renal clearance in the ICU: results of a multicenter observational study of renal function in critically ill patients with normal plasma creatinine concentrations*. <i>Critical Care Medicine</i> , 2014 , 42, 520-7 | 1.4 | 181 |
| 263 | A new marker of sepsis post burn injury?*. Critical Care Medicine, 2014, 42, 2029-36 | 1.4 | 31 |
| 262 | Doripenem population pharmacokinetics and dosing requirements for critically ill patients receiving continuous venovenous haemodiafiltration. <i>Journal of Antimicrobial Chemotherapy</i> , 2014 , 69, 2508-16 | 5.1 | 25 |
| 261 | Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. Lancet Respiratory Medicine, the, 2014 , 2, 380-6 | 35.1 | 566 |
| 260 | Principles of antibiotic use 2014 , 738-742.e1 | | |
| 259 | Extended versus bolus infusion of meropenem and piperacillin: a pharmacokinetic analysis. <i>Minerva Anestesiologica</i> , 2014 , 80, 1302-9 | 1.9 | 23 |
| 258 | What is the relevance of fosfomycin pharmacokinetics in the treatment of serious infections in critically ill patients? A systematic review. <i>International Journal of Antimicrobial Agents</i> , 2013 , 42, 289-93 | 14.3 | 54 |
| 257 | Population pharmacokinetics and dosing simulations of amoxicillin/clavulanic acid in critically ill patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2013 , 68, 2600-8 | 5.1 | 36 |
| 256 | Protein binding of Elactam antibiotics in critically ill patients: can we successfully predict unbound concentrations?. <i>Antimicrobial Agents and Chemotherapy</i> , 2013 , 57, 6165-70 | 5.9 | 137 |
| 255 | Reply to Soman et al. <i>Clinical Infectious Diseases</i> , 2013 , 57, 323-4 | 11.6 | |
| 254 | A comparison of CKD-EPI estimated glomerular filtration rate and measured creatinine clearance in recently admitted critically ill patients with normal plasma creatinine concentrations. <i>BMC Nephrology</i> , 2013 , 14, 250 | 2.7 | 35 |
| 253 | Clinical implications of antibiotic pharmacokinetic principles in the critically ill. <i>Intensive Care Medicine</i> , 2013 , 39, 2070-82 | 14.5 | 150 |

| 252 | Ampicillin/sulbactam: its potential use in treating infections in critically ill patients. <i>International Journal of Antimicrobial Agents</i> , 2013 , 42, 384-9 | 14.3 | 41 |
|-----|---|----------------------|-----|
| 251 | The clinical relevance of plasma protein binding changes. Clinical Pharmacokinetics, 2013, 52, 1-8 | 6.2 | 173 |
| 250 | Continuous infusion of beta-lactam antibiotics in severe sepsis: a multicenter double-blind, randomized controlled trial. <i>Clinical Infectious Diseases</i> , 2013 , 56, 236-44 | 11.6 | 250 |
| 249 | Lactam therapeutic drug monitoring in the critically ill: optimising drug exposure in patients with fluctuating renal function and hypoalbuminaemia. <i>International Journal of Antimicrobial Agents</i> , 2013 , 41, 162-6 | 14.3 | 49 |
| 248 | Phenytoin loading doses in adult critical care patients: does current practice achieve adequate drug levels?. <i>Anaesthesia and Intensive Care</i> , 2013 , 41, 602-9 | 1.1 | 4 |
| 247 | Pharmacokinetics of meropenem and piperacillin in critically ill patients with indwelling surgical drains. <i>International Journal of Antimicrobial Agents</i> , 2013 , 42, 90-3 | 14.3 | 17 |
| 246 | Can population pharmacokinetic modelling guide vancomycin dosing during continuous renal replacement therapy in critically ill patients?. <i>International Journal of Antimicrobial Agents</i> , 2013 , 41, 564 | 1 <mark>14</mark> .3 | 35 |
| 245 | Augmented renal clearance in septic and traumatized patients with normal plasma creatinine concentrations: identifying at-risk patients. <i>Critical Care</i> , 2013 , 17, R35 | 10.8 | 108 |
| 244 | Meropenem and piperacillin/tazobactam prescribing in critically ill patients: does augmented renal clearance affect pharmacokinetic/pharmacodynamic target attainment when extended infusions are used?. Critical Care, 2013, 17, R84 | 10.8 | 133 |
| 243 | Molecular analysis of the Acinetobacter baumannii biofilm-associated protein. <i>Applied and Environmental Microbiology</i> , 2013 , 79, 6535-43 | 4.8 | 49 |
| 242 | B-type natriuretic Peptide: not just a heart throb. A & A Case Reports, 2013, 1, 59-61 | | |
| 241 | A Protocol for the Pharmacokinetics of Enteric Coated Mycophenolate Sodium in Lupus Nephritis (POEMSLUN): an open-label, randomised controlled trial. <i>BMJ Open</i> , 2013 , 3, | 3 | 2 |
| 240 | How should we dose antibiotics for pneumonia in the ICU?. <i>Current Opinion in Infectious Diseases</i> , 2013 , 26, 189-95 | 5.4 | 8 |
| 239 | Toward improved surveillance: the impact of ventilator-associated complications on length of stay and antibiotic use in patients in intensive care units. <i>Clinical Infectious Diseases</i> , 2013 , 56, 471-7 | 11.6 | 113 |
| 238 | Antibacterial therapeutic drug monitoring in cerebrospinal fluid: difficulty in achieving adequate drug concentrations. <i>Journal of Neurosurgery</i> , 2013 , 118, 297-301 | 3.2 | 31 |
| 237 | Defining Antibiotic Dosing in Lung Infections. <i>Clinical Pulmonary Medicine</i> , 2013 , 20, 121-128 | 0.3 | 14 |
| 236 | A new regimen for continuous infusion of vancomycin during continuous renal replacement therapy. <i>Journal of Antimicrobial Chemotherapy</i> , 2013 , 68, 2859-65 | 5.1 | 42 |
| 235 | Optimal doripenem dosing simulations in critically ill nosocomial pneumonia patients with obesity, augmented renal clearance, and decreased bacterial susceptibility. <i>Critical Care Medicine</i> , 2013 , 41, 489- | 9 ₅ 4 | 65 |

| 234 | Identification of novel vaccine candidates against multidrug-resistant Acinetobacter baumannii. <i>PLoS ONE</i> , 2013 , 8, e77631 | 3.7 | 56 |
|-----|---|--------------------|------|
| 233 | A protocol for a multicentre randomised controlled trial of continuous beta-lactam infusion compared with intermittent beta-lactam dosing in critically ill patients with severe sepsis: the BLING II study. Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care | 2.8 | 3 |
| 232 | Review article: Part two: Goal-directed resuscitationwhich goals? Perfusion targets. <i>EMA - Emergency Medicine Australasia</i> , 2012 , 24, 127-35 | 1.5 | 12 |
| 231 | Review article: Part one: Goal-directed resuscitationwhich goals? Haemodynamic targets. <i>EMA - Emergency Medicine Australasia</i> , 2012 , 24, 14-22 | 1.5 | 12 |
| 230 | Elactam pharmacokinetics and pharmacodynamics in critically ill patients and strategies for dose optimization: a structured review. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2012 , 39, 489- | -96 | 59 |
| 229 | Hydroxyethyl starch or saline for fluid resuscitation in intensive care. <i>New England Journal of Medicine</i> , 2012 , 367, 1901-11 | 59.2 | 1127 |
| 228 | Characteristics and determinants of outcome of hospital-acquired bloodstream infections in intensive care units: the EUROBACT International Cohort Study. <i>Intensive Care Medicine</i> , 2012 , 38, 1930- | -4 4 ·5 | 226 |
| 227 | A limited set of molecular biomarkers may provide superior diagnostic outcomes to procalcitonin in sepsis. <i>Critical Care</i> , 2012 , 16, | 10.8 | 1 |
| 226 | Characteristics of bloodstream infections in burn patients: An 11-year retrospective study. <i>Burns</i> , 2012 , 38, 685-90 | 2.3 | 40 |
| 225 | DALI: Defining Antibiotic Levels in Intensive care unit patients: a multi-centre point of prevalence study to determine whether contemporary antibiotic dosing for critically ill patients is therapeutic. <i>BMC Infectious Diseases</i> , 2012 , 12, 152 | 4 | 34 |
| 224 | Using pharmacokinetics and pharmacodynamics to optimise dosing of antifungal agents in critically ill patients: a systematic review. <i>International Journal of Antimicrobial Agents</i> , 2012 , 39, 1-10 | 14.3 | 37 |
| 223 | Improving vancomycin prescription in critical illness through a drug use evaluation process: a weight-based dosing intervention study. <i>International Journal of Antimicrobial Agents</i> , 2012 , 39, 69-72 | 14.3 | 35 |
| 222 | How to optimise antimicrobial prescriptions in the Intensive Care Unit: principles of individualised dosing using pharmacokinetics and pharmacodynamics. <i>International Journal of Antimicrobial Agents</i> , 2012 , 39, 187-92 | 14.3 | 45 |
| 221 | A method for determining the free (unbound) concentration of ten beta-lactam antibiotics in human plasma using high performance liquid chromatography with ultraviolet detection. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012 , 907, 178-84 | 3.2 | 84 |
| 220 | Pharmacokinetics of beta-lactam antibiotics in patients with intra-abdominal disease: a structured review. <i>Surgical Infections</i> , 2012 , 13, 9-17 | 2 | 16 |
| 219 | Editorial commentary: Closing the loopa colistin clinical study to confirm dosing recommendations from PK/PD modeling. <i>Clinical Infectious Diseases</i> , 2012 , 54, 1727-9 | 11.6 | 18 |
| 218 | Continuous beta-lactam infusion in critically ill patients: the clinical evidence. <i>Annals of Intensive Care</i> , 2012 , 2, 37 | 8.9 | 66 |
| 217 | Does Beta-lactam Pharmacokinetic Variability in Critically Ill Patients Justify Therapeutic Drug Monitoring? A Systematic Review. <i>Annals of Intensive Care</i> , 2012 , 2, 35 | 8.9 | 119 |

| 216 | Blood culture collection in patients with acute kidney injury receiving renal replacement therapy: an observational study. <i>Anaesthesia and Intensive Care</i> , 2012 , 40, 813-9 | 1.1 | 3 |
|-----|--|-------------|-----------|
| 215 | A comparison of pulse contour wave analysis and ultrasonic cardiac output monitoring in the critically ill. <i>Anaesthesia and Intensive Care</i> , 2012 , 40, 631-7 | 1.1 | 8 |
| 214 | Importance of High Creatinine Clearance for Antibacterial Treatment in Sepsis 2012, 171-197 | | |
| 213 | Improving antibiotic dosing in special situations in the ICU: burns, renal replacement therapy and extracorporeal membrane oxygenation. <i>Current Opinion in Critical Care</i> , 2012 , 18, 460-71 | 3.5 | 68 |
| 212 | Pharmacokinetics of intraperitoneal gentamicin in peritoneal dialysis patients with peritonitis (GIPD study). <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012 , 7, 1249-56 | 6.9 | 16 |
| 211 | Therapeutic drug monitoring of beta-lactam antibiotics in burns patientsa one-year prospective study. <i>Therapeutic Drug Monitoring</i> , 2012 , 34, 160-4 | 3.2 | 46 |
| 210 | Variability of antibiotic concentrations in critically ill patients receiving continuous renal replacement therapy: a multicentre pharmacokinetic study. <i>Critical Care Medicine</i> , 2012 , 40, 1523-8 | 1.4 | 151 |
| 209 | Subtherapeutic initial flactam concentrations in select critically ill patients: association between augmented renal clearance and low trough drug concentrations. <i>Chest</i> , 2012 , 142, 30-39 | 5.3 | 277 |
| 208 | The ecology of antibiotic use in the ICU: homogeneous prescribing of cefepime but not tazocin selects for antibiotic resistant infection. <i>PLoS ONE</i> , 2012 , 7, e38719 | 3.7 | 19 |
| 207 | Continuous infusion vs. bolus dosing: implications for beta-lactam antibiotics. <i>Minerva Anestesiologica</i> , 2012 , 78, 94-104 | 1.9 | 26 |
| 206 | Therapeutic drug monitoring when using cefepime in continuous renal replacement therapy: seizures associated with cefepime. <i>Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine</i> , 2012 , 14, 312-5 | 2.8 | 13 |
| 205 | Ganciclovir pharmacokinetics and suggested dosing in continuous venovenous haemodiafiltration. <i>International Journal of Antimicrobial Agents</i> , 2011 , 37, 90-2 | 14.3 | 6 |
| | | | |
| 204 | Increased mortality associated with methicillin-resistant Staphylococcus aureus (MRSA) infection in the intensive care unit: results from the EPIC II study. <i>International Journal of Antimicrobial Agents</i> , 2011 , 38, 331-5 | 14.3 | 109 |
| 204 | the intensive care unit: results from the EPIC II study. International Journal of Antimicrobial Agents, | 14.3 4.5 | 109 96 |
| · | the intensive care unit: results from the EPIC II study. <i>International Journal of Antimicrobial Agents</i> , 2011 , 38, 331-5 Antimicrobial pharmacokinetic and pharmacodynamic issues in the critically ill with severe sepsis | | |
| 203 | the intensive care unit: results from the EPIC II study. <i>International Journal of Antimicrobial Agents</i> , 2011 , 38, 331-5 Antimicrobial pharmacokinetic and pharmacodynamic issues in the critically ill with severe sepsis and septic shock. <i>Critical Care Clinics</i> , 2011 , 27, 19-34 The effects of hypoalbuminaemia on optimizing antibacterial dosing in critically ill patients. <i>Clinical</i> | 4.5 | 96 |
| 203 | the intensive care unit: results from the EPIC II study. <i>International Journal of Antimicrobial Agents</i> , 2011 , 38, 331-5 Antimicrobial pharmacokinetic and pharmacodynamic issues in the critically ill with severe sepsis and septic shock. <i>Critical Care Clinics</i> , 2011 , 27, 19-34 The effects of hypoalbuminaemia on optimizing antibacterial dosing in critically ill patients. <i>Clinical Pharmacokinetics</i> , 2011 , 50, 99-110 Pharmacokinetic evaluation of fluconazole in critically ill patients. <i>Expert Opinion on Drug</i> | 4.5 | 96 238 |

| 198 | Experience with high frequency oscillation ventilation during the 2009 H1N1 influenza pandemic in Australia and New Zealand. <i>Anaesthesia and Intensive Care</i> , 2011 , 39, 837-46 | 1.1 | 4 |
|-----|---|-------|-----|
| 197 | Sustained low efficiency dialysis allows rational renal replacement therapy, but does it allow rational drug dosing?. <i>Critical Care Medicine</i> , 2011 , 39, 602-3 | 1.4 | 18 |
| 196 | Determining authorship in multicenter trials: a systematic review. <i>Acta Anaesthesiologica Scandinavica</i> , 2011 , 55, 1037-43 | 1.9 | 11 |
| 195 | Pitfalls of using estimations of glomerular filtration rate in an intensive care population. <i>Internal Medicine Journal</i> , 2011 , 41, 537-43 | 1.6 | 50 |
| 194 | Prior statin use is not associated with improved outcome in emergency patients admitted with infection: a prospective observational study. <i>Academic Emergency Medicine</i> , 2011 , 18, 127-34 | 3.4 | 7 |
| 193 | Implications of augmented renal clearance in critically ill patients. <i>Nature Reviews Nephrology</i> , 2011 , 7, 539-43 | 14.9 | 120 |
| 192 | The role of surveillance cultures in the prediction of susceptibility patterns of Gram-negative bacilli in the intensive care unit. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 2011 , 30, 739 | 9-543 | 12 |
| 191 | The Crystalloid versus Hydroxyethyl Starch Trial: protocol for a multi-centre randomised controlled trial of fluid resuscitation with 6% hydroxyethyl starch (130/0.4) compared to 0.9% sodium chloride (saline) in intensive care patients on mortality. <i>Intensive Care Medicine</i> , 2011 , 37, 816-23 | 14.5 | 29 |
| 190 | Antibiotic dosing in the lat riskLcritically ill patient: Linking pathophysiology with pharmacokinetics/pharmacodynamics in sepsis and trauma patients. <i>BMC Anesthesiology</i> , 2011 , 11, 3 | 2.4 | 33 |
| 189 | The pharmacokinetics of cefazolin in patients undergoing elective & semi-elective abdominal aortic aneurysm open repair surgery. <i>BMC Anesthesiology</i> , 2011 , 11, 5 | 2.4 | 6 |
| 188 | A prospective registry of emergency department patients admitted with infection. <i>BMC Infectious Diseases</i> , 2011 , 11, 27 | 4 | 6 |
| 187 | A comparison of estimates of glomerular filtration in critically ill patients with augmented renal clearance. <i>Critical Care</i> , 2011 , 15, R139 | 10.8 | 138 |
| 186 | Development and validation of a novel molecular biomarker diagnostic test for the early detection of sepsis. <i>Critical Care</i> , 2011 , 15, R149 | 10.8 | 101 |
| 185 | Monte Carlo simulations: maximizing antibiotic pharmacokinetic data to optimize clinical practice for critically ill patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2011 , 66, 227-31 | 5.1 | 104 |
| 184 | Physiological changes in the critically ill patient with sepsis. <i>Current Pharmaceutical Biotechnology</i> , 2011 , 12, 1991-5 | 2.6 | 22 |
| 183 | ARCaugmented renal clearance. Current Pharmaceutical Biotechnology, 2011 , 12, 2020-9 | 2.6 | 27 |
| 182 | Remote care by telemedicine in the ICU: many models of care can be effective. <i>Current Opinion in Critical Care</i> , 2011 , 17, 634-40 | 3.5 | 16 |
| 181 | Plasma and tissue pharmacokinetics of cefazolin in patients undergoing elective and semielective abdominal aortic aneurysm open repair surgery. <i>Antimicrobial Agents and Chemotherapy</i> , 2011 , 55, 5238 | 3-42 | 55 |

| 180 | Antibiotic dosing in multiple organ dysfunction syndrome. <i>Chest</i> , 2011 , 139, 1210-1220 | 5.3 | 70 |
|-----|---|------|-----|
| 179 | Effect of intravenous GLutamine supplementation IN Trauma patients receiving enteral nutrition study protocol (GLINT Study): a prospective, blinded, randomised, placebo-controlled clinical trial. <i>BMJ Open</i> , 2011 , 1, e000334 | 3 | 9 |
| 178 | Population pharmacokinetics of fluconazole in critically ill patients receiving continuous venovenous hemodiafiltration: using Monte Carlo simulations to predict doses for specified pharmacodynamic targets. <i>Antimicrobial Agents and Chemotherapy</i> , 2011 , 55, 5868-73 | 5.9 | 38 |
| 177 | Vancomycin dosing in critically ill patients: robust methods for improved continuous-infusion regimens. <i>Antimicrobial Agents and Chemotherapy</i> , 2011 , 55, 2704-9 | 5.9 | 159 |
| 176 | Candida Pneumonia in Patients with Hematological Neoplasia 2011 , 349-356 | | 1 |
| 175 | Can learning to sustain life be BASIC? Teaching for the initial management of the critically ill in Australia and New Zealand. <i>Anaesthesia and Intensive Care</i> , 2010 , 38, 1043-51 | 1.1 | 3 |
| 174 | Flucloxacillin dosing in critically ill patients with hypoalbuminaemia: special emphasis on unbound pharmacokinetics. <i>Journal of Antimicrobial Chemotherapy</i> , 2010 , 65, 1771-8 | 5.1 | 83 |
| 173 | Meropenem dosing in critically ill patients with sepsis receiving high-volume continuous venovenous hemofiltration. <i>Antimicrobial Agents and Chemotherapy</i> , 2010 , 54, 2974-8 | 5.9 | 56 |
| 172 | Antibiotic use and impact on outcome from bacteraemic critical illness: the BActeraemia Study in Intensive Care (BASIC). <i>Journal of Antimicrobial Chemotherapy</i> , 2010 , 65, 1276-85 | 5.1 | 67 |
| 171 | Using population pharmacokinetics to determine gentamicin dosing during extended daily diafiltration in critically ill patients with acute kidney injury. <i>Antimicrobial Agents and Chemotherapy</i> , 2010 , 54, 3635-40 | 5.9 | 53 |
| 170 | Optimal Antibiotic Therapy in the Management of the Lung of the Critically Ill. <i>Current Respiratory Medicine Reviews</i> , 2010 , 6, 253-263 | 0.3 | |
| 169 | Respiratory Burns: A Clinical Review. Current Respiratory Medicine Reviews, 2010, 6, 285-291 | 0.3 | |
| 168 | Control of an outbreak of carbapenem-resistant Acinetobacter baumannii in Australia after introduction of environmental cleaning with a commercial oxidizing disinfectant. <i>Infection Control and Hospital Epidemiology</i> , 2010 , 31, 418-20 | 2 | 34 |
| 167 | First-dose and steady-state population pharmacokinetics and pharmacodynamics of piperacillin by continuous or intermittent dosing in critically ill patients with sepsis. <i>International Journal of Antimicrobial Agents</i> , 2010 , 35, 156-63 | 14.3 | 133 |
| 166 | Augmented renal clearance in the Intensive Care Unit: an illustrative case series. <i>International Journal of Antimicrobial Agents</i> , 2010 , 35, 606-8 | 14.3 | 69 |
| 165 | Therapeutic drug monitoring of beta-lactams in critically ill patients: proof of concept. <i>International Journal of Antimicrobial Agents</i> , 2010 , 36, 332-9 | 14.3 | 243 |
| 164 | Antimicrobial chemotherapy and lung microdialysis: a review. <i>International Journal of Antimicrobial Agents</i> , 2010 , 36, 491-500 | 14.3 | 33 |
| 163 | Management of invasive candidiasis in the intensive care unit. <i>Drugs</i> , 2010 , 70, 823-39 | 12.1 | 21 |

(2010-2010)

| 162 | Augmented renal clearance: implications for antibacterial dosing in the critically ill. <i>Clinical Pharmacokinetics</i> , 2010 , 49, 1-16 | 6.2 | 245 |
|-----|---|------|-----|
| 161 | Augmented renal clearance in traumatic brain injury. Critical Care, 2010, 14, P521 | 10.8 | 2 |
| 160 | A novel molecular biomarker diagnostic for the early detection of sepsis. Critical Care, 2010, 14, P9 | 10.8 | 1 |
| 159 | Pharmacokinetic evaluation of piperacillin-tazobactam. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2010 , 6, 1017-31 | 5.5 | 50 |
| 158 | Principles of antibacterial dosing in continuous renal replacement therapy. <i>Blood Purification</i> , 2010 , 30, 195-212 | 3.1 | 49 |
| 157 | Preventive Strategies for Ventilator Associated Pneumonia. <i>Current Respiratory Medicine Reviews</i> , 2010 , 6, 238-252 | 0.3 | 3 |
| 156 | Optimizing Antibiotic Use in the Intensive Care Unit. Clinical Pulmonary Medicine, 2010, 17, 162-169 | 0.3 | Ο |
| 155 | Prophylaxis, empirical and preemptive treatment of invasive candidiasis. <i>Current Opinion in Critical Care</i> , 2010 , 16, 470-4 | 3.5 | 26 |
| 154 | Pharmacokinetics and pharmacodynamics in critically ill patients. <i>Current Opinion in Anaesthesiology</i> , 2010 , 23, 472-8 | 2.9 | 37 |
| 153 | Do burn patients cost more? The intensive care unit costs of burn patients compared with controls matched for length of stay and acuity. <i>Journal of Burn Care and Research</i> , 2010 , 31, 598-602 | 0.8 | 21 |
| 152 | Augmented creatinine clearance in traumatic brain injury. Anesthesia and Analgesia, 2010, 111, 1505-10 | 3.9 | 85 |
| 151 | Assessment of peripheral arterial catheters as a source of sepsis in the critically ill: a narrative review. <i>Journal of Hospital Infection</i> , 2010 , 75, 12-8 | 6.9 | 22 |
| 150 | Analysis of 12 beta-lactam antibiotics in human plasma by HPLC with ultraviolet detection. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2010 , 878, 2039-43 | 3.2 | 144 |
| 149 | Pitfalls of using estimations of glomerular filtration rate in an intensive care population. <i>Internal Medicine Journal</i> , 2010 , | 1.6 | 5 |
| 148 | Augmented Renal Clearance: Unraveling the Mystery of Elevated Antibiotic Clearance. <i>Yearbook of Intensive Care and Emergency Medicine</i> , 2010 , 495-506 | | |
| 147 | Augmented Renal Clearance: Unraveling the Mystery of Elevated Antibiotic Clearance 2010 , 495-506 | | O |
| 146 | A detailed feeding algorithm improves delivery of nutrition support in an intensive care unit. Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine, 2010, 12, 149-55 | 2.8 | 2 |
| 145 | A survey of antibiotic prescribing practices in Australian and New Zealand intensive care units. Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine, 2010, 12, 162-70 | 2.8 | 7 |

| 144 | Predictors and outcome associated with an Enterococcus positive isolate during intensive care unit admission. <i>Anaesthesia and Intensive Care</i> , 2009 , 37, 976-82 | 1.1 | 2 |
|-----|--|------|------|
| 143 | Airway obstruction due to a pre-vertebral haematoma following difficult central line insertionimplications for ultrasound guidance and review of the literature. <i>Anaesthesia and Intensive Care</i> , 2009 , 37, 309-13 | 1.1 | 8 |
| 142 | Tissue accumulation of cephalothin in burns: a comparative study by microdialysis of subcutaneous interstitial fluid cephalothin concentrations in burn patients and healthy volunteers. <i>Antimicrobial Agents and Chemotherapy</i> , 2009 , 53, 210-5 | 5.9 | 11 |
| 141 | Unbound cephalothin pharmacokinetics in adult burn patients are related to the elapsed time after injury. <i>Antimicrobial Agents and Chemotherapy</i> , 2009 , 53, 5303-5 | 5.9 | 7 |
| 140 | Assessment of clinical risk predictive rules for invasive candidiasis in a prospective multicentre cohort of ICU patients. <i>Intensive Care Medicine</i> , 2009 , 35, 2141-5 | 14.5 | 39 |
| 139 | Optimising intraperitoneal gentamicin dosing in peritoneal dialysis patients with peritonitis (GIPD) study. <i>BMC Nephrology</i> , 2009 , 10, 42 | 2.7 | 8 |
| 138 | Point of care measurement of plasma creatinine in critically ill patients with acute kidney injury. <i>Anaesthesia</i> , 2009 , 64, 403-7 | 6.6 | 30 |
| 137 | Meropenem dosing in critically ill patients with sepsis and without renal dysfunction: intermittent bolus versus continuous administration? Monte Carlo dosing simulations and subcutaneous tissue distribution. <i>Journal of Antimicrobial Chemotherapy</i> , 2009 , 64, 142-50 | 5.1 | 238 |
| 136 | Temporal trends, risk factors and outcomes in albicans and non-albicans candidaemia: an international epidemiological study in four multidisciplinary intensive care units. <i>International Journal of Antimicrobial Agents</i> , 2009 , 33, 554.e1-7 | 14.3 | 47 |
| 135 | Clearance of intravenous 5-fluorocytosine during continuous venovenous haemodiafiltration in a patient with hepatosplenic candidiasis. <i>International Journal of Antimicrobial Agents</i> , 2009 , 34, 383-4 | 14.3 | 5 |
| 134 | A systematic review of antibiotic dosing regimens for septic patients receiving continuous renal replacement therapy: do current studies supply sufficient data?. <i>Journal of Antimicrobial Chemotherapy</i> , 2009 , 64, 929-37 | 5.1 | 68 |
| 133 | Management of meningitis due to antibiotic-resistant Acinetobacter species. <i>Lancet Infectious Diseases, The</i> , 2009 , 9, 245-55 | 25.5 | 152 |
| 132 | International study of the prevalence and outcomes of infection in intensive care units. <i>JAMA - Journal of the American Medical Association</i> , 2009 , 302, 2323-9 | 27.4 | 2023 |
| 131 | Pharmacokinetic issues for antibiotics in the critically ill patient. <i>Critical Care Medicine</i> , 2009 , 37, 840-51; quiz 859 | 1.4 | 598 |
| 130 | Principles of antibacterial dosing in continuous renal replacement therapy. <i>Critical Care Medicine</i> , 2009 , 37, 2268-82 | 1.4 | 280 |
| 129 | Dose matters: Dose of antibiotics in the critically ill patient depends on the dose of renal replacement therapy. <i>Critical Care Medicine</i> , 2009 , 37, 2491-2492 | 1.4 | |
| 128 | Midazolam Metabolism: Implications for Individualised Dosing?. <i>Journal of Pharmacy Practice and Research</i> , 2009 , 39, 198-201 | 0.7 | 3 |
| 127 | A systematic review on clinical benefits of continuous administration of beta-lactam antibiotics. <i>Critical Care Medicine</i> , 2009 , 37, 2071-8 | 1.4 | 208 |

(2007-2009)

| 126 | Tissue Distribution of Beta-Lactam Antibiotics: Continuous versus Bolus Dosing. <i>Journal of Pharmacy Practice and Research</i> , 2009 , 39, 219-222 | 0.7 | 5 |
|-----|---|------|-----|
| 125 | Respiratory Complications in Burns. Clinical Pulmonary Medicine, 2009, 16, 132-138 | 0.3 | 8 |
| 124 | Piperacillin penetration into tissue of critically ill patients with sepsisbolus versus continuous administration?. <i>Critical Care Medicine</i> , 2009 , 37, 926-33 | 1.4 | 136 |
| 123 | The microcirculation, regional blood flow and tissue oxygenation: will new technologies drive new resuscitation goals?. <i>Anaesthesia and Intensive Care</i> , 2009 , 37, 700-2 | 1.1 | 1 |
| 122 | Principles of Pharmacology in the Critically Ill 2009 , 31-36 | | |
| 121 | A new paradigm for treating infections: "go hard and go home". <i>Critical Care and Resuscitation:</i> Journal of the Australasian Academy of Critical Care Medicine, 2009 , 11, 276-81 | 2.8 | 7 |
| 120 | Lessons learnt in the pharmacokinetic analysis of the effect of haemoperfusion for acute overdose with sustained-release diltiazem. <i>Anaesthesia</i> , 2008 , 63, 714-8 | 6.6 | 10 |
| 119 | A novel way to investigate the effects of plasma exchange on antibiotic levels: use of microdialysis. <i>International Journal of Antimicrobial Agents</i> , 2008 , 31, 240-4 | 14.3 | 9 |
| 118 | Increased fluid resuscitation can lead to adverse outcomes in major-burn injured patients, but low mortality is achievable. <i>Burns</i> , 2008 , 34, 1090-7 | 2.3 | 77 |
| 117 | Antibiotic resistancewhatIs dosing got to do with it?. Critical Care Medicine, 2008, 36, 2433-40 | 1.4 | 239 |
| 116 | Better outcomes through continuous infusion of time-dependent antibiotics to critically ill patients?. <i>Current Opinion in Critical Care</i> , 2008 , 14, 390-6 | 3.5 | 75 |
| 115 | Continuous Infusion of Time-dependent Antibiotics. Clinical Pulmonary Medicine, 2008, 15, 167-172 | 0.3 | 1 |
| 114 | Does severe non-infectious SIRS differ from severe sepsis? Results from a multi-centre Australian and New Zealand intensive care unit study. <i>Intensive Care Medicine</i> , 2008 , 34, 1654-61 | 14.5 | 54 |
| 113 | A comparison of epinephrine and norepinephrine in critically ill patients. <i>Intensive Care Medicine</i> , 2008 , 34, 2226-34 | 14.5 | 216 |
| 112 | Dose Adjustment and Pharmacodynamic Considerations for Antibiotics in Severe Sepsis and Septic Shock 2008 , 97-136 | | 2 |
| 111 | Prehospital Care 2008 , 307-319 | | 1 |
| 110 | Cefepime versus ceftazidime: considerations for empirical use in critically ill patients. <i>International Journal of Antimicrobial Agents</i> , 2007 , 29, 117-28 | 14.3 | 19 |
| 109 | Continuous infusion of beta-lactam antibiotics in severe infections: a review of its role. <i>International Journal of Antimicrobial Agents</i> , 2007 , 30, 11-8 | 14.3 | 132 |

| 108 | Acute respiratory distress in a bleomycin primed patient: a new use for nitric oxide. <i>Anaesthesia and Intensive Care</i> , 2007 , 35, 86-90 | 1.1 | 1 |
|-----|--|------|-----|
| 107 | Autonomic dysfunction in tetanus - what lessons can be learnt with specific reference to alpha-2 agonists?. <i>Anaesthesia</i> , 2007 , 62, 1066-70 | 6.6 | 21 |
| 106 | Population pharmacokinetics and pharmacodynamics of cefpirome in critically ill patients against Gram-negative bacteria. <i>Intensive Care Medicine</i> , 2007 , 33, 781-788 | 14.5 | 32 |
| 105 | Is continuous infusion ceftriaxone better than once-a-day dosing in intensive care? A randomized controlled pilot study. <i>Journal of Antimicrobial Chemotherapy</i> , 2007 , 59, 285-91 | 5.1 | 91 |
| 104 | The entrapped central venous catheter. British Journal of Anaesthesia, 2007, 98, 89-92 | 5.4 | 9 |
| 103 | Optimizing use of beta-lactam antibiotics in the critically ill. <i>Seminars in Respiratory and Critical Care Medicine</i> , 2007 , 28, 579-85 | 3.9 | 16 |
| 102 | Inadequate antimicrobial prophylaxis during surgery: a study of beta-lactam levels during burn debridement. <i>Journal of Antimicrobial Chemotherapy</i> , 2007 , 60, 166-9 | 5.1 | 15 |
| 101 | Entrapped central venous catheter. British Journal of Anaesthesia, 2007, 98, 695; author reply 695 | 5.4 | |
| 100 | Returning to the pre-antibiotic era in the critically ill: the XDR problem. <i>Critical Care Medicine</i> , 2007 , 35, 1789-91 | 1.4 | 28 |
| 99 | Resistance in the intensive care unit: whose problem is it and how can intensivists help?. <i>Critical Care Medicine</i> , 2007 , 35, 299-301 | 1.4 | 4 |
| 98 | Lateral positioning of ventilated intensive care patients: a study of oxygenation, respiratory mechanics, hemodynamics, and adverse events. <i>Heart and Lung: Journal of Acute and Critical Care</i> , 2007 , 36, 277-86 | 2.6 | 32 |
| 97 | Dose Adjustment and Pharmacokinetics of Antibiotics in Severe Sepsis and Septic Shock 2007 , 122-146 | | |
| 96 | Prescription of Antimicrobial Agents in Patients Undergoing Continuous Renal Replacement Therapy 2007 , 147-158 | | |
| 95 | The implications of enterococci for the intensive care unit. <i>Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine</i> , 2007 , 9, 69-75 | 2.8 | 9 |
| 94 | Pharmacokinetic-pharmacodynamic rationale for cefepime dosing regimens in intensive care units. Journal of Antimicrobial Chemotherapy, 2006 , 58, 987-93 | 5.1 | 80 |
| 93 | Positioning practices for ventilated intensive care patients: current practice, indications and contraindications. <i>Australian Critical Care</i> , 2006 , 19, 122-6, 128, 130-2 | 2.9 | 22 |
| 92 | Antibacterial dosing in intensive care: pharmacokinetics, degree of disease and pharmacodynamics of sepsis. <i>Clinical Pharmacokinetics</i> , 2006 , 45, 755-73 | 6.2 | 205 |
| 91 | Effect of drug concentration on adsorption of levofloxacin by polyacrylonitrile haemofilters. <i>International Journal of Antimicrobial Agents</i> , 2006 , 28, 147-50 | 14.3 | 27 |

(2004-2006)

| 90 | Manual hyperinflation causes norepinephrine release. <i>Heart and Lung: Journal of Acute and Critical Care</i> , 2006 , 35, 262-8 | 2.6 | 6 |
|----------------------|---|---------------------------------|-----------------------|
| 89 | Estimate of the number of patients eligible for treatment with drotrecogin alfa (activated) based on differing international indications: post-hoc analysis of an inception cohort study in Australia and New Zealand. <i>Anaesthesia and Intensive Care</i> , 2006 , 34, 184-90 | 1.1 | 1 |
| 88 | Making the wrong choice: Consequences for patients with infections due to methicillin-resistant Staphylococcus aureus. <i>Critical Care Medicine</i> , 2006 , 34, 2239-41 | 1.4 | 4 |
| 87 | Low mannose-binding lectin function is associated with sepsis in adult patients. <i>FEMS Immunology and Medical Microbiology</i> , 2006 , 48, 274-82 | | 43 |
| 86 | The use of prophylactic fluconazole in immunocompetent high-risk surgical patients: a meta-analysis. <i>Critical Care</i> , 2005 , 9, R710-7 | 10.8 | 34 |
| 85 | Fluid shifts have no influence on ciprofloxacin pharmacokinetics in intensive care patients with intra-abdominal sepsis. <i>International Journal of Antimicrobial Agents</i> , 2005 , 26, 50-5 | 14.3 | 40 |
| 84 | Is the end-game penetration (of the airway)?. Critical Care Medicine, 2005, 33, 1654-5 | 1.4 | 25 |
| 83 | Disease risk and mortality prediction in intensive care patients with pneumonia. Australian and New Zealand practice in intensive care (ANZPIC II). <i>Anaesthesia and Intensive Care</i> , 2005 , 33, 101-11 | 1.1 | 17 |
| 82 | The costs of caring for patients in a tertiary referral Australian Intensive Care Unit. <i>Anaesthesia and Intensive Care</i> , 2005 , 33, 477-82 | 1.1 | 40 |
| | | | |
| 81 | Antibiotic resistance in the intensive care unit: a primer in bacteriology. <i>Anaesthesia and Intensive Care</i> , 2005 , 33, 188-95 | 1.1 | |
| 80 | | 1.1 | 9 |
| | Care, 2005, 33, 188-95 Predictors of physician confidence to diagnose pneumonia and determine illness severity in ventilated patients. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia | | 9 |
| 80 | Care, 2005, 33, 188-95 Predictors of physician confidence to diagnose pneumonia and determine illness severity in ventilated patients. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia and Intensive Care, 2005, 33, 112-9 The spectrum of practice in the diagnosis and management of pneumonia in patients requiring mechanical ventilation. Australian and New Zealand practice in intensive care (ANZPIC II). | 1.1 | |
| 80 79 | Care, 2005, 33, 188-95 Predictors of physician confidence to diagnose pneumonia and determine illness severity in ventilated patients. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia and Intensive Care, 2005, 33, 112-9 The spectrum of practice in the diagnosis and management of pneumonia in patients requiring mechanical ventilation. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia and Intensive Care, 2005, 33, 87-100 Pharmacokinetics of colistin methanesulfonate and colistin in a critically ill patient receiving | 1.1 | 15 |
| 80 79 78 | Predictors of physician confidence to diagnose pneumonia and determine illness severity in ventilated patients. Australian and New Zealand practice in intensive care (ANZPIC II). <i>Anaesthesia and Intensive Care</i> , 2005 , 33, 112-9 The spectrum of practice in the diagnosis and management of pneumonia in patients requiring mechanical ventilation. Australian and New Zealand practice in intensive care (ANZPIC II). <i>Anaesthesia and Intensive Care</i> , 2005 , 33, 87-100 Pharmacokinetics of colistin methanesulfonate and colistin in a critically ill patient receiving continuous venovenous hemodiafiltration. <i>Antimicrobial Agents and Chemotherapy</i> , 2005 , 49, 4814-5 Serum procalcitonin and C-reactive protein as markers of sepsis and outcome in patients with | 1.1 1.1 5.9 | 15 |
| 80 79 78 77 | Predictors of physician confidence to diagnose pneumonia and determine illness severity in ventilated patients. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia and Intensive Care, 2005, 33, 112-9 The spectrum of practice in the diagnosis and management of pneumonia in patients requiring mechanical ventilation. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia and Intensive Care, 2005, 33, 87-100 Pharmacokinetics of colistin methanesulfonate and colistin in a critically ill patient receiving continuous venovenous hemodiafiltration. Antimicrobial Agents and Chemotherapy, 2005, 49, 4814-5 Serum procalcitonin and C-reactive protein as markers of sepsis and outcome in patients with neurotrauma and subarachnoid haemorrhage. Anaesthesia and Intensive Care, 2004, 32, 465-70 Vancomycin pharmacokinetics in critically ill patients receiving continuous venovenous | 1.1 1.1 5.9 | 15 108 47 |
| 80 79 78 77 76 | Predictors of physician confidence to diagnose pneumonia and determine illness severity in ventilated patients. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia and Intensive Care, 2005, 33, 112-9 The spectrum of practice in the diagnosis and management of pneumonia in patients requiring mechanical ventilation. Australian and New Zealand practice in intensive care (ANZPIC II). Anaesthesia and Intensive Care, 2005, 33, 87-100 Pharmacokinetics of colistin methanesulfonate and colistin in a critically ill patient receiving continuous venovenous hemodiafiltration. Antimicrobial Agents and Chemotherapy, 2005, 49, 4814-5 Serum procalcitonin and C-reactive protein as markers of sepsis and outcome in patients with neurotrauma and subarachnoid haemorrhage. Anaesthesia and Intensive Care, 2004, 32, 465-70 Vancomycin pharmacokinetics in critically ill patients receiving continuous venovenous haemodiafiltration. British Journal of Clinical Pharmacology, 2004, 58, 259-68 Adult-population incidence of severe sepsis in Australian and New Zealand intensive care units. | 1.1 1.1 5.9 1.1 3.8 | 15 108 47 86 |

| 72 | The effect of adsorption, filter material and point of dilution on antibiotic elimination by haemofiltration an in vitro study of levofloxacin. <i>International Journal of Antimicrobial Agents</i> , 2004 , 24, 468-72 | 14.3 | 23 |
|----|--|------|-----|
| 71 | The effect of adsorption, filter material and point of dilution on antibiotic elimination by haemofiltrationAn in vitro study of levofloxacin. <i>International Journal of Antimicrobial Agents</i> , 2004 , 24, 468-472 | 14.3 | 41 |
| 70 | Cefepime versus cefpirome: the importance of creatinine clearance. <i>Anesthesia and Analgesia</i> , 2003 , 97, 1149-1154 | 3.9 | 68 |
| 69 | Pharmacological principles of antibiotic prescription in the critically ill. <i>Anaesthesia and Intensive Care</i> , 2002 , 30, 134-44 | 1.1 | 88 |
| 68 | High dependency units: issues to consider in their planning. <i>Anaesthesia and Intensive Care</i> , 2002 , 30, 348-54 | 1.1 | 16 |
| 67 | Ciprofloxacin pharmacokinetic profiles in paediatric sepsis: how much ciprofloxacin is enough?. <i>Intensive Care Medicine</i> , 2002 , 28, 493-500 | 14.5 | 28 |
| 66 | Intravascular administration sets are accurate and in appropriate condition after 7 days of continuous use: an in vitro study. <i>Journal of Advanced Nursing</i> , 2002 , 37, 330-7 | 3.1 | 6 |
| 65 | Continuous measurement of cerebral blood flow velocity using transcranial Doppler reveals significant moment-to-moment variability of data in healthy volunteers and in patients with subarachnoid hemorrhage. <i>Critical Care Medicine</i> , 2002 , 30, 563-9 | 1.4 | 27 |
| 64 | Low cefpirome levels during twice daily dosing in critically ill septic patients: pharmacokinetic modelling calls for more frequent dosing. <i>Intensive Care Medicine</i> , 2001 , 27, 363-70 | 14.5 | 60 |
| 63 | Pharmacokinetics of ciprofloxacin in ICU patients on continuous veno-venous haemodiafiltration. <i>Intensive Care Medicine</i> , 2001 , 27, 665-72 | 14.5 | 37 |
| 62 | The pharmacokinetics of once-daily dosing of ceftriaxone in critically ill patients. <i>Journal of Antimicrobial Chemotherapy</i> , 2001 , 47, 421-9 | 5.1 | 140 |
| 61 | Subcutaneous oxygen tensions provide similar information to ileal luminal CO2 tensions in an animal model of haemorrhagic shock. <i>Intensive Care Medicine</i> , 2000 , 26, 592-600 | 14.5 | 30 |
| 60 | The treatment of tetanus with intrathecal baclofen. <i>Anaesthesia and Intensive Care</i> , 2000 , 28, 438-42 | 1.1 | 15 |
| 59 | Can we achieve consensus on central venous catheter-related infections?. <i>Anaesthesia and Intensive Care</i> , 2000 , 28, 475-90 | 1.1 | 31 |
| 58 | Cerebrospinal fluid penetration of high doses of intravenous ciprofloxacin in meningitis. <i>Clinical Infectious Diseases</i> , 2000 , 31, 1131-3 | 11.6 | 43 |
| 57 | Towards better ICU antibiotic dosing. <i>Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine</i> , 2000 , 2, 282-9 | 2.8 | 1 |
| 56 | Low plasma cefepime levels in critically ill septic patients: pharmacokinetic modeling indicates improved troughs with revised dosing. <i>Antimicrobial Agents and Chemotherapy</i> , 1999 , 43, 2559-61 | 5.9 | 98 |
| 55 | Small tidal volume ventilation using a zero deadspace tracheal tube. <i>British Journal of Anaesthesia</i> , 1999 , 82, 213-6 | 5.4 | 4 |

| 54 | Continuous infusion ceftazidime in intensive care: a randomized controlled trial. <i>Journal of Antimicrobial Chemotherapy</i> , 1999 , 43, 309-11 | 5.1 | 72 |
|----|---|----------------|----|
| 53 | Acute respiratory distress syndrome precipitated by transjugular intrahepatic porto-systemic shunting for severe hepatic veno-occlusive disease. Is it due to pulmonary leucostasis?. <i>Intensive Care Medicine</i> , 1999 , 25, 1332-3 | 14.5 | 5 |
| 52 | Inter observer variability of the transcranial Doppler ultrasound technique: impact of lack of practice on the accuracy of measurement. <i>Journal of Clinical Monitoring and Computing</i> , 1999 , 15, 179-8. | 4 ² | 47 |
| 51 | Keeping relatives informed by the internet. <i>Lancet, The</i> , 1999 , 354, 517-8 | 40 | |
| 50 | Vancomycin resistance in gram-positive cocciold bugs, new tricks, confusing terminology. <i>Anaesthesia and Intensive Care</i> , 1999 , 27, 396-9 | 1.1 | |
| 49 | Critical care research and pre-emptive informed consent: a practical approach used in Chris Hani Baragwanath ICU. <i>Intensive Care Medicine</i> , 1998 , 24, 353-7 | 14.5 | 10 |
| 48 | Good clinical research practice: what is it and is it possible in the intensive care unit?. <i>Anaesthesia and Intensive Care</i> , 1998 , 26, 568-74 | 1.1 | 1 |
| 47 | Pharmacokinetic profiles of high-dose intravenous ciprofloxacin in severe sepsis. The Baragwanath Ciprofloxacin Study Group. <i>Antimicrobial Agents and Chemotherapy</i> , 1998 , 42, 2235-9 | 5.9 | 77 |
| 46 | Intermittent bolus dosing of ceftazidime in critically ill patients. <i>Journal of Antimicrobial Chemotherapy</i> , 1997 , 40, 269-73 | 5.1 | 70 |
| 45 | Critical care in Africa. North to south and the future with special reference to southern Africa. <i>Critical Care Clinics</i> , 1997 , 13, 255-65 | 4.5 | 14 |
| 44 | Gastric intramucosal pH and blood lactate in severe sepsis. <i>Anaesthesia</i> , 1997 , 52, 726-32 | 6.6 | 19 |
| 43 | Performance of the Pediatric Risk of Mortality Score. <i>Critical Care Medicine</i> , 1997 , 25, 717 | 1.4 | |
| 42 | Pitfalls in the prediction of renal function in the intensive care unit. A review. <i>South African Journal of Surgery</i> , 1997 , 35, 16-9 | 0.4 | 8 |
| 41 | Measurements of glomerular filtration in the intensive care unit are only a rough guide to renal function. <i>South African Journal of Surgery</i> , 1997 , 35, 20-3 | 0.4 | 27 |
| 40 | Predictors of mortality in severe malaria: a two year experience in a non-endemic area. <i>Anaesthesia and Intensive Care</i> , 1996 , 24, 217-23 | 1.1 | 33 |
| 39 | Acute stridor as a presentation of bilateral abductor vocal cord paralysis. <i>Anaesthesia</i> , 1996 , 51, 1046-8 | 6.6 | 4 |
| 38 | A guideline for competency of the critical care nurse. <i>American Journal of Critical Care</i> , 1996 , 5, 217-226 | 1.7 | 13 |
| 37 | Development and validation of a critical care patient classification system. <i>American Journal of Critical Care</i> , 1996 , 5, 282-288 | 1.7 | 2 |

| 36 | Poor discriminatory performance of the Pediatric Risk of Mortality (PRISM) score in a South African intensive care unit. <i>Critical Care Medicine</i> , 1996 , 24, 1507-13 | 1.4 | 37 |
|----|---|--------------|-----|
| 35 | Accidental intra-arterial adrenaline administration. <i>Anaesthesia and Intensive Care</i> , 1995 , 23, 488-9 | 1.1 | 2 |
| 34 | Failed nasogastric intubation in undiagnosed carcinoma of the oesophagus. <i>Anaesthesia and Intensive Care</i> , 1995 , 23, 370-2 | 1.1 | |
| 33 | Changes in vancomycin pharmacokinetics in critically ill infants. <i>Anaesthesia and Intensive Care</i> , 1995 , 23, 678-82 | 1.1 | 45 |
| 32 | Emergency percutaneous balloon mitral valvotomy in a patient with septic shock. <i>Chest</i> , 1995 , 108, 570 | -3 .3 | 2 |
| 31 | Paralytic ileus as a complication of atropine therapy following severe organophosphate poisoning. <i>Anaesthesia</i> , 1994 , 49, 791-3 | 6.6 | 6 |
| 30 | A double-blind comparison of vecuronium administered by the Springfusor infusion device to vecuronium by intermittent bolus injection in critically ill adult patients. <i>Anaesthesia and Intensive Care</i> , 1994 , 22, 580-5 | 1.1 | 4 |
| 29 | Malignant astrocytoma of the cervico-medullary junction masquerading as Guillain-Barrßyndrome. <i>Postgraduate Medical Journal</i> , 1994 , 70, 499-502 | 2 | O |
| 28 | Haemodynamic and oxygen transport response during exchange transfusion for severe falciparum malaria. <i>Postgraduate Medical Journal</i> , 1994 , 70, 801-4 | 2 | 9 |
| 27 | Decreased cardiac index as an indicator of tension pneumothorax in the ventilated patient. <i>Anaesthesia</i> , 1994 , 49, 137-41 | 6.6 | 14 |
| 26 | A comparison of arterial lines and insertion techniques in critically ill patients. <i>Anaesthesia</i> , 1994 , 49, 968-73 | 6.6 | 21 |
| 25 | Esmolol in a case of severe tetanus. Adequate haemodynamic control achieved despite markedly elevated catecholamine levels. <i>South African Journal of Surgery</i> , 1994 , 32, 33-5 | 0.4 | 4 |
| 24 | Successful use of continuous veno-venous haemofiltration to treat profound fluid retention in severe peripartum cardiomyopathy. <i>Anaesthesia</i> , 1993 , 48, 1065-7 | 6.6 | 10 |
| 23 | Acute lung injury at Baragwanath ICU. An eight-month audit and call for consensus for other organ failure in the adult respiratory distress syndrome. <i>Chest</i> , 1993 , 103, 1832-6 | 5.3 | 16 |
| 22 | Hydrocortisone and tumor necrosis factor in severe community-acquired pneumonia. A randomized controlled study. <i>Chest</i> , 1993 , 104, 389-92 | 5.3 | 117 |
| 21 | Intensive care utilisation: the Baragwanath experience. <i>Anaesthesia and Intensive Care</i> , 1993 , 21, 396-9 | 1.1 | 7 |
| 20 | High-dose adrenaline in adult in-hospital asystolic cardiopulmonary resuscitation: a double-blind randomised trial. <i>Anaesthesia and Intensive Care</i> , 1993 , 21, 192-6 | 1.1 | 40 |
| 19 | Renal replacement therapy for the critically illprecarious progress. Part II. Technical aspects and clinical application. <i>South African Journal of Surgery</i> , 1993 , 31, 147-51 | 0.4 | 4 |

| 18 | Renal replacement therapy for the critically illprecarious progress. Part I. Definitions and physiological aspects. <i>South African Journal of Surgery</i> , 1993 , 31, 114-20 | 0.4 | 4 |
|----|--|------------------|-----|
| 17 | Septic shock: does adrenaline have a role as a first-line inotropic agent?. <i>Anaesthesia and Intensive Care</i> , 1992 , 20, 470-4 | 1.1 | 37 |
| 16 | Erythropoietin in a patient following multiple trauma. <i>Anaesthesia</i> , 1992 , 47, 962-4 | 6.6 | 20 |
| 15 | Comparison of disease severity scoring systems in septic shock. <i>Critical Care Medicine</i> , 1991 , 19, 1165-7 | 1 _{1.4} | 30 |
| 14 | The pharmacokinetic of amikacin in critically ill adult and paediatric patients: comparison of onceversus twice-daily dosing regimens. <i>Journal of Antimicrobial Chemotherapy</i> , 1991 , 27 Suppl C, 81-9 | 5.1 | 55 |
| 13 | A prospective randomized study comparing once- versus twice-daily amikacin dosing in critically ill adult and paediatric patients. <i>Journal of Antimicrobial Chemotherapy</i> , 1991 , 28, 753-64 | 5.1 | 107 |
| 12 | Typhoid fever: an undescribed cause of adult respiratory distress syndrome. <i>Anaesthesia and Intensive Care</i> , 1991 , 19, 124-6 | 1.1 | 5 |
| 11 | Vasoconstrictor effects of adrenaline in human septic shock. <i>Anaesthesia and Intensive Care</i> , 1991 , 19, 61-5 | 1.1 | 46 |
| 10 | Acute preload effects of furosemide. <i>Chest</i> , 1990 , 98, 124-8 | 5.3 | 23 |
| 9 | Coronary embolism causing myocardial infarction. <i>Intensive Care Medicine</i> , 1990 , 16, 215-6 | 14.5 | 30 |
| 8 | Difficulty in weaning from respiratory support in a patient with the intermediate syndrome of organophosphate poisoning. <i>Critical Care Medicine</i> , 1989 , 17, 1075-6 | 1.4 | 12 |
| 7 | Catheter related infection. A plea for consensus with review and guidelines. <i>Intensive Care Medicine</i> , 1988 , 14, 503-9 | 14.5 | 34 |
| 6 | Fatal infection caused by a multiply resistant type 3 pneumococcus. <i>Journal of Clinical Microbiology</i> , 1988 , 26, 1590-1 | 9.7 | 8 |
| 5 | Autonomic dysfunction in severe tetanus. <i>Critical Care Medicine</i> , 1987 , 15, 987-988 | 1.4 | 56 |
| 4 | Pitfalls in the interpretation of pulmonary capillary wedge pressure. <i>South African Medical Journal</i> , 1985 , 67, 174-7 | 1.5 | 7 |
| 3 | Vitamin B12 levels in the prolonged use of sodium nitroprusside. <i>Critical Care Medicine</i> , 1984 , 12, 161-3 | 1.4 | 4 |
| 2 | Intravenous Calcium chloride as an antidote to verapamil-induced hypotension. <i>Intensive Care Medicine</i> , 1982 , 8, 55-7 | 14.5 | 35 |
| 1 | Use and Abuse of Antibiotics128-136 | | |