

Catalina Ruiz PÃ©rez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5251459/publications.pdf>

Version: 2024-02-01

253
papers

10,031
citations

32410

55
h-index

60403

85
g-index

271
all docs

271
docs citations

271
times ranked

6735
citing authors

#	ARTICLE	IF	CITATIONS
1	Metal organic framework composite, nano-Fe ₃ O ₄ @Fe-(benzene-1,3,5-tricarboxylic acid), for solid phase extraction of blood lipid regulators from water. <i>Talanta</i> , 2020, 207, 120275.	2.9	32
2	Electroswitching of the single-molecule magnet behaviour in an octahedral spin crossover cobalt(II) complex with a redox-active pyridinediimine ligand. <i>Chemical Communications</i> , 2020, 56, 12242-12245.	2.2	8
3	Synthesis, crystal structure and magnetic properties of the cyclic tetranuclear compound [Cu ₄ (pz) ₄ (hppa) ₂ (H ₂ O) ₄] [pz ⁻ = ⁻ pyrazolate; hppa ⁻ = ⁻ R,S-2-hydroxo-2-phenyl-2-(1-pyrazolyl)acetate]. <i>Polyhedron</i> , 2019, 170, 217-222.	1.0	4
4	Application of a Pillared-Layer Zn-Triazolate Metal-Organic Framework in the Dispersive Miniaturized Solid-Phase Extraction of Personal Care Products from Wastewater Samples. <i>Molecules</i> , 2019, 24, 690.	1.7	20
5	Ferromagnetic coupling through the oxalate bridge in heterobimetallic Cr(III)-M(II) (M = Mn and Co) assemblies. <i>Comptes Rendus Chimie</i> , 2019, 22, 452-465.	0.2	7
6	One-dimensional oxalato-bridged heterobimetallic coordination polymers by using [the [Cr(pyim)(C ₂ O ₄) ₂] ⁺ complex as metalloligand [pyim ⁻ = ⁻ 2-(2-pyridyl)imidazole]. <i>Inorganica Chimica Acta</i> , 2019, 486, 150-157.		4
7	Anion-π Interactions in Hollow Crystals of a Copper(II)-Cyamelurate Coordination Complex. <i>Crystal Growth and Design</i> , 2018, 18, 2636-2644.	1.4	12
8	Halogen bonding two-point recognition with terphenyl derivatives. <i>New Journal of Chemistry</i> , 2018, 42, 10476-10480.	1.4	17
9	Dicopper(II) metallacyclophanes with photoswitchable oligoacene spacers: a joint experimental and computational study on molecular magnetic photoswitches. <i>Journal of Coordination Chemistry</i> , 2018, 71, 675-692.	0.8	3
10	Insights in the analytical performance of neat metal-organic frameworks in the determination of pollutants of different nature from waters using dispersive miniaturized solid-phase extraction and liquid chromatography. <i>Talanta</i> , 2018, 179, 775-783.	2.9	52
11	Magnetostructural relationships in polymorphic ethylmalonate-containing copper(II) coordination polymers. <i>CrystEngComm</i> , 2018, 20, 7464-7472.	1.3	3
12	Influence of Ligand Functionalization of UiO-66-Based Metal-Organic Frameworks When Used as Sorbents in Dispersive Solid-Phase Analytical Microextraction for Different Aqueous Organic Pollutants. <i>Molecules</i> , 2018, 23, 2869.	1.7	40
13	A green metal-organic framework to monitor water contaminants. <i>RSC Advances</i> , 2018, 8, 31304-31310.	1.7	34
14	Crystal structure analysis of supramolecular arrangements in bis(1-isoquinolinecarboxamide)alkanes and their Ag(I) complexes. <i>CrystEngComm</i> , 2017, 19, 1076-1088.	1.3	2
15	Photoluminescent and Slow Magnetic Relaxation Studies on Lanthanide(III)-2,5-pyrazinedicarboxylate Frameworks. <i>Inorganic Chemistry</i> , 2017, 56, 2108-2123.	1.9	49
16	Reversible solvatomagnetic switching in a single-ion magnet from an entatic state. <i>Chemical Science</i> , 2017, 8, 3694-3702.	3.7	67
17	Tailor-made copper(II) coordination polymers based on the C ₃ symmetric methanetriacetate as a ligand. <i>CrystEngComm</i> , 2017, 19, 376-390.	1.3	1
18	Cadmium(II) coordination polymers based on substituted malonic acid: synthesis, characterization and photoluminescence properties. <i>Inorganic Chemistry Frontiers</i> , 2017, 4, 1384-1392.	3.0	10

#	ARTICLE	IF	CITATIONS
19	Metal-organic and supramolecular lead(II) networks assembled from isomeric nicotinoylhydrazone blocks: the effects of ligand geometry and counter-ion on topology and supramolecular assembly. <i>CrystEngComm</i> , 2016, 18, 5375-5385.	1.3	40
20	Crystal growth and structural remarks on malonate-based lanthanide coordination polymers. <i>CrystEngComm</i> , 2016, 18, 7831-7842.	1.3	11
21	Inorganic-organic hybrid materials based on PbBr ₂ and pyridine-hydrazone blocks: structural and theoretical study. <i>RSC Advances</i> , 2016, 6, 60385-60393.	1.7	24
22	Bioinspired Citrate-Apatite Nanocrystals Doped with Divalent Transition Metal Ions. <i>Crystal Growth and Design</i> , 2016, 16, 145-153.	1.4	32
23	Adsorption of silver nanoparticles from aqueous solution on copper-based metal organic frameworks (HKUST-1). <i>Chemosphere</i> , 2016, 150, 659-666.	4.2	39
24	A magnetic-based dispersive micro-solid-phase extraction method using the metal-organic framework HKUST-1 and ultra-high-performance liquid chromatography with fluorescence detection for determining polycyclic aromatic hydrocarbons in waters and fruit tea infusions. <i>Journal of Chromatography A</i> , 2016, 1436, 42-50.	1.8	100
25	Extending the halogen-bonded supramolecular synthon concept to 1,3,4-oxadiazole derivatives. <i>CrystEngComm</i> , 2016, 18, 42-47.	1.3	11
26	Synthesis and structural characterization of six Cu(II)-based coordination polymers using the thermally tunable 1,2,3,4-cyclobutanetetracarboxylic acid. <i>CrystEngComm</i> , 2015, 17, 5081-5093.	1.3	7
27	Metallosupramolecular approach toward multifunctional magnetic devices for molecular spintronics. <i>Coordination Chemistry Reviews</i> , 2015, 303, 110-138.	9.5	64
28	The metal-organic framework HKUST-1 as efficient sorbent in a vortex-assisted dispersive micro solid-phase extraction of parabens from environmental waters, cosmetic creams, and human urine. <i>Talanta</i> , 2015, 139, 13-20.	2.9	144
29	Influence of the coligand in the magnetic properties of a series of copper(II)-phenylmalonate complexes. <i>CrystEngComm</i> , 2014, 16, 8106-8118.	1.3	14
30	Synthesis, Crystal Structure, and Magnetic Characterization of the Three-Dimensional Compound [Co ₂ (cbut)(H ₂ O) ₃] _n (H ₄ cbut = Tj ETQq0100 rgBT 16 verlock 1		
31	Syntheses, crystal structures and magnetic properties of five new manganese(II) complexes: influence of the conformation of different alkyl/aryl substituted malonate ligands on the crystal packing. <i>CrystEngComm</i> , 2014, 16, 2766.	1.3	27
32	Two-Dimensional 3d-4f Heterometallic Coordination Polymers: Syntheses, Crystal Structures, and Magnetic Properties of Six New Co(II)-Ln(III) Compounds. <i>Inorganic Chemistry</i> , 2014, 53, 6299-6308.	1.9	20
33	Three new europium(III) methanetriacetate metal-organic frameworks: the influence of synthesis on the product topology. <i>Acta Crystallographica Section B: Structural Science, Crystal Engineering and Materials</i> , 2014, 70, 19-27.	0.5	5
34	Conformational influence of quinoline moieties in the crystal packing of bis(quinolinecarboxamide)alkane derivatives. <i>CrystEngComm</i> , 2013, 15, 7212.	1.3	3
35	Low-Dimensional 3d-4f Complexes Assembled by Low-Spin [Fe ^{III} (phen)(CN) ₄] ⁿ⁺ Anions. <i>Inorganic Chemistry</i> , 2013, 52, 1525-1537.	1.9	35
36	Synthesis, Crystal Structure and Magnetic Characterization of a Series of Compounds with an Unusual Single Crystal to Single Crystal Phase Transition. <i>Crystal Growth and Design</i> , 2013, 13, 4735-4745.	1.4	16

#	ARTICLE	IF	CITATIONS
37	Neutron Diffraction Studies of the Molecular Compound [Co ₂ (bta)] _n (H ₄ bta = 1,2,4,5-Benzenetetracarboxylic Acid): In the Quest of Canted Ferromagnetism. <i>Inorganic Chemistry</i> , 2013, 52, 12818-12827.	1.9	10
38	A step further in the comprehension of the magnetic coupling in gadolinium(III)-based carboxylate complexes. <i>Polyhedron</i> , 2013, 52, 321-332.	1.0	23
39	Synthesis, crystal structure and magnetic properties of a new cyanide-bridged mixed-valence copper(I)/copper(II) clathrate. <i>Inorganic Chemistry Communication</i> , 2013, 35, 252-254.	1.8	5
40	Synthesis, crystal structures and tautomerism in novel oximes based on hydroxyalkylpyrazolones. <i>New Journal of Chemistry</i> , 2013, 37, 2002.	1.4	6
41	Copper(ii) assembling with bis(2-pyridylcarbonyl)amidate and N,N'-2,2-phenylenebis(oxamate). <i>Dalton Transactions</i> , 2013, 42, 5778.	1.6	35
42	Magnetic coupling and spin topology in linear oxalato-bridged tetranuclear chromium(III)-copper(II) complexes with aromatic diimine ligands. <i>Polyhedron</i> , 2013, 52, 1246-1255.	1.0	14
43	Self-assembly of a chiral three-dimensional manganese(ii)-copper(ii) coordination polymer with a double helical architecture. <i>CrystEngComm</i> , 2013, 15, 9312.	1.3	17
44	Dicopper(II) Metallacyclophanes with Electroswitchable Polymethyl-Substituted <i>para</i> -Phenylene Spacers. <i>Chemistry - A European Journal</i> , 2013, 19, 12124-12137.	1.7	25
45	Self-assembly, metal binding ability, and magnetic properties of dinickel(ii) and dicobalt(ii) triple mesocates. <i>CrystEngComm</i> , 2012, 14, 5639.	1.3	14
46	Redox switching of the antiferromagnetic coupling in permethylated dicopper(ii) paracyclophanes. <i>Chemical Communications</i> , 2012, 48, 8401.	2.2	22
47	Ligand effects on the dimensionality of oxamato-bridged mixed-metal open-framework magnets. <i>Chemical Communications</i> , 2012, 48, 3539.	2.2	15
48	Influence of the alkaline earth cations on the topology of M ^{II} /Cu ^{II} mixed-metal-organic frameworks (M = Ca, Sr and Ba). <i>CrystEngComm</i> , 2012, 14, 761-764.	1.3	17
49	Molecular Engineering To Control the Magnetic Interaction between Single-Chain Magnets Assembled in a Two-Dimensional Network. <i>Journal of the American Chemical Society</i> , 2012, 134, 15265-15268.	6.6	67
50	Pillaring Role of 4,4'-Azobis(pyridine) in Substituted Malonate-Containing Manganese(II) Complexes: Syntheses, Crystal Structures, and Magnetic Properties. <i>Crystal Growth and Design</i> , 2012, 12, 4505-4518.	1.4	20
51	Highly Selective Chemical Sensing in a Luminescent Nanoporous Magnet. <i>Advanced Materials</i> , 2012, 24, 5625-5629.	11.1	131
52	Dryness sensitive porous 3d-4f metal-organic framework with unusual dynamic behaviour. <i>CrystEngComm</i> , 2012, 14, 765-767.	1.3	20
53	Synthesis, structural analysis, and thermal and spectroscopic studies of methylmalonate-containing zinc(II) complexes. <i>Comptes Rendus Chimie</i> , 2012, 15, 911-923.	0.2	6
54	Novel Malonate-Containing Coordination Compounds with Ligands Having N- and NO-Donors: Synthesis, Structures, and Magnetic Properties. <i>Crystal Growth and Design</i> , 2012, 12, 599-614.	1.4	27

#	ARTICLE	IF	CITATIONS
55	[FeIII(dmbpy)(CN)4]•: a new building block for designing single-chain magnets. Dalton Transactions, 2012, 41, 13716.	1.6	33
56	Two-dimensional (6,3) networks obtained with the {Cu3(Hmesox)3}• secondary building unit (H4mesox = mesoxalic acid). CrystEngComm, 2012, 14, 4289.	1.3	11
57	Selective Gas and Vapor Sorption and Magnetic Sensing by an Isoreticular Mixed-Metal•Organic Framework. Journal of the American Chemical Society, 2012, 134, 15301-15304.	6.6	109
58	Theoretical Insights into the Ferromagnetic Coupling in Oxalato-Bridged Chromium(III)-Cobalt(II) and Chromium(III)-Manganese(II) Dinuclear Complexes with Aromatic Diimine Ligands. Inorganic Chemistry, 2012, 51, 3289-3301.	1.9	17
59	Slow Relaxation of the Magnetization in a 4,2-Wavelike Fe^{III}₂Co^{II} Heterobimetallic Chain. Inorganic Chemistry, 2012, 51, 1216-1218.	1.9	33
60	Solid-State Aggregation of Metallacyclophane-Based Mn^{II}Cu^{II} One-Dimensional Ladders. Inorganic Chemistry, 2012, 51, 7019-7021.	1.9	15
61	Copper(II) complexes with 2,5-bis(2-pyridyl)pyrazine and oxalate and croconate: Synthesis, crystal structure and magnetic properties. Inorganica Chimica Acta, 2012, 389, 52-59.	1.2	8
62	Pentacyanido(L)ferrate(III) complexes: Crystal structures, electrochemical and DFT studies (L=pyrazine, 4,4•bipyridine, azide). Polyhedron, 2012, 31, 539-547.	1.0	7
63	Reversible Solvatomagnetic Switching in a Spongelike Manganese(II)•Copper(II) 3D Open Framework with a Pillared Square/Octagonal Layer Architecture. Chemistry - A European Journal, 2012, 18, 1608-1617.	1.7	86
64	SYNTHESIS AND CHARACTERIZATION OF BIS- AND TOS-(4-CARBOXYBENZOYL)-ALKANEAMINES. Journal of the Chilean Chemical Society, 2012, 57, 1305-1308.	0.5	1
65	Photoswitching of the antiferromagnetic coupling in an oxamato-based dicopper(ii) anthracenophane. Chemical Communications, 2011, 47, 11035.	2.2	39
66	Solid-State Dinuclear-to-Trinuclear Conversion in an Oxalato-Bridged Chromium(III)•Cobalt(II) Complex as a New Route toward Single-Molecule Magnets. Inorganic Chemistry, 2011, 50, 2073-2075.	1.9	30
67	Spin Control in Oxamato-Based Manganese(II)•Copper(II) Coordination Polymers with Brick-Wall Layer Architectures. Inorganic Chemistry, 2011, 50, 8694-8696.	1.9	33
68	Synthesis, Structural Analysis, and Magnetic Properties of Ethylmalonate-Manganese(II) Complexes. Inorganic Chemistry, 2011, 50, 10765-10776.	1.9	15
69	New cyanide-bridged MnIII•MIII heterometallic dinuclear complexes constructed from [MIII(AA)(CN)4]• building blocks (M = Cr and Fe): synthesis, crystal structures and magnetic properties. Dalton Transactions, 2011, 40, 4898.	1.6	27
70	A three-dimensional copper(ii) 12-metallacrown-4 complex with malonomonohydroxamic acid (H3mmh) as a ligand. New Journal of Chemistry, 2011, 35, 1817.	1.4	26
71	Copper(II)-phenylmalonate complexes with the bifunctional ligands nicotinamide and isonicotinamide. Polyhedron, 2011, 30, 2451-2458.	1.0	10
72	Chromium(III) complexes with 2-(2•pyridyl)imidazole: Synthesis, crystal structure and magnetic properties. Inorganica Chimica Acta, 2011, 376, 358-366.	1.2	21

#	ARTICLE	IF	CITATIONS
73	Synthesis, Crystal Structures and Magnetic Properties of $M^{II}Cu^{II}$ Chains (M=Mn and Co) with Sterically Hindered Alkyl-Substituted Phenyloxamate Bridging Ligands. Chemistry - A European Journal, 2011, 17, 2176-2188.	1.7	58
74	Rational Enantioselective Design of Chiral Heterobimetallic Single-Chain Magnets: Synthesis, Crystal Structures and Magnetic Properties of Oxamato-Bridged $M^{II}Cu^{II}$ Chains (M=Mn, Co). Chemistry - A European Journal, 2011, 17, 12482-12494.	1.7	78
75	Metal-organic coordination frameworks based on mixed methylmalonate and 4,4'-bipyridine ligands: synthesis, crystal structure and magnetic properties. New Journal of Chemistry, 2010, 34, 2515.	1.4	24
76	Single chain magnet behaviour in an enantiopure chiral cobalt(ii)-copper(ii) one-dimensional compound. Chemical Communications, 2010, 46, 2322.	2.2	100
77	Supramolecular coordination chemistry of aromatic polyoxalamide ligands: A metallosupramolecular approach toward functional magnetic materials. Coordination Chemistry Reviews, 2010, 254, 2281-2296.	9.5	178
78	The Construction of Open Gd^{III} Metal-Organic Frameworks Based on Methanetricetic Acid: New Objects with an Old Ligand. Chemistry - A European Journal, 2010, 16, 4037-4047.	1.7	37
79	Oligo- <i>m</i> -phenyleneoxalamide Copper(II) Mesocates as Electro-Switchable Ferromagnetic Metal-Organic Wires. Chemistry - A European Journal, 2010, 16, 12838-12851.	1.7	30
80	Inside Cover: Oligo- <i>m</i> -phenyleneoxalamide Copper(II) Mesocates as Electro-Switchable Ferromagnetic Metal-Organic Wires (Chem. Eur. J. 43/2010). Chemistry - A European Journal, 2010, 16, 12762-12762.	1.7	0
81	All-cis-1,2,3,4,5,6-cyclohexanehexacarboxylate two-dimensional gadolinium(III) complexes: Synthesis, X-ray crystal structure and magnetic properties. Polyhedron, 2010, 29, 188-195.	1.0	13
82	Topological control of the spin coupling in dinuclear copper(II) complexes with meta- and para-phenylenediamine bridging ligands. Inorganica Chimica Acta, 2010, 363, 1984-1994.	1.2	23
83	Long-distance magnetic coupling in dinuclear copper(II) complexes with oligo-para-phenylenediamine bridging ligands. Inorganica Chimica Acta, 2010, 363, 1666-1678.	1.2	34
84	$[Cu_3(Hmesox)_3]^{3+}$: a Precursor for the Rational Design of Chiral Molecule-Based Magnets ($H_4mesox = 2$ -dihydroxymalonic acid). Inorganic Chemistry, 2010, 49, 7880-7889.	1.9	18
85	Tuning the Spin Ground State in Heterononanuclear Nickel(II)-Copper(II) Cylinders with a Triangular Metallacyclophane Core. Inorganic Chemistry, 2010, 49, 11264-11266.	1.9	5
86	Building-block process for the synthesis of new chromium(III) malonate complexes. CrystEngComm, 2010, 12, 2711.	1.3	19
87	Magnetic coupling in discrete cyano-bridged $M^{III}Fe^{III}$ motifs: Synthesis, crystal structure, magnetic properties and theoretical study. Dalton Transactions, 2010, 39, 5028.	1.6	39
88	Variation of the ground spin state in homo- and hetero-octanuclear copper(II) and nickel(II) double-star complexes with a meso-helicate-type metallacryptand core. Dalton Transactions, 2010, 39, 4786.	1.6	11
89	Ferromagnetic coupling and magnetic anisotropy in oxalato-bridged trinuclear chromium(III)-cobalt(II) complexes with aromatic diimine ligands. Dalton Transactions, 2010, 39, 2350-2358.	1.6	42
90	Dinuclear, tetranuclear and one-dimensional pyrazine-based copper(II) complexes: preparation, X-ray structure and magnetic properties. Dalton Transactions, 2010, 39, 167-179.	1.6	24

#	ARTICLE	IF	CITATIONS
91	Intramolecular ferro- and antiferromagnetic interactions in oxo-carboxylate bridged digadolinium(III) complexes. Dalton Transactions, 2010, 39, 7286.	1.6	51
92	[Fe(phen)(CN) ₄] ³⁻ : a suitable metalloligand unit to build 3d ⁶ -4f heterobimetallic complexes with mixed bpym-cyano bridges (phen = 1,10-phenanthroline, bpym = 2,2'-bipyrimidine). CrystEngComm, 2010, 12, 2454.	1.3	41
93	Trialkylamine versus Trialkylphosphine: Catalytic Conjugate Addition of Alcohols to Alkyl Propiolates. Synlett, 2009, 2009, 1223-1226.	1.0	25
94	Novel Lactone Chamigrene α -Derived Metabolites from <i>Laurencia majuscula</i> . European Journal of Organic Chemistry, 2009, 2009, 1407-1411.	1.2	18
95	Copper(II)-methylmalonate complexes with unidentate N-donor ligands: Syntheses, structural characterization and magnetic properties. Polyhedron, 2009, 28, 1802-1807.	1.0	15
96	New 3d ⁶ -4f supramolecular systems constructed by [Fe(bipy)(CN) ₄] ³⁻ and partially blocked lanthanide cations. Polyhedron, 2009, 28, 851-859.	1.0	18
97	Study of the Influence of the Bridge on the Magnetic Coupling in Cobalt(II) Complexes. Inorganic Chemistry, 2009, 48, 11342-11351.	1.9	81
98	Unprecedented coexistence of cyano-bridged Mn ₄ ^{III} Cr ^{III} and Mn ₂ ^{III} Cr ^{III} heterobimetallic complexes in one single crystal. Dalton Transactions, 2009, , 37-40.	1.6	25
99	Cobalt(II) Sheet-Like Systems Based on Diacetic Ligands: from Subtle Structural Variances to Different Magnetic Behaviors. Inorganic Chemistry, 2009, 48, 6086-6095.	1.9	51
100	Dinuclear and two- and three-dimensional gadolinium(III) complexes with mono- and dicarboxylate ligands: synthesis, structure and magnetic properties. CrystEngComm, 2009, 11, 2131.	1.3	64
101	Redox Switch-Off of the Ferromagnetic Coupling in a Mixed-Spin Tricobalt(II) Triple Mesocate. Journal of the American Chemical Society, 2009, 131, 14614-14615.	6.6	39
102	Low-Dimensional Copper(II) Complexes with the Trinucleating Ligand 2,4,6-Tris(di-2-pyridylamine)-1,3,5-triazine: Synthesis, Crystal Structures, and Magnetic Properties. Inorganic Chemistry, 2009, 48, 6630-6640.	1.9	33
103	Molecular-Programmed Self-Assembly of Homo- and Heterometallic Tetranuclear Coordination Compounds: Synthesis, Crystal Structures, and Magnetic Properties of Rack-Type Cu ₂ M ₂ Complexes (M = Cu and Ni) with Tetranucleating Phenylenedioxamate Bridging Ligands. Inorganic Chemistry, 2009, 48, 4661-4673.	1.9	22
104	Influence of the Aliphatic Wrapping in the Crystal Structure of Benzene Tricarboxamide Supramolecular Polymers. Crystal Growth and Design, 2009, 9, 4987-4989.	1.4	31
105	Novel cobalt(II) coordination polymers based on 1,2,4,5-benzenetetracarboxylic acid and extended bis-monodentate ligands. CrystEngComm, 2009, 11, 2169.	1.3	36
106	One-dimensional and two-dimensional coordination polymers constructed from copper(II) nodes and polycarboxylate spacers: Synthesis, crystal structures and magnetic properties. Polyhedron, 2008, 27, 574-582.	1.0	41
107	Insertion of a Spin Crossover Fe ^{III} Complex into an Oxalate-Based Layered Material: Coexistence of Spin Canting and Spin Crossover in a Hybrid Magnet. Inorganic Chemistry, 2008, 47, 9111-9120.	1.9	59
108	Synthesis, crystal structures and magnetic properties of tricyanomethanide-containing copper(II) complexes. Dalton Transactions, 2008, , 1583.	1.6	43

#	ARTICLE	IF	CITATIONS
109	(4,4) Rectangular Lattices of Cobalt(II) with 1,2,4,5-Benzenetetracarboxylic Acid: Influence of the Packing in the Crystal Structure. <i>Crystal Growth and Design</i> , 2008, 8, 3984-3992.	1.4	48
110	Crystal Structure and Magnetic Properties of Two Isomeric Three-Dimensional Pyromellitate-Containing Cobalt(II) Complexes. <i>Inorganic Chemistry</i> , 2008, 47, 8053-8061.	1.9	70
111	Well-resolved unusual alternating cyclic water tetramers embedded in a crystal host. <i>CrystEngComm</i> , 2008, 10, 1743.	1.3	43
112	Supramolecular Networks in Copper(II) Malonate Complexes. <i>Crystal Growth and Design</i> , 2008, 8, 3219-3232.	1.4	48
113	Crystal Engineering of Complexes of Propane-1,2,3-tricarboxylic Acid (H3tca) with Lanthanide(III) Cations. <i>Crystal Growth and Design</i> , 2008, 8, 1313-1318.	1.4	22
114	[MIII(bpym)(CN)4] ³⁻ : a suitable building block to design ferrimagnetic cyano-bridged heterobimetallic chains (M = Fe, Cr; bpym = 2,2'-bipyrimidine). <i>Dalton Transactions</i> , 2008, , 4103.	1.6	34
115	1,2,4,5-Benzenetetracarboxylate- and 2,2'-Bipyrimidine-Containing Cobalt(II) Coordination Polymers: Preparation, Crystal Structure, and Magnetic Properties. <i>Inorganic Chemistry</i> , 2008, 47, 3568-3576.	1.9	101
116	Heterobinuclear Complexes as Tectons in Designing Coordination Polymers. <i>Crystal Growth and Design</i> , 2008, 8, 941-949.	1.4	87
117	Heterotrimetallic Oxalato-Bridged ReIV2MII Complexes (M = Mn, Co, Ni, Cu): Synthesis, Crystal Structure, and Magnetic Properties. <i>Inorganic Chemistry</i> , 2007, 46, 3523-3530.	1.9	36
118	New anionic cobalt complexes using highly hindered bis-amides with varying donor abilities as ligands. <i>Dalton Transactions</i> , 2007, , 2135-2144.	1.6	13
119	Crystal engineering of 3-D coordination polymers by pillaring ferromagnetic copper(ii)-methylmalonate layers. <i>CrystEngComm</i> , 2007, 9, 478-487.	1.3	92
120	4,2-Ribbon like ferromagnetic cyano-bridged FeIII2NiII chains: a magneto-structural study. <i>Dalton Transactions</i> , 2007, , 3690.	1.6	43
121	Molecular-Programmed Self-Assembly of Homo- and Heterometallic Penta- and Hexanuclear Coordination Compounds: Synthesis, Crystal Structures, and Magnetic Properties of Ladder-Type Cull2MIIx(M = Cu, Ni;x= 3, 4) Oxamate Complexes with Cull2Metallacyclophane Cores. <i>Inorganic Chemistry</i> , 2007, 46, 4504-4514.	1.9	45
122	Unusual (1/4-aqua)bis(1/4-carboxylate) Bridge in Homometallic M(II) (M = Mn, Co and Ni) Two-Dimensional Compounds Based on the 1,2,3,4-Butanetetracarboxylic Acid: Synthesis, Structure, and Magnetic Properties. <i>Inorganic Chemistry</i> , 2007, 46, 7458-7465.	1.9	85
123	Structural changes in the crystal packing of highly hindered symmetrical vicinal bis-amides. <i>CrystEngComm</i> , 2007, 9, 746.	1.3	2
124	Structural versatility in cobalt(ii) complexes with 1,2,4,5-benzenetetracarboxylic acid (H4bta) and 4,4'-bipyridine-N,N'-dioxide (dpo). <i>CrystEngComm</i> , 2007, 9, 815.	1.3	69
125	Ligand Design for Heterobimetallic Single-Chain Magnets: Synthesis, Crystal Structures, and Magnetic Properties of MIIICuII (M=Mn, Co) Chains with Sterically Hindered Methyl-Substituted Phenylloxamate Bridging Ligands. <i>Chemistry - A European Journal</i> , 2007, 13, 2054-2066.	1.7	105
126	Solid-State Anion Guest Encapsulation by Metallosupramolecular Capsules Made from Two Tetranuclear Copper(II) Complexes. <i>European Journal of Inorganic Chemistry</i> , 2007, 2007, 4569-4573.	1.0	9

#	ARTICLE	IF	CITATIONS
127	Malonate-Containing Manganese(III) Complexes: Synthesis, Crystal Structure, and Magnetic Properties of $\text{AsPh}_4[\text{Mn}(\text{mal})_2(\text{H}_2\text{O})_2]$. <i>Inorganic Chemistry</i> , 2006, 45, 1012-1020.	1.9	50
128	Two- and Three-Dimensional Networks of Gadolinium(III) with Dicarboxylate Ligands: Synthesis, Crystal Structure, and Magnetic Properties. <i>Inorganic Chemistry</i> , 2006, 45, 10585-10594.	1.9	89
129	Versatile supramolecular self-assembly : Part II. Network formation and magnetic behaviour of copper(ii) malonate anions in ammonium derivatives. <i>CrystEngComm</i> , 2006, 8, 530-544.	1.3	38
130	Metamagnetism in hydrophobically induced carboxylate (phenylmalonate)-bridged copper(ii) layers. <i>Chemical Communications</i> , 2006, , 2857-2859.	2.2	32
131	Zeolite-like Nanoporous Gadolinium Complexes Incorporating Alkaline Cations. <i>Crystal Growth and Design</i> , 2006, 6, 87-93.	1.4	44
132	$[\text{Fe}(\text{bpym})(\text{CN})_4]$: A New Building Block for Designing Single-Chain Magnets. <i>Journal of the American Chemical Society</i> , 2006, 128, 4842-4853.	6.6	248
133	Influence of the presence of divalent first-row transition metal ions on the structure of sodium(i) salts of 1,2,3,4-benzenetetracarboxylic acid (H_4bta). <i>CrystEngComm</i> , 2006, 8, 338-345.	1.3	28
134	Versatile supramolecular self-assembly. Part I. Network formation and magnetic behaviour of the alkaline salts of the bis(malonate)cuprate(ii) anion. <i>CrystEngComm</i> , 2006, 8, 507-529.	1.3	40
135	A unique tautomer of 1-n-hexyl-3-phenyl-1H-pyrazol-5-ol. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2006, 62, o599-o601.	0.4	3
136	Trans-dicyanobis(acetylacetonato)ruthenate(III) as a precursor to build novel cyanide-bridged Ru(II) bimetallic compounds $[\text{M}=\text{Co}$ and $\text{Ni}]$. <i>Coordination Chemistry Reviews</i> , 2006, 250, 2176-2193.	9.5	73
137	A new synthetic route towards heterotrimetallic complexes. Synthesis, crystal structure and magnetic properties of a $[\text{CuII}(\text{MnII}(\text{CrIII}))]$ trinuclear complex. <i>Inorganica Chimica Acta</i> , 2006, 359, 433-440.	1.2	46
138	Design of single chain magnets through cyanide-bearing six-coordinate complexes. <i>Coordination Chemistry Reviews</i> , 2005, 249, 2691-2729.	9.5	417
139	9-Oxabicyclo[3.3.1]nona-2, 6-diene. Short Access and Allylic Bromination. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2005, 60, 1143-1148.	0.3	1
140	Hybrid materials containing organometallic cations and 3-D anionic metal dicyanamide networks of type $[\text{Cp}^*\text{2M}][\text{M}(\text{dca})_3]$. <i>Dalton Transactions</i> , 2005, , 285-290.	1.6	32
141	Hydrogen Bond-Directed Frameworks Based on 1,2,4,5-Benzene-Tetracarboxylate. <i>Crystal Growth and Design</i> , 2005, 5, 1163-1167.	1.4	52
142	Polymeric Networks of Copper(II) Phenylmalonate with Heteroaromatic N-donor Ligands: Synthesis, Crystal Structure, and Magnetic Properties. <i>Inorganic Chemistry</i> , 2005, 44, 7794-7801.	1.9	52
143	Synthesis, crystal structure and magnetic properties of two-dimensional malonato-bridged cobalt(ii) and nickel(ii) compounds. <i>CrystEngComm</i> , 2004, 6, 106-111.	1.3	103
144	Cobalt(II)-Copper(II) Bimetallic Chains as a New Class of Single-Chain Magnets. <i>Advanced Materials</i> , 2004, 16, 1597-1600.	11.1	135

#	ARTICLE	IF	CITATIONS
145	1,2,4,5-Benzenetetracarboxylic Acid and 4,4'-Bipyridine as Ligands in Designing Low-Dimensional Coordination Polymers. <i>European Journal of Inorganic Chemistry</i> , 2004, 2004, 3873-3879.	1.0	67
146	Phenylmalonate-Containing Copper(II) Complexes: Synthesis, Crystal Structure and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2004, 2004, 4081-4090.	1.0	57
147	Malonic Acid: A Multi-Modal Bridging Ligand for New Architectures and Properties on Molecule-Based Magnets. <i>ChemInform</i> , 2004, 35, no.	0.1	0
148	Nuclearity Controlled Cyanide-Bridged Bimetallic Cr(III)-Mn(II) Compounds: Synthesis, Crystal Structures, Magnetic Properties and Theoretical Calculations. <i>Chemistry - A European Journal</i> , 2004, 10, 6130-6145.	1.7	94
149	Synthesis and crystal structure of the low-spin iron(II) complex [Fe(bpz) ₃](ClO ₄) ₂ ·H ₂ O (bpz=2,2'-bipyrazine). <i>Inorganica Chimica Acta</i> , 2004, 357, 2609-2614.	1.2	10
150	Synthesis, crystal structures and magnetic properties of single and double cyanide-bridged bimetallic Fe ₂ III Cu ₁ zigzag chains. <i>Dalton Transactions</i> , 2004, , 2836.	1.6	78
151	Preparation and crystal structure of the oxalato-bridged Cr(III)-Ag(I) two-dimensional compound {Ag ₃ (H ₂ O)[Cr(dpa)(ox) ₂] ₃ ·nH ₂ O (dpa = 2,2'-dipyridylamine). <i>CrystEngComm</i> , 2004, 6, 11-18.	1.3	19
152	Self-assembly and magnetic properties of a double-propeller octanuclear copper(II) complex with a meso-helicate-type metallacryptand core. <i>Chemical Communications</i> , 2004, , 920-921.	2.2	28
153	Supramolecular Loop-Chain Network of Pillared Layers via 4,4'-Bipyridine. <i>Crystal Growth and Design</i> , 2004, 4, 57-61.	1.4	36
154	Spin Control in Ladderlike Hexanuclear Copper(II) Complexes with Metallacyclophane Cores. <i>Inorganic Chemistry</i> , 2004, 43, 2768-2770.	1.9	43
155	Magnetic Anisotropy of a High-Spin Octanuclear Nickel(II) Complex with a meso-Helicate Core. <i>Inorganic Chemistry</i> , 2004, 43, 7594-7596.	1.9	41
156	Protonated malonate: the influence of the hydrogen bonds on the magnetic behaviour. <i>CrystEngComm</i> , 2004, 6, 443-450.	1.3	32
157	Hybrid Organic/Inorganic Molecular Materials Formed by Tetrathiafulvalene Radicals and Magnetic Trimeric Clusters of Dimetallic Oxalate-Bridged Complexes: The Series (TTF) ₄ {M(II)(H ₂ O) ₂ [M(III)(ox) ₃] ₂ ·nH ₂ O (M(II) = Mn, Fe, Co, Ni, Cu and Zn; M(III) = Cr and Fe; ox = C ₂ O ₄ ²⁻). <i>European Journal of Inorganic Chemistry</i> , 2003, 2003, 2290-2298.	1.0	48
158	Title is missing!. <i>Angewandte Chemie</i> , 2003, 115, 1521-1524.	1.6	79
159	Cyanide-Bridged Iron(III)-Cobalt(II) Double Zigzag Ferromagnetic Chains: Two New Molecular Magnetic Nanowires. <i>Angewandte Chemie - International Edition</i> , 2003, 42, 1483-1486.	7.2	353
160	Cover Picture: Cyanide-Bridged Iron(III)-Cobalt(II) Double Zigzag Ferromagnetic Chains: Two New Molecular Magnetic Nanowires (<i>Angew. Chem. Int. Ed.</i> 13/2003). <i>Angewandte Chemie - International Edition</i> , 2003, 42, 1427-1427.	7.2	4
161	Malonate-based copper(II) coordination compounds: ferromagnetic coupling controlled by dicarboxylates. <i>Polyhedron</i> , 2003, 22, 2143-2153.	1.0	104
162	Malonic acid: a multi-modal bridging ligand for new architectures and properties on molecule-based magnets. <i>Polyhedron</i> , 2003, 22, 2111-2123.	1.0	80

#	ARTICLE	IF	CITATIONS
163	Formation of High-Dimensional Frameworks through Counterions: Structure of $[\text{Cr}_2(\text{Bipy})_2(\text{H}_2\text{O})_2(\text{H}_2\text{O})_2(\text{NO}_3)_3\text{H}_2\text{O}] \cdot \text{C}_6\text{H}_4\text{-2-NH}_2 \cdot \text{H}_2\text{O}$. <i>Crystal Growth and Design</i> , 2003, 3, 273-277.	1.4	6
164	Ferromagnetic Coupling in the Three-Dimensional Malonato-Bridged Gadolinium(III) Complex $[\text{Gd}_2(\text{mal})_3(\text{H}_2\text{O})_6]$ (H_2mal = Malonic Acid). <i>Inorganic Chemistry</i> , 2003, 42, 5456-5458.	1.9	122
165	High-dimensional malonate-based materials: Synthesis, crystal structures and magnetic properties of $[\text{M}_2(\text{mal})_2(\text{L})(\text{H}_2\text{O})_2]_n \cdot n(\text{H}_2\text{O})$ $\text{M} = \text{Zn}(\text{ii}), \text{Co}(\text{ii})$; H_2mal = malonic acid, L = pyrimidine, pyrazine. <i>CrystEngComm</i> , 2003, 5, 280-284.	1.3	53
166	Crystal structures and magnetic properties of two- and three-dimensional malonato-bridged manganese(ii) complexes. <i>Dalton Transactions</i> , 2003, , 2359-2365.	1.6	69
167	$\{[\text{Cu}(\text{H}_2\text{O})_3][\text{Cu}(\text{phmal})_2]\}_n$: a new two-dimensional copper(ii) complex with intralayer ferromagnetic interactions ($\text{phmal} = \text{phenylmalonate dianion}$). <i>New Journal of Chemistry</i> , 2003, 27, 1557-1562.	1.4	51
168	Combining Coordination Chemistry and Hydrogen Bonds: Synthesis, Crystal Structures and Thermal Behavior of the Complexes $\text{Aqua}(2,2\text{-Bipyridine-}N, N\text{-}(\text{Malonamide-}O, O\text{-}M(\text{II})\text{Dinitrate})$ ($\text{M} = \text{Cu}, \text{Ni}$). <i>Journal of Coordination Chemistry</i> , 2003, 56, 181-191.	0.8	2
169	Design of High-Dimensional Copper(II) Malonate Complexes with Exo-Polydentate N-Donor Ligands. <i>Inorganic Chemistry</i> , 2003, 42, 5938-5948.	1.9	119
170	Structural versatility of the malonate ligand as a tool for crystal engineering in the design of molecular magnets. <i>CrystEngComm</i> , 2002, 4, 522-535.	1.3	136
171	Ferromagnetic coupling in the malonato-bridged copper(ii) chains $[\text{Cu}(\text{Im})_2(\text{mal})]_n$ and $[\text{Cu}(2\text{-Melm})_2(\text{mal})]_n$ ($\text{H}_2\text{mal} = \text{malonic acid}$, $\text{Im} = \text{imidazole}$ and $2\text{-Melm} = 2\text{-methylimidazole}$). <i>New Journal of Chemistry</i> , 2002, 26, 1624-1628.	0.8	2
172	A phase transition in the novel three-dimensional compound $[\text{Eu}_2(\text{mal})_3(\text{H}_2\text{O})_6]$ (H_2mal = malonic acid). <i>Dalton Transactions RSC</i> , 2002, , 3462-3470.	2.3	40
173	The flexibility of molecular components as a suitable tool in designing extended magnetic systems. <i>CrystEngComm</i> , 2002, 4, 440-446.	1.3	59
174	Alternating cationic/anionic layers in the $[\text{M}(\text{H}_2\text{O})_6][\text{Cu}(\text{mal})_2(\text{H}_2\text{O})]$ complexes linked through hydrogen bonds ($\text{M} = \text{Mn}, \text{Co}, \text{Ni}, \text{Cu}$ and Zn ; H_2mal = malonic acid). <i>CrystEngComm</i> , 2002, 4, 631-637.	1.3	64
175	Extended network via hydrogen bond linkages of coordination compounds: synthesis, crystal structure and thermal behaviour of the complexes $[\text{M}(\text{L})_2(\text{NO}_3)_2]$ ($\text{M} = \text{Cu}, \text{Co}$) and $[\text{Ni}(\text{L})_2(\text{H}_2\text{O})_2][(\text{NO}_3)_2]$ ($\text{L} = \text{malonamide}$). <i>Inorganica Chimica Acta</i> , 2002, 328, 169-178.	1.2	23
176	Dimensionally controlled hydrogen-bonded nanostructures: synthesis, structure, thermal and magnetic behaviour of the tris-(chelated)nickel(II) complex $[\text{Ni}(\text{bipy})_3]\text{Cl}_2 \cdot 5.5\text{H}_2\text{O}$ ($\text{bipy} = 2,2\text{-bipyridyl}$). <i>Inorganica Chimica Acta</i> , 2002, 336, 131-136.	1.2	36
177	APREFIS 1.0: a computer program for learning physics. <i>European Journal of Physics</i> , 2001, 22, 205-210.	0.3	1
178	A new coordination mode in oxalate-bridged polymers: molecular and crystal structure of $[\text{Cu}_3(\text{C}_2\text{O}_4)_2(\text{C}_2\text{H}_8\text{N}_2)_4(\text{ClO}_4)_2]_n \cdot 2\text{H}_2\text{O}$. <i>CrystEngComm</i> , 2001, 3, 60-63.	1.3	25
179	Geometric isomeric pairs: synthesis, structure, thermal decomposition and magnetic properties of cis- and trans- $[\text{Cr}(\text{mal})(\text{bpy})(\text{H}_2\text{O})_2][\text{NO}_3]_x \cdot x\text{H}_2\text{O}$ ($\text{H}_2\text{mal} = \text{malonic acid}$, $\text{bpy} = 2,2\text{-bipyridine}$). <i>Inorganica Chimica Acta</i> , 2001, 313, 87-94.	1.2	17
180	A new eight-coordinate complex of manganese(II): synthesis, crystal structure, spectroscopy and magnetic properties of $[\text{Mn}(\text{Hoxam})_2(\text{H}_2\text{O})_4]$ ($\text{H}_2\text{oxam} = \text{oxamic acid}$). <i>Inorganica Chimica Acta</i> , 2001, 315, 120-125.	1.2	14

#	ARTICLE	IF	CITATIONS
181	Crystal structure and magnetic properties of the flexible self-assembled two-dimensional square network complex $[\text{Cu}_2(\text{mal})_2(\text{H}_2\text{O})_2(4,4\text{-bpy})]$ (H_2mal =malonic acid and $4,4\text{-bpy}$ =4,4-bipyridine). <i>Inorganica Chimica Acta</i> , 2001, 318, 159-165.	1.2	132
182	Synthesis, crystal structure and magnetic properties of $[\text{Cu}(\text{bpym})(\text{mal})(\text{H}_2\text{O})] \cdot 6\text{H}_2\text{O}$ and $[\text{Cu}_2(\text{bpym})(\text{mal})_2(\text{H}_2\text{O})_2] \cdot 4\text{H}_2\text{O}$ (bpym =2,2-bipyrimidine, H_2mal =malonic acid). <i>Inorganica Chimica Acta</i> , 2001, 326, 20-26.	1.2	59
183	Synthesis, crystal structure and magnetic properties of the malonato-bridged bimetallic chain $[\text{Mn}(\text{II})\text{Cu}(\text{II})(\text{mal})_2(\text{H}_2\text{O})_4] \cdot 2\text{H}_2\text{O}$. <i>Inorganica Chimica Acta</i> , 2000, 298, 202-208.	1.2	67
184	Synthesis, crystal structure and magnetic properties of the three-dimensional compound $[\text{Na}_2\text{Ni}(\text{mal})_2(\text{H}_2\text{O})_6]_n$ (H_2mal =malonic acid). <i>Inorganica Chimica Acta</i> , 2000, 298, 245-250.	1.2	52
185	Generation of lanthanide coordination polymers with dicarboxylate ligands: synthesis, structure, thermal decomposition and magnetic properties of the two-dimensional triaquatrismalonatodiprasedymium(III) dihydrate $\{[\text{Pr}_2(\text{C}_3\text{H}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}\}$. <i>CrystEngComm</i> , 2000, 2, 169-173.	1.3	21
186	Magnetic Coupling through the Carbon Skeleton of Malonate in Two Polymorphs of $\{[\text{Cu}(\text{bpy})(\text{H}_2\text{O})][\text{Cu}(\text{bpy})(\text{mal})(\text{H}_2\text{O})]\}(\text{ClO}_4)_2$ (H_2mal = Malonic Acid; bpy = 2,2'-Bipyridine). <i>Inorganic Chemistry</i> , 2000, 39, 3845-3852.	1.9	129
187	Ferromagnetism in Malonato-Bridged Copper(II) Complexes. Synthesis, Crystal Structures, and Magnetic Properties of $\{[\text{Cu}(\text{H}_2\text{O})_3][\text{Cu}(\text{mal})_2(\text{H}_2\text{O})]\}_n$ and $\{[\text{Cu}(\text{H}_2\text{O})_4]_2[\text{Cu}(\text{mal})_2(\text{H}_2\text{O})]\}[\text{Cu}(\text{mal})_2(\text{H}_2\text{O})_2]\{[\text{Cu}(\text{H}_2\text{O})_4][\text{Cu}(\text{mal})_2(\text{H}_2\text{O})_2]\}$ (H_2mal = malonic Acid). <i>Inorganic Chemistry</i> , 2000, 39, 1363-1370.	1.9	218
188	Organic-inorganic hybrid solids: control of perhalometallate solid state structures. <i>Dalton Transactions RSC</i> , 2000, , 3897-3905.	2.3	107
189	VALMAP2.0: contour maps using the bond-valence-sum method. <i>Journal of Applied Crystallography</i> , 1999, 32, 341-344.	1.9	15
190	Aquabis(2,2'-bipyridine-N,N')(perchlorato-O)manganese(II) perchlorate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999, 55, 186-188.	0.4	1
191	BariumL-tartrate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999, 55, 740-742.	0.4	7
192	Polymeric aqua-1 $\frac{1}{2}$ -bis[$\frac{1}{4}$ -(R,R)-tartrato-1 $\frac{1}{2}$ O:1, O2:2 $\frac{1}{2}$ O:3, O4]dicadmium(II) trihydrate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999, 55, 710-712.	0.4	8
193	Diaquabis(2,2'-bipyridine-N,N')nickel(II) diperchlorate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999, 55, 1087-1090.	0.4	1
194	4 $\frac{1}{2}$,15-Dihydro-3-dehydrosolstitialin A. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999, 55, 1837-1839.	0.4	6
195	Filic-3-ene, a pentacyclic triterpene from <i>Davallia canariensis</i> . <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1999, 55, 1835-1837.	0.4	2
196	Crystal structure and magnetic properties of the single- $\frac{1}{4}$ -chloro copper(II) chain $[\text{Cu}(\text{bipy})\text{Cl}_2]$ (bipy =2,2'-bipyridine). <i>Inorganica Chimica Acta</i> , 1999, 284, 258-265.	1.2	85
197	Molybdotellurates containing the 1-methyl-, 2-methyl- and 4-methyl imidazolium cations. <i>Transition Metal Chemistry</i> , 1999, 24, 141-151.	0.7	13
198	Tungstotellurates of the imidazolium and 4-methylimidazolium cations. <i>Transition Metal Chemistry</i> , 1999, 24, 686-692.	0.7	26

#	ARTICLE	IF	CITATIONS
199	Cation-controlled formation of $[MCl_4]_n$ chains in $[4,4\text{-H}_2\text{bipy}][MCl_4]$ (M = Mn, Cd): an alternative to the $A_2MCl_4 \cdot 100H_2O$ layer perovskite structure. <i>Chemical Communications</i> , 1999, , 2287-2288.	2.2	71
200	Anderson-Evans type molybdotellurates with diazoliun cations: preparation, structure and properties of $(H_2imz)_6 [TeMo_6O_{24}] \cdot 4H_2O$ and $(H_2pyz)_6 [TeMo_6O_{24}] \cdot Te(OH)_6$. <i>Inorganica Chimica Acta</i> , 1998, 277, 139-150.	1.2	20
201	Structural and dielectric characterization of cadmium tartrate. <i>Journal of Applied Physics</i> , 1998, 84, 5729-5732.	1.1	53
202	Weak Ferromagnetism in Chiral 3-Dimensional Oxalato-Bridged Cobalt(II) Compounds. Crystal Structure of $[Co(bpy)_3][Co_2(ox)_3]ClO_4$. <i>Inorganic Chemistry</i> , 1998, 37, 4131-4135.	1.9	132
203	Different Ground Spin States in Iron(III) Complexes with Quadridentate Schiff Bases: Synthesis, Crystal Structures, and Magnetic Properties. <i>Inorganic Chemistry</i> , 1998, 37, 5102-5108.	1.9	65
204	The paraelectric-incommensurate phase transition and crystal structure of the mixed crystals. <i>Journal of Physics Condensed Matter</i> , 1998, 10, 5245-5253.	0.7	5
205	X-ray diffraction, thermal analysis, and Raman scattering study of K_2BeF_4 and comparison to other members of the A_2SO_4 family with ferroelectric-paraelectric transitions. <i>Physical Review B</i> , 1998, 57, 5122-5125.	1.1	3
206	Study on the mixed crystals. <i>Journal of Physics Condensed Matter</i> , 1997, 9, 2657-2669.	0.7	8
207	Dimer species in dimethyl sulfoxide-water (80% v/v) solution of N,N'-bis(salicylideneimine)-m-phenylenediamine ($H_2sal\text{-}m\text{-phen}$) and similar Schiff bases with Cu(I), Ni(II), Co(II) and Zn(II). Crystal structure of $[Co_2(sal\text{-}m\text{-phen})_2] \cdot CHCl_3$. <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 4327-4334.	1.1	37
208	Preparation and characterization of pyridinium-n-carboxylate trioxochromate (VI) (n=3, 4) and pyridinium-4-carboxylic pyridine-4 carboxylate trioxochromate (VI) hemihydrate. <i>Inorganica Chimica Acta</i> , 1997, 258, 53-63.	1.2	10
209	Spectroscopic properties and crystal structure of an oxovanadium(V) alkoxo complex with the ligand: 5-chloro-o-hydroxyacetophenone-salicylhydrazide. <i>Polyhedron</i> , 1997, 16, 1143-1147.	1.0	15
210	3-Acetamido-4-(cyclooctylamino)nitrobenzene. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1997, 53, 651-654.	0.4	0
211	A Structural Study on the Rochelle Salt. <i>Journal of Solid State Chemistry</i> , 1997, 131, 350-357.	1.4	45
212	Structure and properties of Cu(II)(o-hydroxypropiophenone-salicylhydrazide)(2,2'-bipyridyl) $[Cu(OHP\text{-}SHZ)(bipy)]$ and Cu(II)(o-hydroxypropiophenone-salicylhydrazide)(2,2'-bipyridylamine) monohydrate $[Cu(OHP\text{-}SHZ)(bipyam)]H_2O$. <i>Inorganica Chimica Acta</i> , 1997, 255, 139-148.	1.2	13
213	Dimer complexes of 2,4-toluenediamine-N,N,N',N'-tetraacetic acid (2,4-TDTA) with copper(II), nickel(II), cobalt(II), zinc(II) and manganese(II). Studies in aqueous solution and solid state. X-ray crystal structures of $Na_4[Ni_2(2,4\text{-TDTA})_2] \cdot 15H_2O$ and $Na_4[Cu_2(2,4\text{-TDTA})_2] \cdot 20H_2O$. <i>Inorganica Chimica Acta</i> , 1997, 255, 367-380.	1.2	29
214	Study on the mixed crystals $(NH_4)_2(BeF_4)_{1-x}(SO_4)_x$. <i>Ferroelectrics</i> , 1996, 175, 207-218.	0.3	4
215	Hybrid molecular materials formed by alternating layers of bimetallic oxalate complexes and tetrathiafulvalene molecules: Synthesis, structure, and magnetic properties of	11.1	59
216	Five New Chlorochromates of Organic Bases. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1996, 52, 1441-1448.	0.4	9

#	ARTICLE	IF	CITATIONS
217	Disodium [N,N'-(1,3-Propanediyl)bis(oxamato-N,O)]cuprate(II) Hexahydrate. Acta Crystallographica Section C: Crystal Structure Communications, 1996, 52, 1415-1417.	0.4	1
218	Polymeric (Aqua-1 \hat{c} O)bis[(R,R)-tartrato-1 \hat{c} 2O1,O2:2 \hat{c} 2O3,O4]dimanganese(II) Trihydrate. Acta Crystallographica Section C: Crystal Structure Communications, 1996, 52, 2473-2475.	0.4	9
219	Study on K ₂ SeO ₄ at different temperatures. Ferroelectrics, 1996, 177, 191-199.	0.3	11
220	4,4'-Dimethoxy-2,2'-[dithiobis(o-phenylenenitrilomethylidene)]diphenol. Acta Crystallographica Section C: Crystal Structure Communications, 1995, 51, 1016-1018.	0.4	2
221	Preparation and characterization of the dichromates of the organic bases: 2,4 \hat{c} -bipyridine, 2,2 \hat{c} -dipyridylamine and 4,4 \hat{c} -bipyridine. Polyhedron, 1995, 14, 2907-2917.	1.0	29
222	Synthesis and characterization of the new peroxo complex aquaoxodiperoxoimidazole molybdenum(VI). Inorganica Chimica Acta, 1994, 223, 173-175.	1.2	19
223	Structural and Electrophysiological Analysis of Annexin V Mutants. Journal of Molecular Biology, 1994, 237, 479-499.	2.0	59
224	Crystal structure and bonding of bis-(1H-indole-3-ethylene-3'-methoxy-salicylaldehyde)copper(II) monohydrate. Zeitschrift Fur Kristallographie - Crystalline Materials, 1992, 198, 61-66.	0.4	0
225	Crystal structure and bonding of bis-(1H-indole-3-ethylene-3 \hat{c} -methoxy-salicylaldehyde)copper(II) monohydrate. Zeitschrift F \hat{A} ¼r Kristallographie, 1992, 198, 61-66.	1.1	4
226	A new sesquiterpene from Maytenus boaria (Celastraceae); crystal structure and absolute configuration. Tetrahedron Letters, 1992, 33, 1921-1924.	0.7	15
227	Complexes of copper(II) and nickel(II) with the new ligand N-[2-(3-ethyl-indole)]pyridoxaldehyde. X-ray crystal structure of bis{N-[2-(3-ethyl-indole)]pyridoxaldehyde}nickel(II). Polyhedron, 1992, 11, 2171-2178.	1.0	15
228	Synthesis and structure of tetra-aqua-4-chloro-1,2-phenylenediamine-N,N,N \hat{c} ,N \hat{c} -tetra-acetato-O1,O2,O3,O4,N,N \hat{c} -cobalt(II) disodium; Na ₂ [Co(4-Cl-o-PDTA)] \hat{A} ·4H ₂ O. Polyhedron, 1992, 11, 1965-1971.	1.0	11
229	Diterpenoids from Calceolaria thyrsoflora. Phytochemistry, 1991, 30, 589-592.	1.4	38
230	Preparation of oligonucleotides containing dAICA using an unexpected side-reaction observed on a protected derivative of 2-aza-2 \hat{c} -deoxyinosine.. Tetrahedron, 1991, 47, 8917-8930.	1.0	13
231	Structure of 6-chloro-10-oxabicyclo[5.2.1]decane-2,3-diol. Acta Crystallographica Section C: Crystal Structure Communications, 1990, 46, 1655-1657.	0.4	5
232	Structure of 5-hydroxymethyl-7,7-dimethyl-6-oxabicyclo[3.2.1]octane-1-carboxylic acid. Acta Crystallographica Section C: Crystal Structure Communications, 1990, 46, 1507-1509.	0.4	1
233	Tetraaqua-1 \hat{c} /4-(phenylenediaminetetracetato-O1:O2:O3,O4,N:O5,O6,N1)-dicopper(II) dihydrate. Acta Crystallographica Section C: Crystal Structure Communications, 1990, 46, 1405-1407.	0.4	6
234	Structure of triaqua[N-(5-bromosalicylidene)-L-tryptophanato]nickel(II) dihydrate. Acta Crystallographica Section C: Crystal Structure Communications, 1990, 46, 1414-1416.	0.4	2

#	ARTICLE	IF	CITATIONS
235	(+)-Lepidozene a new bicyclic sesquiterpene from the gorgonia lophogorgia ruberrima. Tetrahedron, 1990, 46, 8237-8242.	1.0	10
236	New metabolites from the marine mollusc siphonaria grisea. Tetrahedron, 1990, 46, 1669-1678.	1.0	44
237	New xenicane diterpenes from the brown algae of dictyotaceae. Tetrahedron, 1990, 46, 6125-6132.	1.0	14
238	Synthesis, characterization and crystal structure of		

#	ARTICLE	IF	CITATIONS
253	As4S5N10â€“A Diaminosulfane Derivative with Two Bicyclic Arsino Sulfur Diimides. <i>Angewandte Chemie International Edition in English</i> , 1987, 26, 82-83.	4.4	7