## André L Simão

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5260799/publications.pdf Version: 2024-02-01



ΔΝΙΟΡΑΘΙ SIMAfo

#	Article	IF	CITATIONS
1	Isolation of Mitochondria from Liver and Extraction of Total RNA and Protein: Analyses of miRNA and Protein Expression. Methods in Molecular Biology, 2021, 2310, 1-15.	0.9	2
2	Impact of aging on primary liver cancer: epidemiology, pathogenesis and therapeutics. Aging, 2021, 13, 23416-23434.	3.1	17
3	RIPK3 acts as a lipid metabolism regulator contributing to inflammation and carcinogenesis in non-alcoholic fatty liver disease. Gut, 2021, 70, 2359-2372.	12.1	56
4	Administration of lactobacillus alleviates experimental NASH by reducing miR-21 in the liver. Journal of Hepatology, 2020, 73, S235.	3.7	0
5	Hepatic RIPK3 signalling differentially modulates lipid metabolism and inflammation in non-alcoholic fatty liver disease. Journal of Hepatology, 2020, 73, S672.	3.7	0
6	The role of RIPK3 in non-alcoholic fatty liver disease: a multi-omics perspective. Journal of Hepatology, 2020, 73, S673.	3.7	0
7	mIR-21 is increased in patients with NASH-associated HCC and contributes to hepatocarcinogenesis in mice with NAFLD. Journal of Hepatology, 2020, 73, S677-S678.	3.7	1
8	Composite targeting of nuclear receptors protects against diet-induced NAFLD. Journal of Hepatology, 2020, 73, S458.	3.7	0
9	Liquid Biopsies in Hepatocellular Carcinoma: Are We Winning?. Journal of Clinical Medicine, 2020, 9, 1541.	2.4	38
10	Extracellular Vesicles in Non-alcoholic Fatty Liver Disease: Key Players in Disease Pathogenesis and Promising Biomarker Tools. , 2020, , 157-180.		0
11	FRI-278-RIP3-dependent signalling exerts divergent effects on liver steatosis and carcinogenesis in experimental non-alcholic fatty liver disease. Journal of Hepatology, 2019, 70, e518.	3.7	0
12	Skeletal muscle miR-34a/SIRT1:AMPK axis is activated in experimental and human non-alcoholic steatohepatitis. Journal of Molecular Medicine, 2019, 97, 1113-1126.	3.9	21
13	PS-003-Activation of the miR-34a/SIRT1:AMPK axis contributes for insulin resistance and mitochondrial dysfunction in the NAFLD muscle. Journal of Hepatology, 2019, 70, e7.	3.7	1
14	miRNA-21 ablation protects against liver injury and necroptosis in cholestasis. Cell Death and Differentiation, 2018, 25, 857-872.	11.2	92
15	miR-21 ablation and obeticholic acid ameliorate nonalcoholic steatohepatitis in mice. Cell Death and Disease, 2017, 8, e2748-e2748.	6.3	78
16	Circulating microRNAs as Potential Biomarkers in Non-Alcoholic Fatty Liver Disease and Hepatocellular Carcinoma. Journal of Clinical Medicine, 2016, 5, 30.	2.4	77
17	Activation of necroptosis in human and experimental cholestasis. Cell Death and Disease, 2016, 7, e2390-e2390.	6.3	107
18	Inhibition of NF-κB by deoxycholic acid induces miR-21/PDCD4-dependent hepatocellular apoptosis. Scientific Reports, 2015, 5, 17528.	3.3	24

#	Article	IF	CITATIONS
19	c-Jun N-Terminal Kinase 1/c-Jun Activation of the p53/MicroRNA 34a/Sirtuin 1 Pathway Contributes to Apoptosis Induced by Deoxycholic Acid in Rat Liver. Molecular and Cellular Biology, 2014, 34, 1100-1120.	2.3	61
20	Revisiting the metabolic syndrome and paving the way for micro <scp>RNA</scp> s in nonâ€alcoholic fatty liver disease. FEBS Journal, 2014, 281, 2503-2524.	4.7	55