

Stephan Ruhrmann

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5277935/stephan-ruhrmann-publications-by-citations.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

196 papers	10,794 citations	59 h-index	100 g-index
256 ext. papers	12,880 ext. citations	5.1 avg, IF	6.07 L-index

#	Paper	IF	Citations
196	The psychosis high-risk state: a comprehensive state-of-the-art review. <i>JAMA Psychiatry</i> , 2013 , 70, 107-204.5	204.5	965
195	Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. <i>Archives of General Psychiatry</i> , 2010 , 67, 241-51		491
194	Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. <i>Cell</i> , 2019 , 179, 1469-1482.e11	56.2	402
193	Genome-wide association study of obsessive-compulsive disorder. <i>Molecular Psychiatry</i> , 2013 , 18, 788-9815.1	15.1	244
192	EPA guidance on the early detection of clinical high risk states of psychoses. <i>European Psychiatry</i> , 2015 , 30, 405-16	6	235
191	EPA guidance on the early intervention in clinical high risk states of psychoses. <i>European Psychiatry</i> , 2015 , 30, 388-404	6	233
190	Prediction of psychosis by mismatch negativity. <i>Biological Psychiatry</i> , 2011 , 69, 959-66	7.9	219
189	Intervention in individuals at ultra-high risk for psychosis: a review and future directions. <i>Journal of Clinical Psychiatry</i> , 2009 , 70, 1206-12	4.6	219
188	Revealing the complex genetic architecture of obsessive-compulsive disorder using meta-analysis. <i>Molecular Psychiatry</i> , 2018 , 23, 1181-1188	15.1	205
187	Partitioning the heritability of Tourette syndrome and obsessive compulsive disorder reveals differences in genetic architecture. <i>PLoS Genetics</i> , 2013 , 9, e1003864	6	189
186	Identifying gene-environment interactions in schizophrenia: contemporary challenges for integrated, large-scale investigations. <i>Schizophrenia Bulletin</i> , 2014 , 40, 729-36	1.3	186
185	Sensory gating in schizophrenia: P50 and N100 gating in antipsychotic-free subjects at risk, first-episode, and chronic patients. <i>Biological Psychiatry</i> , 2008 , 64, 376-84	7.9	181
184	Disorder, not just state of risk: meta-analysis of functioning and quality of life in people at high risk of psychosis. <i>British Journal of Psychiatry</i> , 2015 , 207, 198-206	5.4	180
183	Preventing progression to first-episode psychosis in early initial prodromal states. <i>British Journal of Psychiatry</i> , 2012 , 200, 22-9	5.4	162
182	Neurocognitive indicators for a conversion to psychosis: comparison of patients in a potentially initial prodromal state who did or did not convert to a psychosis. <i>Schizophrenia Research</i> , 2007 , 92, 116-23 ⁶	23.6	161
181	Prediction Models of Functional Outcomes for Individuals in the Clinical High-Risk State for Psychosis or With Recent-Onset Depression: A Multimodal, Multisite Machine Learning Analysis. <i>JAMA Psychiatry</i> , 2018 , 75, 1156-1172	14.5	156
180	Basic symptoms and ultrahigh risk criteria: symptom development in the initial prodromal state. <i>Schizophrenia Bulletin</i> , 2010 , 36, 182-91	1.3	152

179	Early detection and secondary prevention of psychosis: facts and visions. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2004 , 254, 117-28	5.1	146
178	Anandamide elevation in cerebrospinal fluid in initial prodromal states of psychosis. <i>British Journal of Psychiatry</i> , 2009 , 194, 371-2	5.4	140
177	Application of transcranial magnetic stimulation in treatment of drug-resistant major depression—report of two cases. <i>Human Psychopharmacology</i> , 1993 , 8, 361-365	2.3	138
176	Neural correlates of working memory dysfunction in first-episode schizophrenia patients: an fMRI multi-center study. <i>Schizophrenia Research</i> , 2007 , 89, 198-210	3.6	130
175	Impaired mismatch negativity generation in prodromal subjects and patients with schizophrenia. <i>Schizophrenia Research</i> , 2005 , 73, 297-310	3.6	130
174	Basic symptoms and the prediction of first-episode psychosis. <i>Current Pharmaceutical Design</i> , 2012 , 18, 351-7	3.3	121
173	Increased neural response related to neutral faces in individuals at risk for psychosis. <i>NeuroImage</i> , 2008 , 40, 289-97	7.9	120
172	Gray matter abnormalities in subjects at ultra-high risk for schizophrenia and first-episode schizophrenic patients compared to healthy controls. <i>Psychiatry Research - Neuroimaging</i> , 2009 , 173, 163-9	2.9	116
171	Probably at-risk, but certainly ill—advocating the introduction of a psychosis spectrum disorder in DSM-V. <i>Schizophrenia Research</i> , 2010 , 120, 23-37	3.6	114
170	Neurocognitive functioning in subjects at risk for a first episode of psychosis compared with first- and multiple-episode schizophrenia. <i>Journal of Clinical and Experimental Neuropsychology</i> , 2006 , 28, 1388-407	2.1	108
169	Neuropsychological profiles in different at-risk states of psychosis: executive control impairment in the early—and additional memory dysfunction in the late—prodromal state. <i>Schizophrenia Bulletin</i> , 2011 , 37, 861-73	1.3	105
168	Cross-disorder genome-wide analyses suggest a complex genetic relationship between Tourette's syndrome and OCD. <i>American Journal of Psychiatry</i> , 2015 , 172, 82-93	11.9	101
167	Disability in people clinically at high risk of psychosis. <i>British Journal of Psychiatry</i> , 2010 , 197, 278-84	5.4	101
166	Acute effects of treatment for prodromal symptoms for people putatively in a late initial prodromal state of psychosis. <i>British Journal of Psychiatry</i> , 2007 , 51, s88-95	5.4	100
165	Impulsiveness in obsessive-compulsive disorder: results from a family study. <i>Acta Psychiatrica Scandinavica</i> , 2007 , 115, 41-7	6.5	97
164	Early detection and intervention in the initial prodromal phase of schizophrenia. <i>Pharmacopsychiatry</i> , 2003 , 36 Suppl 3, S162-7	2	97
163	The European Prediction of Psychosis Study (EPOS): integrating early recognition and intervention in Europe. <i>World Psychiatry</i> , 2005 , 4, 161-7	14.4	97
162	Improving the clinical prediction of psychosis by combining ultra-high risk criteria and cognitive basic symptoms. <i>Schizophrenia Research</i> , 2014 , 154, 100-6	3.6	94

161	Copy number variation in obsessive-compulsive disorder and tourette syndrome: a cross-disorder study. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2014 , 53, 910-9	7.2	86
160	Axis I diagnoses and transition to psychosis in clinical high-risk patients EPOS project: prospective follow-up of 245 clinical high-risk outpatients in four countries. <i>Schizophrenia Research</i> , 2012 , 138, 192-7 ^{3.6}	3.6	85
159	Subjective quality of life in subjects at risk for a first episode of psychosis: a comparison with first episode schizophrenia patients and healthy controls. <i>Schizophrenia Research</i> , 2005 , 79, 137-43	3.6	81
158	Remembering or knowing: electrophysiological evidence for an episodic memory deficit in schizophrenia. <i>Psychological Medicine</i> , 2002 , 32, 1261-71	6.9	81
157	CSF metabolic and proteomic profiles in patients prodromal for psychosis. <i>PLoS ONE</i> , 2007 , 2, e756	3.7	80
156	Auditory P300 in individuals clinically at risk for psychosis. <i>International Journal of Psychophysiology</i> , 2008 , 70, 192-205	2.9	79
155	O6.5. INVESTIGATING VARIABLES FROM THE NAPLS RISK CALCULATOR FOR PSYCHOSIS IN THE EU-GEI HIGH RISK STUDY. <i>Schizophrenia Bulletin</i> , 2019 , 45, S177-S178	1.3	78
154	S13. DO PATIENTS WITH RECENT-ONSET DEPRESSION DIFFER CLINICALLY AND NEUROBIOLOGICALLY FROM DEPRESSED PATIENTS WITH A CLINICAL HIGH-RISK STATE FOR PSYCHOSIS?. <i>Schizophrenia Bulletin</i> , 2018 , 44, S328-S329	1.3	78
153	S94. PREDICTION OF CANNABIS RELAPSE IN CLINICAL HIGH-RISK INDIVIDUALS AND RECENT ONSET PSYCHOSIS - PRELIMINARY RESULTS FROM THE PRONIA STUDY. <i>Schizophrenia Bulletin</i> , 2020 , 46, S69-S70	1.3	78
152	T239. SINGLE-SUBJECT PREDICTION OF FUNCTIONAL OUTCOMES IN CLINICAL HIGH RISK SUBJECTS USING CLINICAL DATA. <i>Schizophrenia Bulletin</i> , 2018 , 44, S209-S210	1.3	78
151	Forecasting psychosis by event-related potentials-systematic review and specific meta-analysis. <i>Biological Psychiatry</i> , 2015 , 77, 951-8	7.9	76
150	Psychosis prediction: stratification of risk estimation with information-processing and premorbid functioning variables. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1482-90	1.3	75
149	Whither the attenuated psychosis syndrome?. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1130-4	1.3	74
148	Binocular depth inversion as a paradigm of reduced visual information processing in prodromal state, antipsychotic-naïve and treated schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2009 , 259, 195-202	5.1	74
147	Familiarity of obsessive-compulsive disorder in nonclinical and clinical subjects. <i>American Journal of Psychiatry</i> , 2006 , 163, 1986-92	11.9	74
146	Prediction and prevention of schizophrenia: what has been achieved and where to go next?. <i>World Psychiatry</i> , 2011 , 10, 165-74	14.4	73
145	Prevalence and clinical significance of DSM-5-attenuated psychosis syndrome in adolescents and young adults in the general population: the Bern Epidemiological At-Risk (BEAR) study. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1499-508	1.3	70
144	Rationale and first results of developing at-risk (prodromal) criteria for bipolar disorder. <i>Current Pharmaceutical Design</i> , 2012 , 18, 358-75	3.3	65

143	Interrelated neuropsychological and anatomical evidence of hippocampal pathology in the at-risk mental state. <i>Psychological Medicine</i> , 2008 , 38, 843-51	6.9	65
142	Effects of fluoxetine versus bright light in the treatment of seasonal affective disorder. <i>Psychological Medicine</i> , 1998 , 28, 923-33	6.9	65
141	Reduced subjective quality of life in persons at risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2008 , 117, 357-68	6.5	64
140	Cannabis use and age at onset of symptoms in subjects at clinical high risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2012 , 125, 45-53	6.5	61
139	Dimensions of working memory dysfunction in schizophrenia. <i>Schizophrenia Research</i> , 2003 , 62, 259-68	3.6	60
138	Resilience as a multimodal dynamic process. <i>Microbial Biotechnology</i> , 2019 , 13, 725-732	3.3	59
137	Basic symptoms in early psychotic and depressive disorders. <i>British Journal of Psychiatry</i> , 2007 , 51, s31-7	5.4	59
136	The near Babylonian speech confusion in early detection of psychosis. <i>Schizophrenia Bulletin</i> , 2011 , 37, 653-5	1.3	54
135	A stratified model for psychosis prediction in clinical practice. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1533-42	1.3	53
134	A rose is a rose is a rose, but at-risk criteria differ. <i>Psychopathology</i> , 2013 , 46, 75-87	3.4	51
133	Neurophysiological correlates of impaired facial affect recognition in individuals at risk for schizophrenia. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1021-9	1.3	51
132	Interventions in the initial prodromal states of psychosis in Germany: concept and recruitment. <i>British Journal of Psychiatry</i> , 2005 , 48, s45-8	5.4	50
131	Harm avoidance in subjects with obsessive-compulsive disorder and their families. <i>Journal of Affective Disorders</i> , 2008 , 107, 265-9	6.6	48
130	Randomized controlled multicentre trial of cognitive behaviour therapy in the early initial prodromal state: effects on social adjustment post treatment. <i>Microbial Biotechnology</i> , 2007 , 1, 71-8	3.3	48
129	Self-reported psychotic-like experiences are a poor estimate of clinician-rated attenuated and frank delusions and hallucinations. <i>Psychopathology</i> , 2014 , 47, 194-201	3.4	47
128	Psychosis-predictive value of self-reported schizotypy in a clinical high-risk sample. <i>Journal of Abnormal Psychology</i> , 2016 , 125, 923-932	7	46
127	No human tryptophan hydroxylase-2 gene R441H mutation in a large cohort of psychiatric patients and control subjects. <i>Biological Psychiatry</i> , 2006 , 60, 202-3	7.9	46
126	Pharmacological intervention in the initial prodromal phase of psychosis. <i>European Psychiatry</i> , 2005 , 20, 1-6	6	45

125	Course of clinical high-risk states for psychosis beyond conversion. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2018 , 268, 39-48	5.1	43
124	Support for the association between the rare functional variant I425V of the serotonin transporter gene and susceptibility to obsessive compulsive disorder. <i>Molecular Psychiatry</i> , 2005 , 10, 1059-61	15.1	43
123	Alexithymia in obsessive-compulsive disorder - results from a family study. <i>Psychotherapy and Psychosomatics</i> , 2006 , 75, 312-8	9.4	42
122	Integrating evolutionary and regulatory information with a multispecies approach implicates genes and pathways in obsessive-compulsive disorder. <i>Nature Communications</i> , 2017 , 8, 774	17.4	41
121	Duration of unspecific prodromal and clinical high risk states, and early help-seeking in first-admission psychosis patients. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015 , 50, 1831-41	4.5	41
120	Revisiting the Basic Symptom Concept: Toward Translating Risk Symptoms for Psychosis into Neurobiological Targets. <i>Frontiers in Psychiatry</i> , 2016 , 7, 9	5	40
119	Low plasma thyroid indices of depressed patients are attenuated by antidepressant drugs and influence treatment outcome. <i>Pharmacopsychiatry</i> , 1996 , 29, 180-6	2	39
118	Can quantitative EEG measures predict clinical outcome in subjects at Clinical High Risk for psychosis? A prospective multicenter study. <i>Schizophrenia Research</i> , 2014 , 153, 42-7	3.6	38
117	Antisaccade performance in patients with obsessive-compulsive disorder and unaffected relatives: further evidence for impaired response inhibition as a candidate endophenotype. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2012 , 262, 625-34	5.1	38
116	Characterization of SLITRK1 variation in obsessive-compulsive disorder. <i>PLoS ONE</i> , 2013 , 8, e70376	3.7	37
115	Kraepelin and psychotic prodromal conditions. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2008 , 258 Suppl 2, 74-84	5.1	37
114	Prediction of psychosis in clinical high-risk patients by the Schizotypal Personality Questionnaire. Results of the EPOS project. <i>European Psychiatry</i> , 2013 , 28, 469-75	6	36
113	Early onset of obsessive-compulsive disorder and associated comorbidity. <i>Depression and Anxiety</i> , 2009 , 26, 1012-7	8.4	36
112	Relationship between subjective and objective cognitive function in the early and late prodrome. <i>British Journal of Psychiatry</i> , 2007 , 51, s43-51	5.4	36
111	Early detection of psychosis - establishing a service for persons at risk. <i>European Psychiatry</i> , 2009 , 24, 1-10	6	35
110	Personality disorders and accentuations in at-risk persons with and without conversion to first-episode psychosis. <i>Microbial Biotechnology</i> , 2012 , 6, 389-98	3.3	34
109	A promoter variant of SHANK1 affects auditory working memory in schizophrenia patients and in subjects clinically at risk for psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2012 , 262, 117-24	5.1	33
108	Prevalence and clinical relevance of interview-assessed psychosis-risk symptoms in the young adult community. <i>Psychological Medicine</i> , 2018 , 48, 1167-1178	6.9	32

107	The initial prodrome of schizophrenia: different duration, different underlying deficits?. <i>Comprehensive Psychiatry</i> , 2007 , 48, 479-88	7.3	31
106	Multimodal Machine Learning Workflows for Prediction of Psychosis in Patients With Clinical High-Risk Syndromes and Recent-Onset Depression. <i>JAMA Psychiatry</i> , 2021 , 78, 195-209	14.5	31
105	Perceived parental rearing in subjects with obsessive-compulsive disorder and their siblings. <i>Acta Psychiatrica Scandinavica</i> , 2010 , 121, 280-8	6.5	30
104	The interaction of working memory and emotion in persons clinically at risk for psychosis: an fMRI pilot study. <i>Schizophrenia Research</i> , 2010 , 120, 167-76	3.6	29
103	DAOA/G72 predicts the progression of prodromal syndromes to first episode psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2010 , 260, 209-15	5.1	29
102	Obsessive-compulsive disorder and posttraumatic stress disorder. <i>Psychopathology</i> , 2008 , 41, 129-34	3.4	28
101	Predictors for symptom re-exacerbation after targeted stepwise drug discontinuation in first-episode schizophrenia: Results of the first-episode study within the German research network on schizophrenia. <i>Schizophrenia Research</i> , 2016 , 170, 168-76	3.6	27
100	Psychosocial outcome in patients at clinical high risk of psychosis: a prospective follow-up. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013 , 48, 303-11	4.5	27
99	Orienting of attention in unmedicated patients with schizophrenia, prodromal subjects and healthy relatives. <i>Schizophrenia Research</i> , 2007 , 97, 35-42	3.6	27
98	Child Maltreatment and Clinical Outcome in Individuals at Ultra-High Risk for Psychosis in the EU-GEI High Risk Study. <i>Schizophrenia Bulletin</i> , 2018 , 44, 584-592	1.3	25
97	Recurrent brief depression and its relationship to seasonal affective disorder. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1992 , 242, 20-6	5.1	25
96	Sex differences in the genetic architecture of obsessive-compulsive disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2019 , 180, 351-364	3.5	25
95	Short-term functional outcome and premorbid adjustment in clinical high-risk patients. Results of the EPOS project. <i>European Psychiatry</i> , 2014 , 29, 371-80	6	24
94	The Strauss and Carpenter Prognostic Scale in subjects clinically at high risk of psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2013 , 127, 53-61	6.5	23
93	Intervention in the at-risk state to prevent transition to psychosis. <i>Current Opinion in Psychiatry</i> , 2009 , 22, 177-83	4.9	22
92	Restricted attention to social cues in schizophrenia patients. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2016 , 266, 649-61	5.1	21
91	Pathways to care in subjects at high risk for psychotic disorders - a European perspective. <i>Schizophrenia Research</i> , 2014 , 152, 400-7	3.6	21
90	Morphometry of structural disconnectivity indicators in subjects at risk and in age-matched patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2013 , 263, 15-24	5.1	20

89	Efficacy of flupentixol and risperidone in chronic schizophrenia with predominantly negative symptoms. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2007 , 31, 1012-22	5.5	20
88	The interrelationship between schizotypy, clinical high risk for psychosis and related symptoms: Cognitive disturbances matter. <i>Schizophrenia Research</i> , 2019 , 210, 188-196	3.6	19
87	Development of Proteomic Prediction Models for Transition to Psychotic Disorder in the Clinical High-Risk State and Psychotic Experiences in Adolescence. <i>JAMA Psychiatry</i> , 2021 , 78, 77-90	14.5	19
86	Individualized Prediction of Transition to Psychosis in 1,676 Individuals at Clinical High Risk: Development and Validation of a Multivariable Prediction Model Based on Individual Patient Data Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2019 , 10, 345	5	18
85	5-HT3 receptor influences the washing phenotype and visual organization in obsessive-compulsive disorder supporting 5-HT3 receptor antagonists as novel treatment option. <i>European Neuropsychopharmacology</i> , 2014 , 24, 86-94	1.2	18
84	Depression predicts persistence of paranoia in clinical high-risk patients to psychosis: results of the EPOS project. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016 , 51, 247-57	4.5	17
83	Prediction and prevention of psychosis: current progress and future tasks. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2014 , 264 Suppl 1, S9-16	5.1	16
82	Pharmacological prevention and treatment in clinical at-risk states for psychosis. <i>Current Pharmaceutical Design</i> , 2012 , 18, 550-7	3.3	16
81	Chances and risks of predicting psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2012 , 262 Suppl 2, S85-90	5.1	16
80	Gender differences of patients at-risk for psychosis regarding symptomatology, drug use, comorbidity and functioning - Results from the EU-GEI study. <i>European Psychiatry</i> , 2019 , 59, 52-59	6	15
79	Traces of Trauma: A Multivariate Pattern Analysis of Childhood Trauma, Brain Structure, and Clinical Phenotypes. <i>Biological Psychiatry</i> , 2020 , 88, 829-842	7.9	15
78	Atypical processing of uncertainty in individuals at risk for psychosis. <i>NeuroImage: Clinical</i> , 2020 , 26, 102239	3.9	14
77	A controlled study of the efficacy and safety of mianserin and amitriptyline in depressive inpatients. <i>Pharmacopsychiatry</i> , 1995 , 28, 249-52	2	14
76	Association of Adverse Outcomes With Emotion Processing and Its Neural Substrate in Individuals at Clinical High Risk for Psychosis. <i>JAMA Psychiatry</i> , 2020 , 77, 190-200	14.5	14
75	Comparing the prodrome of schizophrenia-spectrum psychoses and affective disorders with and without psychotic features. <i>Schizophrenia Research</i> , 2012 , 138, 218-22	3.6	13
74	Intervention in at-risk states for developing psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2010 , 260 Suppl 2, S90-4	5.1	13
73	Search for copy number variants in chromosomes 15q11-q13 and 22q11.2 in obsessive compulsive disorder. <i>BMC Medical Genetics</i> , 2010 , 11, 100	2.1	13
72	The German Research Network on Schizophrenia--impact on the management of schizophrenia. <i>Dialogues in Clinical Neuroscience</i> , 2006 , 8, 115-21	5.7	13

71	Towards clinical application of prediction models for transition to psychosis: A systematic review and external validation study in the PRONIA sample. <i>Neuroscience and Biobehavioral Reviews</i> , 2021 , 125, 478-492	9	13
70	Antisaccade performance is related to genetic loading for schizophrenia. <i>Journal of Psychiatric Research</i> , 2009 , 43, 291-7	5.2	12
69	Development of a stage-dependent prognostic model to predict psychosis in ultra-high-risk patients seeking treatment for co-morbid psychiatric disorders. <i>Psychological Medicine</i> , 2016 , 46, 1839-51	6.9	12
68	Age effects on basic symptoms in the community: A route to gain new insight into the neurodevelopment of psychosis?. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2020 , 270, 311-324	5.1	11
67	Neurocognitive functioning in parents of schizophrenia patients: Attentional and executive performance vary with genetic loading. <i>Psychiatry Research</i> , 2015 , 230, 885-91	9.9	10
66	Perceived negative attitude of others predicts transition to psychosis in patients at risk of psychosis. <i>European Psychiatry</i> , 2012 , 27, 264-6	6	10
65	Antisaccade and prosaccade eye movements in individuals clinically at risk for psychosis: comparison with first-episode schizophrenia and prediction of conversion. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2019 , 269, 921-930	5.1	10
64	Dysregulated Lipid Metabolism Precedes Onset of Psychosis. <i>Biological Psychiatry</i> , 2021 , 89, 288-297	7.9	10
63	Genome-wide association study of pediatric obsessive-compulsive traits: shared genetic risk between traits and disorder. <i>Translational Psychiatry</i> , 2021 , 11, 91	8.6	10
62	Expressed emotion as a predictor of the first psychotic episode - Results of the European prediction of psychosis study. <i>Schizophrenia Research</i> , 2018 , 199, 346-352	3.6	9
61	Perceived negative attitude of others as an early sign of psychosis. <i>European Psychiatry</i> , 2009 , 24, 233-8	6	9
60	The Psychopathology and Neuroanatomical Markers of Depression in Early Psychosis. <i>Schizophrenia Bulletin</i> , 2021 , 47, 249-258	1.3	9
59	Multimodal prevention of first psychotic episode through N-acetyl-l-cysteine and integrated preventive psychological intervention in individuals clinically at high risk for psychosis: Protocol of a randomized, placebo-controlled, parallel-group trial. <i>Microbial Biotechnology</i> , 2019 , 13, 1404-1415	3.3	8
58	Sex differences in cognitive functioning of patients at-risk for psychosis and healthy controls: Results from the European Gene-Environment Interactions study. <i>European Psychiatry</i> , 2020 , 63, e25	6	8
57	General psychopathology links burden of recent life events and psychotic symptoms in a network approach. <i>NPJ Schizophrenia</i> , 2020 , 6, 40	5.5	8
56	Toward Generalizable and Transdiagnostic Tools for Psychosis Prediction: An Independent Validation and Improvement of the NAPLS-2 Risk Calculator in the Multisite PRONIA Cohort. <i>Biological Psychiatry</i> , 2021 , 90, 632-642	7.9	8
55	Emotion Recognition and Adverse Childhood Experiences in Individuals at Clinical High Risk of Psychosis. <i>Schizophrenia Bulletin</i> , 2020 , 46, 823-833	1.3	7
54	Childhood adversity predicts persistence of suicidal thoughts differently in females and males at clinical high-risk patients of psychosis. Results of the EPOS project. <i>Microbial Biotechnology</i> , 2019 , 13, 935-942	3.3	7

53	Clinical response to sleep deprivation and auditory-evoked potentials--preliminary results. <i>Pharmacopsychiatry</i> , 1994 , 27, 70-1	2	7
52	Clinical, cognitive and neuroanatomical associations of serum NMDAR autoantibodies in people at clinical high risk for psychosis. <i>Molecular Psychiatry</i> , 2021 , 26, 2590-2604	15.1	7
51	Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings. <i>Molecular Psychiatry</i> , 2021 , 26, 4529-4543	15.1	7
50	Personality dimensions in persons symptomatically at risk of psychosis: pronounced but lacking a characteristic profile. <i>Microbial Biotechnology</i> , 2015 , 9, 242-7	3.3	6
49	The functional coding variant Asn107Ile of the neuropeptide S receptor gene (NPSR1) influences age at onset of obsessive-compulsive disorder. <i>International Journal of Neuropsychopharmacology</i> , 2013 , 16, 1951-8	5.8	6
48	Evidence for a seasonal form of recurrent brief depression (RBD-seasonal). <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1994 , 244, 205-10	5.1	6
47	Basic disturbances of information processing in psychosis prediction. <i>Frontiers in Psychiatry</i> , 2013 , 4, 93	5	5
46	Pre-training inter-rater reliability of clinical instruments in an international psychosis research project. <i>Schizophrenia Research</i> , 2021 , 230, 104-107	3.6	5
45	Association between age of cannabis initiation and gray matter covariance networks in recent onset psychosis. <i>Neuropsychopharmacology</i> , 2021 , 46, 1484-1493	8.7	5
44	Sex differences in symptomatology of psychosis-risk patients and in prediction of psychosis. <i>Archives of Women's Mental Health</i> , 2020 , 23, 339-349	5	5
43	Health literacy in clinical-high-risk individuals for psychosis: A systematic mixed-methods review. <i>Microbial Biotechnology</i> , 2019 , 13, 1293-1309	3.3	4
42	Validation of the Bullying Scale for Adults - Results of the PRONIA-study. <i>Journal of Psychiatric Research</i> , 2020 , 129, 88-97	5.2	4
41	Relationship between jumping to conclusions and clinical outcomes in people at clinical high-risk for psychosis. <i>Psychological Medicine</i> , 2020 , 1-9	6.9	4
40	Peculiarities of health literacy in people with mental disorders: A cross-sectional study. <i>International Journal of Social Psychiatry</i> , 2020 , 66, 10-22	8.5	4
39	Physical illnesses, developmental risk factors and psychiatric diagnoses among subjects at risk of psychosis. <i>European Psychiatry</i> , 2013 , 28, 135-40	6	3
38	Validierung einer deutschen Version der Resilience Scale for Adults (RSA). <i>Diagnostica</i> , 2019 , 65, 205-215	5.8	3
37	Multimodal prognosis of negative symptom severity in individuals at increased risk of developing psychosis. <i>Translational Psychiatry</i> , 2021 , 11, 312	8.6	3
36	Heterogeneity and Classification of Recent Onset Psychosis and Depression: A Multimodal Machine Learning Approach. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1130-1140	1.3	3

35	Basic Symptoms Are Associated With Age in Patients With a Clinical High-Risk State for Psychosis: Results From the PRONIA Study. <i>Frontiers in Psychiatry</i> , 2020 , 11, 552175	5	2
34	A randomized Phase II trial evaluating efficacy, safety, and tolerability of oral BI 409306 in attenuated psychosis syndrome: Design and rationale. <i>Microbial Biotechnology</i> , 2021 , 15, 1315-1325	3.3	2
33	Prediction and Early Detection of First-Episode Psychosis 2011 , 207-267		2
32	Cognitive subtypes in recent onset psychosis: distinct neurobiological fingerprints?. <i>Neuropsychopharmacology</i> , 2021 , 46, 1475-1483	8.7	2
31	Obsessive-Compulsive Symptoms and Other Symptoms of the At-risk Mental State for Psychosis: A Network Perspective. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1018-1028	1.3	2
30	Impact of Comorbid Affective Disorders on Longitudinal Clinical Outcomes in Individuals at Ultra-high Risk for Psychosis. <i>Schizophrenia Bulletin</i> , 2021 ,	1.3	2
29	PREVENTIVE PSYCHOTHERAPY. <i>Schizophrenia Research</i> , 2014 , 153, S42-S43	3.6	1
28	Clinical High Risk Symptoms and Criteria in the Community: Prevalence, Clinical Significance and Risk Factors for Their Occurrence. <i>European Psychiatry</i> , 2017 , 41, S226-S226	6	1
27	Basic Symptoms in the Community and their Association with Age. <i>European Psychiatry</i> , 2017 , 41, S84-S86		1
26	Dysregulated lipid metabolism precedes onset of psychosis in people at clinical high risk		1
25	The non-specific nature of mental health and structural brain outcomes following childhood trauma. <i>Psychological Medicine</i> , 2021 , 1-10	6.9	1
24	Novel Gyrfication Networks Reveal Links with Psychiatric Risk Factors in Early Illness. <i>Cerebral Cortex</i> , 2021 ,	5.1	1
23	Impact of smoking Behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study. <i>European Psychiatry</i> , 2021 , 64, e60	6	1
22	Association between tobacco use and symptomatology in individuals at ultra-high risk to develop a psychosis: A longitudinal study. <i>Schizophrenia Research</i> , 2021 , 236, 48-53	3.6	1
21	Attitudes towards Risk Prediction in a Help Seeking Population of Early Detection Centers for Mental Disorders-A Qualitative Approach. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	1
20	Relationships between global functioning and neuropsychological predictors in subjects at high risk of psychosis or with a recent onset of depression.. <i>World Journal of Biological Psychiatry</i> , 2022 , 1-9	3.8	0
19	Verbal memory performance predicts remission and functional outcome in people at clinical high-risk for psychosis.. <i>Schizophrenia Research: Cognition</i> , 2022 , 28, 100222	2.8	0
18	The clinical relevance of formal thought disorder in the early stages of psychosis: results from the PRONIA study. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021 , 1	5.1	0

- 17 Using combined environmental-clinical classification models to predict role functioning outcome in clinical high-risk states for psychosis and recent-onset depression.. *British Journal of Psychiatry*, **2022**, 1-17 5.4 ○
- 16 Electroencephalographic Predictors of Psychosis. *Key Issues in Mental Health*, 133-141
- 15 Intervention in Clinical High Risk States - Current Status and Future Perspectives. *European Psychiatry*, **2017**, 41, S27-S28 6
- 14 Poster #202 COMPARING THE PRODROME OF SCHIZOPHRENIA-SPECTRUM PSYCHOSES AND BIPOLAR DISORDERS WITH AND WITHOUT PSYCHOTIC FEATURES. *Schizophrenia Research*, **2012**, 136, S163-S164 3.6
- 13 Subjective quality of life and its changes in patients at-risk for psychosis results of the EPOS study. *European Psychiatry*, **2011**, 26, 2039-2039 6
- 12 Course of psychopathology in the at-risk mental state - outcomes beyond transition to psychosis. *European Psychiatry*, **2011**, 26, 2037-2037 6
- 11 The EPOS prediction model improves ability to predict transition to first episode psychosis in individuals at high risk. *Evidence-Based Mental Health*, **2010**, 13, 77 11.1
- 10 Klinische Prädiktion und Prävention von Psychosen. *Psychiatrie Und Psychotherapie Up2date*, **2010**, 4, 237-252
- 9 PHARMACOLOGICAL INTERVENTION IN THE PRODRAMAL STATE OF PSYCHOSIS. *Schizophrenia Research*, **2008**, 102, 47-48 3.6
- 8 P.3.f.001 Does anandamide elevation in cerebrospinal fluid protect against transition into frank psychosis?. *European Neuropsychopharmacology*, **2008**, 18, S474 1.2
- 7 Detailed clinical phenotyping and generalisability in prognostic models of functioning in at-risk populations.. *British Journal of Psychiatry*, **2021**, 1-4 5.4
- 6 Intervención farmacológica en la fase prodromática inicial de la psicosis. *European Psychiatry (Ed Española)*, **2005**, 12, 217-222
- 5 Prevention and Early Intervention in At-Risk States for Developing Psychosis **2011**, 81-92
- 4 Is there a diagnosis-specific influence of childhood trauma on later educational attainment? A machine learning analysis in a large help-seeking sample. *Journal of Psychiatric Research*, **2021**, 138, 591-597 5.3
- 3 AuthorsReply. *British Journal of Psychiatry*, **2016**, 208, 197-8 5.4
- 2 Impact of smoking behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study - CORRIGENDUM.. *European Psychiatry*, **2022**, 65, e17 6
- 1 Pattern of predictive features of continued cannabis use in patients with recent-onset psychosis and clinical high-risk for psychosis.. *NPJ Schizophrenia*, **2022**, 8, 19 5.5