

Stephan Ruhrmann

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5277935/stephan-ruhrmann-publications-by-year.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

196 papers	10,794 citations	59 h-index	100 g-index
256 ext. papers	12,880 ext. citations	5.1 avg, IF	6.07 L-index

#	Paper	IF	Citations
196	Relationships between global functioning and neuropsychological predictors in subjects at high risk of psychosis or with a recent onset of depression.. <i>World Journal of Biological Psychiatry</i> , 2022 , 1-9	3.8	0
195	Verbal memory performance predicts remission and functional outcome in people at clinical high-risk for psychosis.. <i>Schizophrenia Research: Cognition</i> , 2022 , 28, 100222	2.8	0
194	Impact of smoking behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study - CORRIGENDUM.. <i>European Psychiatry</i> , 2022 , 65, e17	6	
193	Using combined environmental-clinical classification models to predict role functioning outcome in clinical high-risk states for psychosis and recent-onset depression.. <i>British Journal of Psychiatry</i> , 2022 , 1-17	5.4	0
192	Pattern of predictive features of continued cannabis use in patients with recent-onset psychosis and clinical high-risk for psychosis.. <i>NPJ Schizophrenia</i> , 2022 , 8, 19	5.5	
191	Detailed clinical phenotyping and generalisability in prognostic models of functioning in at-risk populations.. <i>British Journal of Psychiatry</i> , 2021 , 1-4	5.4	
190	A randomized Phase II trial evaluating efficacy, safety, and tolerability of oral BI 409306 in attenuated psychosis syndrome: Design and rationale. <i>Microbial Biotechnology</i> , 2021 , 15, 1315-1325	3.3	2
189	Clinical, cognitive and neuroanatomical associations of serum NMDAR autoantibodies in people at clinical high risk for psychosis. <i>Molecular Psychiatry</i> , 2021 , 26, 2590-2604	15.1	7
188	Pre-training inter-rater reliability of clinical instruments in an international psychosis research project. <i>Schizophrenia Research</i> , 2021 , 230, 104-107	3.6	5
187	Association between age of cannabis initiation and gray matter covariance networks in recent onset psychosis. <i>Neuropsychopharmacology</i> , 2021 , 46, 1484-1493	8.7	5
186	Cognitive subtypes in recent onset psychosis: distinct neurobiological fingerprints?. <i>Neuropsychopharmacology</i> , 2021 , 46, 1475-1483	8.7	2
185	Multimodal prognosis of negative symptom severity in individuals at increased risk of developing psychosis. <i>Translational Psychiatry</i> , 2021 , 11, 312	8.6	3
184	Is there a diagnosis-specific influence of childhood trauma on later educational attainment? A machine learning analysis in a large help-seeking sample. <i>Journal of Psychiatric Research</i> , 2021 , 138, 591-597	5.2	7
183	Towards clinical application of prediction models for transition to psychosis: A systematic review and external validation study in the PRONIA sample. <i>Neuroscience and Biobehavioral Reviews</i> , 2021 , 125, 478-492	9	13
182	Toward Generalizable and Transdiagnostic Tools for Psychosis Prediction: An Independent Validation and Improvement of the NAPLS-2 Risk Calculator in the Multisite PRONIA Cohort. <i>Biological Psychiatry</i> , 2021 , 90, 632-642	7.9	8
181	The Psychopathology and Neuroanatomical Markers of Depression in Early Psychosis. <i>Schizophrenia Bulletin</i> , 2021 , 47, 249-258	1.3	9
180	Dysregulated Lipid Metabolism Precedes Onset of Psychosis. <i>Biological Psychiatry</i> , 2021 , 89, 288-297	7.9	10

179	Development of Proteomic Prediction Models for Transition to Psychotic Disorder in the Clinical High-Risk State and Psychotic Experiences in Adolescence. <i>JAMA Psychiatry</i> , 2021 , 78, 77-90	14.5	19
178	Genome-wide association study of pediatric obsessive-compulsive traits: shared genetic risk between traits and disorder. <i>Translational Psychiatry</i> , 2021 , 11, 91	8.6	10
177	Heterogeneity and Classification of Recent Onset Psychosis and Depression: A Multimodal Machine Learning Approach. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1130-1140	1.3	3
176	Multimodal Machine Learning Workflows for Prediction of Psychosis in Patients With Clinical High-Risk Syndromes and Recent-Onset Depression. <i>JAMA Psychiatry</i> , 2021 , 78, 195-209	14.5	31
175	Obsessive-Compulsive Symptoms and Other Symptoms of the At-risk Mental State for Psychosis: A Network Perspective. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1018-1028	1.3	2
174	The non-specific nature of mental health and structural brain outcomes following childhood trauma. <i>Psychological Medicine</i> , 2021 , 1-10	6.9	1
173	Impact of Comorbid Affective Disorders on Longitudinal Clinical Outcomes in Individuals at Ultra-high Risk for Psychosis. <i>Schizophrenia Bulletin</i> , 2021 ,	1.3	2
172	Novel Cytification Networks Reveal Links with Psychiatric Risk Factors in Early Illness. <i>Cerebral Cortex</i> , 2021 ,	5.1	1
171	Impact of smoking Behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study. <i>European Psychiatry</i> , 2021 , 64, e60	6	1
170	The clinical relevance of formal thought disorder in the early stages of psychosis: results from the PRONIA study. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021 , 1	5.1	0
169	Association between tobacco use and symptomatology in individuals at ultra-high risk to develop a psychosis: A longitudinal study. <i>Schizophrenia Research</i> , 2021 , 236, 48-53	3.6	1
168	Attitudes towards Risk Prediction in a Help Seeking Population of Early Detection Centers for Mental Disorders-A Qualitative Approach. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	1
167	Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings. <i>Molecular Psychiatry</i> , 2021 , 26, 4529-4543	15.1	7
166	Traces of Trauma: A Multivariate Pattern Analysis of Childhood Trauma, Brain Structure, and Clinical Phenotypes. <i>Biological Psychiatry</i> , 2020 , 88, 829-842	7.9	15
165	Atypical processing of uncertainty in individuals at risk for psychosis. <i>NeuroImage: Clinical</i> , 2020 , 26, 102339	3.9	14
164	Sex differences in cognitive functioning of patients at-risk for psychosis and healthy controls: Results from the European Gene-Environment Interactions study. <i>European Psychiatry</i> , 2020 , 63, e25	6	8
163	Emotion Recognition and Adverse Childhood Experiences in Individuals at Clinical High Risk of Psychosis. <i>Schizophrenia Bulletin</i> , 2020 , 46, 823-833	1.3	7
162	Basic Symptoms Are Associated With Age in Patients With a Clinical High-Risk State for Psychosis: Results From the PRONIA Study. <i>Frontiers in Psychiatry</i> , 2020 , 11, 552175	5	2

161	Validation of the Bullying Scale for Adults - Results of the PRONIA-study. <i>Journal of Psychiatric Research</i> , 2020 , 129, 88-97	5.2	4
160	General psychopathology links burden of recent life events and psychotic symptoms in a network approach. <i>NPJ Schizophrenia</i> , 2020 , 6, 40	5.5	8
159	Association of Adverse Outcomes With Emotion Processing and Its Neural Substrate in Individuals at Clinical High Risk for Psychosis. <i>JAMA Psychiatry</i> , 2020 , 77, 190-200	14.5	14
158	S94. PREDICTION OF CANNABIS RELAPSE IN CLINICAL HIGH-RISK INDIVIDUALS AND RECENT ONSET PSYCHOSIS - PRELIMINARY RESULTS FROM THE PRONIA STUDY. <i>Schizophrenia Bulletin</i> , 2020 , 46, S69-S70	1.3	78
157	Relationship between jumping to conclusions and clinical outcomes in people at clinical high-risk for psychosis. <i>Psychological Medicine</i> , 2020 , 1-9	6.9	4
156	Age effects on basic symptoms in the community: A route to gain new insight into the neurodevelopment of psychosis?. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2020 , 270, 311-324	5.1	11
155	Peculiarities of health literacy in people with mental disorders: A cross-sectional study. <i>International Journal of Social Psychiatry</i> , 2020 , 66, 10-22	8.5	4
154	Sex differences in symptomatology of psychosis-risk patients and in prediction of psychosis. <i>Archives of Women's Mental Health</i> , 2020 , 23, 339-349	5	5
153	The interrelationship between schizotypy, clinical high risk for psychosis and related symptoms: Cognitive disturbances matter. <i>Schizophrenia Research</i> , 2019 , 210, 188-196	3.6	19
152	Health literacy in clinical-high-risk individuals for psychosis: A systematic mixed-methods review. <i>Microbial Biotechnology</i> , 2019 , 13, 1293-1309	3.3	4
151	Individualized Prediction of Transition to Psychosis in 1,676 Individuals at Clinical High Risk: Development and Validation of a Multivariable Prediction Model Based on Individual Patient Data Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2019 , 10, 345	5	18
150	Gender differences of patients at-risk for psychosis regarding symptomatology, drug use, comorbidity and functioning - Results from the EU-GEI study. <i>European Psychiatry</i> , 2019 , 59, 52-59	6	15
149	Multimodal prevention of first psychotic episode through N-acetyl-L-cysteine and integrated preventive psychological intervention in individuals clinically at high risk for psychosis: Protocol of a randomized, placebo-controlled, parallel-group trial. <i>Microbial Biotechnology</i> , 2019 , 13, 1404-1415	3.3	8
148	O6.5. INVESTIGATING VARIABLES FROM THE NAPLS RISK CALCULATOR FOR PSYCHOSIS IN THE EU-GEI HIGH RISK STUDY. <i>Schizophrenia Bulletin</i> , 2019 , 45, S177-S178	1.3	78
147	Childhood adversity predicts persistence of suicidal thoughts differently in females and males at clinical high-risk patients of psychosis. Results of the EPOS project. <i>Microbial Biotechnology</i> , 2019 , 13, 935-942	3.3	7
146	Resilience as a multimodal dynamic process. <i>Microbial Biotechnology</i> , 2019 , 13, 725-732	3.3	59
145	Validierung einer deutschen Version der Resilience Scale for Adults (RSA). <i>Diagnostica</i> , 2019 , 65, 205-215.	5.8	3
144	Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. <i>Cell</i> , 2019 , 179, 1469-1482.e11	56.2	402

143	Sex differences in the genetic architecture of obsessive-compulsive disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2019 , 180, 351-364	3.5	25
142	Antisaccade and prosaccade eye movements in individuals clinically at risk for psychosis: comparison with first-episode schizophrenia and prediction of conversion. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2019 , 269, 921-930	5.1	10
141	Expressed emotion as a predictor of the first psychotic episode - Results of the European prediction of psychosis study. <i>Schizophrenia Research</i> , 2018 , 199, 346-352	3.6	9
140	Course of clinical high-risk states for psychosis beyond conversion. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2018 , 268, 39-48	5.1	43
139	Child Maltreatment and Clinical Outcome in Individuals at Ultra-High Risk for Psychosis in the EU-GEI High Risk Study. <i>Schizophrenia Bulletin</i> , 2018 , 44, 584-592	1.3	25
138	Prevalence and clinical relevance of interview-assessed psychosis-risk symptoms in the young adult community. <i>Psychological Medicine</i> , 2018 , 48, 1167-1178	6.9	32
137	S13. DO PATIENTS WITH RECENT-ONSET DEPRESSION DIFFER CLINICALLY AND NEUROBIOLOGICALLY FROM DEPRESSED PATIENTS WITH A CLINICAL HIGH-RISK STATE FOR PSYCHOSIS?. <i>Schizophrenia Bulletin</i> , 2018 , 44, S328-S329	1.3	78
136	Revealing the complex genetic architecture of obsessive-compulsive disorder using meta-analysis. <i>Molecular Psychiatry</i> , 2018 , 23, 1181-1188	15.1	205
135	T239. SINGLE-SUBJECT PREDICTION OF FUNCTIONAL OUTCOMES IN CLINICAL HIGH RISK SUBJECTS USING CLINICAL DATA. <i>Schizophrenia Bulletin</i> , 2018 , 44, S209-S210	1.3	78
134	Prediction Models of Functional Outcomes for Individuals in the Clinical High-Risk State for Psychosis or With Recent-Onset Depression: A Multimodal, Multisite Machine Learning Analysis. <i>JAMA Psychiatry</i> , 2018 , 75, 1156-1172	14.5	156
133	Integrating evolutionary and regulatory information with a multispecies approach implicates genes and pathways in obsessive-compulsive disorder. <i>Nature Communications</i> , 2017 , 8, 774	17.4	41
132	Intervention in Clinical High Risk States - Current Status and Future Perspectives. <i>European Psychiatry</i> , 2017 , 41, S27-S28	6	
131	Clinical High Risk Symptoms and Criteria in the Community: Prevalence, Clinical Significance and Risk Factors for Their Occurrence. <i>European Psychiatry</i> , 2017 , 41, S226-S226	6	1
130	Basic Symptoms in the Community and their Association with Age. <i>European Psychiatry</i> , 2017 , 41, S84-S85		1
129	Restricted attention to social cues in schizophrenia patients. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2016 , 266, 649-61	5.1	21
128	Psychosis-predictive value of self-reported schizotypy in a clinical high-risk sample. <i>Journal of Abnormal Psychology</i> , 2016 , 125, 923-932	7	46
127	Depression predicts persistence of paranoia in clinical high-risk patients to psychosis: results of the EPOS project. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016 , 51, 247-57	4.5	17
126	Predictors for symptom re-exacerbation after targeted stepwise drug discontinuation in first-episode schizophrenia: Results of the first-episode study within the German research network on schizophrenia. <i>Schizophrenia Research</i> , 2016 , 170, 168-76	3.6	27

125	Revisiting the Basic Symptom Concept: Toward Translating Risk Symptoms for Psychosis into Neurobiological Targets. <i>Frontiers in Psychiatry</i> , 2016 , 7, 9	5	40
124	Development of a stage-dependent prognostic model to predict psychosis in ultra-high-risk patients seeking treatment for co-morbid psychiatric disorders. <i>Psychological Medicine</i> , 2016 , 46, 1839-51	6.9	12
123	Authors'Reply. <i>British Journal of Psychiatry</i> , 2016 , 208, 197-8	5.4	
122	EPA guidance on the early detection of clinical high risk states of psychoses. <i>European Psychiatry</i> , 2015 , 30, 405-16	6	235
121	Personality dimensions in persons symptomatically at risk of psychosis: pronounced but lacking a characteristic profile. <i>Microbial Biotechnology</i> , 2015 , 9, 242-7	3.3	6
120	Duration of unspecific prodromal and clinical high risk states, and early help-seeking in first-admission psychosis patients. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015 , 50, 1831-41	4.5	41
119	Disorder, not just state of risk: meta-analysis of functioning and quality of life in people at high risk of psychosis. <i>British Journal of Psychiatry</i> , 2015 , 207, 198-206	5.4	180
118	Forecasting psychosis by event-related potentials-systematic review and specific meta-analysis. <i>Biological Psychiatry</i> , 2015 , 77, 951-8	7.9	76
117	EPA guidance on the early intervention in clinical high risk states of psychoses. <i>European Psychiatry</i> , 2015 , 30, 388-404	6	233
116	Neurocognitive functioning in parents of schizophrenia patients: Attentional and executive performance vary with genetic loading. <i>Psychiatry Research</i> , 2015 , 230, 885-91	9.9	10
115	Cross-disorder genome-wide analyses suggest a complex genetic relationship between Tourette's syndrome and OCD. <i>American Journal of Psychiatry</i> , 2015 , 172, 82-93	11.9	101
114	Psychosis prediction: stratification of risk estimation with information-processing and premorbid functioning variables. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1482-90	1.3	75
113	5-HT3 receptor influences the washing phenotype and visual organization in obsessive-compulsive disorder supporting 5-HT3 receptor antagonists as novel treatment option. <i>European Neuropsychopharmacology</i> , 2014 , 24, 86-94	1.2	18
112	Prediction and prevention of psychosis: current progress and future tasks. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2014 , 264 Suppl 1, S9-16	5.1	16
111	Copy number variation in obsessive-compulsive disorder and tourette syndrome: a cross-disorder study. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2014 , 53, 910-9	7.2	86
110	A stratified model for psychosis prediction in clinical practice. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1533-42	1.3	53
109	Prevalence and clinical significance of DSM-5-attenuated psychosis syndrome in adolescents and young adults in the general population: the Bern Epidemiological At-Risk (BEAR) study. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1499-508	1.3	70
108	Improving the clinical prediction of psychosis by combining ultra-high risk criteria and cognitive basic symptoms. <i>Schizophrenia Research</i> , 2014 , 154, 100-6	3.6	94

107	PREVENTIVE PSYCHOTHERAPY. <i>Schizophrenia Research</i> , 2014 , 153, S42-S43	3.6	1
106	Self-reported psychotic-like experiences are a poor estimate of clinician-rated attenuated and frank delusions and hallucinations. <i>Psychopathology</i> , 2014 , 47, 194-201	3.4	47
105	Identifying gene-environment interactions in schizophrenia: contemporary challenges for integrated, large-scale investigations. <i>Schizophrenia Bulletin</i> , 2014 , 40, 729-36	1.3	186
104	Short-term functional outcome and premorbid adjustment in clinical high-risk patients. Results of the EPOS project. <i>European Psychiatry</i> , 2014 , 29, 371-80	6	24
103	Can quantitative EEG measures predict clinical outcome in subjects at Clinical High Risk for psychosis? A prospective multicenter study. <i>Schizophrenia Research</i> , 2014 , 153, 42-7	3.6	38
102	Pathways to care in subjects at high risk for psychotic disorders - a European perspective. <i>Schizophrenia Research</i> , 2014 , 152, 400-7	3.6	21
101	Psychosocial outcome in patients at clinical high risk of psychosis: a prospective follow-up. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013 , 48, 303-11	4.5	27
100	Morphometry of structural disconnectivity indicators in subjects at risk and in age-matched patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2013 , 263, 15-24	5.1	20
99	Physical illnesses, developmental risk factors and psychiatric diagnoses among subjects at risk of psychosis. <i>European Psychiatry</i> , 2013 , 28, 135-40	6	3
98	Prediction of psychosis in clinical high-risk patients by the Schizotypal Personality Questionnaire. Results of the EPOS project. <i>European Psychiatry</i> , 2013 , 28, 469-75	6	36
97	The psychosis high-risk state: a comprehensive state-of-the-art review. <i>JAMA Psychiatry</i> , 2013 , 70, 107-20	4.5	965
96	The Strauss and Carpenter Prognostic Scale in subjects clinically at high risk of psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2013 , 127, 53-61	6.5	23
95	Genome-wide association study of obsessive-compulsive disorder. <i>Molecular Psychiatry</i> , 2013 , 18, 788-98	5.1	244
94	The functional coding variant Asn107Ile of the neuropeptide S receptor gene (NPSR1) influences age at onset of obsessive-compulsive disorder. <i>International Journal of Neuropsychopharmacology</i> , 2013 , 16, 1951-8	5.8	6
93	Partitioning the heritability of Tourette syndrome and obsessive compulsive disorder reveals differences in genetic architecture. <i>PLoS Genetics</i> , 2013 , 9, e1003864	6	189
92	A rose is a rose is a rose, but at-risk criteria differ. <i>Psychopathology</i> , 2013 , 46, 75-87	3.4	51
91	Characterization of SLITRK1 variation in obsessive-compulsive disorder. <i>PLoS ONE</i> , 2013 , 8, e70376	3.7	37
90	Basic disturbances of information processing in psychosis prediction. <i>Frontiers in Psychiatry</i> , 2013 , 4, 93	5	5

89	Personality disorders and accentuations in at-risk persons with and without conversion to first-episode psychosis. <i>Microbial Biotechnology</i> , 2012 , 6, 389-98	3.3	34
88	A promoter variant of SHANK1 affects auditory working memory in schizophrenia patients and in subjects clinically at risk for psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2012 , 262, 117-24	5.1	33
87	Antisaccade performance in patients with obsessive-compulsive disorder and unaffected relatives: further evidence for impaired response inhibition as a candidate endophenotype. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2012 , 262, 625-34	5.1	38
86	Perceived negative attitude of others predicts transition to psychosis in patients at risk of psychosis. <i>European Psychiatry</i> , 2012 , 27, 264-6	6	10
85	Pharmacological prevention and treatment in clinical at-risk states for psychosis. <i>Current Pharmaceutical Design</i> , 2012 , 18, 550-7	3.3	16
84	Whither the attenuated psychosis syndrome?. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1130-4	1.3	74
83	Preventing progression to first-episode psychosis in early initial prodromal states. <i>British Journal of Psychiatry</i> , 2012 , 200, 22-9	5.4	162
82	Poster #202 COMPARING THE PRODROME OF SCHIZOPHRENIA-SPECTRUM PSYCHOSES AND BIPOLAR DISORDERS WITH AND WITHOUT PSYCHOTIC FEATURES. <i>Schizophrenia Research</i> , 2012 , 136, S163-S164	3.6	
81	Axis I diagnoses and transition to psychosis in clinical high-risk patients EPOS project: prospective follow-up of 245 clinical high-risk outpatients in four countries. <i>Schizophrenia Research</i> , 2012 , 138, 192-7	3.6	85
80	Comparing the prodrome of schizophrenia-spectrum psychoses and affective disorders with and without psychotic features. <i>Schizophrenia Research</i> , 2012 , 138, 218-22	3.6	13
79	Chances and risks of predicting psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2012 , 262 Suppl 2, S85-90	5.1	16
78	Cannabis use and age at onset of symptoms in subjects at clinical high risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2012 , 125, 45-53	6.5	61
77	Basic symptoms and the prediction of first-episode psychosis. <i>Current Pharmaceutical Design</i> , 2012 , 18, 351-7	3.3	121
76	Neurophysiological correlates of impaired facial affect recognition in individuals at risk for schizophrenia. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1021-9	1.3	51
75	Rationale and first results of developing at-risk (prodromal) criteria for bipolar disorder. <i>Current Pharmaceutical Design</i> , 2012 , 18, 358-75	3.3	65
74	Subjective quality of life and its changes in patients at-risk for psychosis results of the EPOS study. <i>European Psychiatry</i> , 2011 , 26, 2039-2039	6	
73	Course of psychopathology in the at-risk mental state - outcomes beyond transition to psychosis. <i>European Psychiatry</i> , 2011 , 26, 2037-2037	6	
72	Prediction of psychosis by mismatch negativity. <i>Biological Psychiatry</i> , 2011 , 69, 959-66	7.9	219

71	The near Babylonian speech confusion in early detection of psychosis. <i>Schizophrenia Bulletin</i> , 2011 , 37, 653-5	1.3	54
70	Neuropsychological profiles in different at-risk states of psychosis: executive control impairment in the early--and additional memory dysfunction in the late--prodromal state. <i>Schizophrenia Bulletin</i> , 2011 , 37, 861-73	1.3	105
69	Prediction and prevention of schizophrenia: what has been achieved and where to go next?. <i>World Psychiatry</i> , 2011 , 10, 165-74	14.4	73
68	Prediction and Early Detection of First-Episode Psychosis 2011 , 207-267		2
67	Prevention and Early Intervention in At-Risk States for Developing Psychosis 2011 , 81-92		
66	Perceived parental rearing in subjects with obsessive-compulsive disorder and their siblings. <i>Acta Psychiatrica Scandinavica</i> , 2010 , 121, 280-8	6.5	30
65	The EPOS prediction model improves ability to predict transition to first episode psychosis in individuals at high risk. <i>Evidence-Based Mental Health</i> , 2010 , 13, 77	11.1	
64	Basic symptoms and ultrahigh risk criteria: symptom development in the initial prodromal state. <i>Schizophrenia Bulletin</i> , 2010 , 36, 182-91	1.3	152
63	Klinische Prädiktion und Prävention von Psychosen. <i>Psychiatrie Und Psychotherapie Up2date</i> , 2010 , 4, 237-252		
62	Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. <i>Archives of General Psychiatry</i> , 2010 , 67, 241-51		491
61	The interaction of working memory and emotion in persons clinically at risk for psychosis: an fMRI pilot study. <i>Schizophrenia Research</i> , 2010 , 120, 167-76	3.6	29
60	Probably at-risk, but certainly ill--advocating the introduction of a psychosis spectrum disorder in DSM-V. <i>Schizophrenia Research</i> , 2010 , 120, 23-37	3.6	114
59	Disability in people clinically at high risk of psychosis. <i>British Journal of Psychiatry</i> , 2010 , 197, 278-84	5.4	101
58	DAOA/G72 predicts the progression of prodromal syndromes to first episode psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2010 , 260, 209-15	5.1	29
57	Intervention in at-risk states for developing psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2010 , 260 Suppl 2, S90-4	5.1	13
56	Search for copy number variants in chromosomes 15q11-q13 and 22q11.2 in obsessive compulsive disorder. <i>BMC Medical Genetics</i> , 2010 , 11, 100	2.1	13
55	Anandamide elevation in cerebrospinal fluid in initial prodromal states of psychosis. <i>British Journal of Psychiatry</i> , 2009 , 194, 371-2	5.4	140
54	Gray matter abnormalities in subjects at ultra-high risk for schizophrenia and first-episode schizophrenic patients compared to healthy controls. <i>Psychiatry Research - Neuroimaging</i> , 2009 , 173, 163-9	2.9	116

53	Antisaccade performance is related to genetic loading for schizophrenia. <i>Journal of Psychiatric Research</i> , 2009 , 43, 291-7	5.2	12
52	Early onset of obsessive-compulsive disorder and associated comorbidity. <i>Depression and Anxiety</i> , 2009 , 26, 1012-7	8.4	36
51	Binocular depth inversion as a paradigm of reduced visual information processing in prodromal state, antipsychotic-naïve and treated schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2009 , 259, 195-202	5.1	74
50	Early detection of psychosis - establishing a service for persons at risk. <i>European Psychiatry</i> , 2009 , 24, 1-10	6	35
49	Perceived negative attitude of others as an early sign of psychosis. <i>European Psychiatry</i> , 2009 , 24, 233-8	6	9
48	Intervention in the at-risk state to prevent transition to psychosis. <i>Current Opinion in Psychiatry</i> , 2009 , 22, 177-83	4.9	22
47	Intervention in individuals at ultra-high risk for psychosis: a review and future directions. <i>Journal of Clinical Psychiatry</i> , 2009 , 70, 1206-12	4.6	219
46	Harm avoidance in subjects with obsessive-compulsive disorder and their families. <i>Journal of Affective Disorders</i> , 2008 , 107, 265-9	6.6	48
45	PHARMACOLOGICAL INTERVENTION IN THE PRODRAMAL STATE OF PSYCHOSIS. <i>Schizophrenia Research</i> , 2008 , 102, 47-48	3.6	
44	Sensory gating in schizophrenia: P50 and N100 gating in antipsychotic-free subjects at risk, first-episode, and chronic patients. <i>Biological Psychiatry</i> , 2008 , 64, 376-84	7.9	181
43	Auditory P300 in individuals clinically at risk for psychosis. <i>International Journal of Psychophysiology</i> , 2008 , 70, 192-205	2.9	79
42	P.3.f.001 Does anandamide elevation in cerebrospinal fluid protect against transition into frank psychosis?. <i>European Neuropsychopharmacology</i> , 2008 , 18, S474	1.2	
41	Increased neural response related to neutral faces in individuals at risk for psychosis. <i>NeuroImage</i> , 2008 , 40, 289-97	7.9	120
40	Obsessive-compulsive disorder and posttraumatic stress disorder. <i>Psychopathology</i> , 2008 , 41, 129-34	3.4	28
39	Interrelated neuropsychological and anatomical evidence of hippocampal pathology in the at-risk mental state. <i>Psychological Medicine</i> , 2008 , 38, 843-51	6.9	65
38	Kraepelin and psychotic prodromal conditions. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2008 , 258 Suppl 2, 74-84	5.1	37
37	Reduced subjective quality of life in persons at risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2008 , 117, 357-68	6.5	64
36	Neural correlates of working memory dysfunction in first-episode schizophrenia patients: an fMRI multi-center study. <i>Schizophrenia Research</i> , 2007 , 89, 198-210	3.6	130

35	Neurocognitive indicators for a conversion to psychosis: comparison of patients in a potentially initial prodromal state who did or did not convert to a psychosis. <i>Schizophrenia Research</i> , 2007 , 92, 116-23	3.6	161
34	Orienting of attention in unmedicated patients with schizophrenia, prodromal subjects and healthy relatives. <i>Schizophrenia Research</i> , 2007 , 97, 35-42	3.6	27
33	Randomized controlled multicentre trial of cognitive behaviour therapy in the early initial prodromal state: effects on social adjustment post treatment. <i>Microbial Biotechnology</i> , 2007 , 1, 71-8	3.3	48
32	Impulsiveness in obsessive-compulsive disorder: results from a family study. <i>Acta Psychiatrica Scandinavica</i> , 2007 , 115, 41-7	6.5	97
31	Relationship between subjective and objective cognitive function in the early and late prodrome. <i>British Journal of Psychiatry</i> , 2007 , 51, s43-51	5.4	36
30	Basic symptoms in early psychotic and depressive disorders. <i>British Journal of Psychiatry</i> , 2007 , 51, s31-7	5.4	59
29	Acute effects of treatment for prodromal symptoms for people putatively in a late initial prodromal state of psychosis. <i>British Journal of Psychiatry</i> , 2007 , 51, s88-95	5.4	100
28	Efficacy of flupentixol and risperidone in chronic schizophrenia with predominantly negative symptoms. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2007 , 31, 1012-22	5.5	20
27	The initial prodrome of schizophrenia: different duration, different underlying deficits?. <i>Comprehensive Psychiatry</i> , 2007 , 48, 479-88	7.3	31
26	CSF metabolic and proteomic profiles in patients prodromal for psychosis. <i>PLoS ONE</i> , 2007 , 2, e756	3.7	80
25	Neurocognitive functioning in subjects at risk for a first episode of psychosis compared with first- and multiple-episode schizophrenia. <i>Journal of Clinical and Experimental Neuropsychology</i> , 2006 , 28, 1388-407	2.1	108
24	Alexithymia in obsessive-compulsive disorder - results from a family study. <i>Psychotherapy and Psychosomatics</i> , 2006 , 75, 312-8	9.4	42
23	No human tryptophan hydroxylase-2 gene R441H mutation in a large cohort of psychiatric patients and control subjects. <i>Biological Psychiatry</i> , 2006 , 60, 202-3	7.9	46
22	Familiarity of obsessive-compulsive disorder in nonclinical and clinical subjects. <i>American Journal of Psychiatry</i> , 2006 , 163, 1986-92	11.9	74
21	The German Research Network on Schizophrenia--impact on the management of schizophrenia. <i>Dialogues in Clinical Neuroscience</i> , 2006 , 8, 115-21	5.7	13
20	Pharmacological intervention in the initial prodromal phase of psychosis. <i>European Psychiatry</i> , 2005 , 20, 1-6	6	45
19	Impaired mismatch negativity generation in prodromal subjects and patients with schizophrenia. <i>Schizophrenia Research</i> , 2005 , 73, 297-310	3.6	130
18	Subjective quality of life in subjects at risk for a first episode of psychosis: a comparison with first episode schizophrenia patients and healthy controls. <i>Schizophrenia Research</i> , 2005 , 79, 137-43	3.6	81

17	Support for the association between the rare functional variant I425V of the serotonin transporter gene and susceptibility to obsessive compulsive disorder. <i>Molecular Psychiatry</i> , 2005 , 10, 1059-61	15.1	43
16	Interventions in the initial prodromal states of psychosis in Germany: concept and recruitment. <i>British Journal of Psychiatry</i> , 2005 , 48, s45-8	5.4	50
15	The European Prediction of Psychosis Study (EPOS): integrating early recognition and intervention in Europe. <i>World Psychiatry</i> , 2005 , 4, 161-7	14.4	97
14	Intervención farmacológica en la fase prodromática inicial de la psicosis. <i>European Psychiatry (Ed Española)</i> , 2005 , 12, 217-222		
13	Early detection and secondary prevention of psychosis: facts and visions. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2004 , 254, 117-28	5.1	146
12	Early detection and intervention in the initial prodromal phase of schizophrenia. <i>Pharmacopsychiatry</i> , 2003 , 36 Suppl 3, S162-7	2	97
11	Dimensions of working memory dysfunction in schizophrenia. <i>Schizophrenia Research</i> , 2003 , 62, 259-68	3.6	60
10	Remembering or knowing: electrophysiological evidence for an episodic memory deficit in schizophrenia. <i>Psychological Medicine</i> , 2002 , 32, 1261-71	6.9	81
9	Effects of fluoxetine versus bright light in the treatment of seasonal affective disorder. <i>Psychological Medicine</i> , 1998 , 28, 923-33	6.9	65
8	Low plasma thyroid indices of depressed patients are attenuated by antidepressant drugs and influence treatment outcome. <i>Pharmacopsychiatry</i> , 1996 , 29, 180-6	2	39
7	A controlled study of the efficacy and safety of mianserin and amitriptyline in depressive inpatients. <i>Pharmacopsychiatry</i> , 1995 , 28, 249-52	2	14
6	Clinical response to sleep deprivation and auditory-evoked potentials--preliminary results. <i>Pharmacopsychiatry</i> , 1994 , 27, 70-1	2	7
5	Evidence for a seasonal form of recurrent brief depression (RBD-seasonal). <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1994 , 244, 205-10	5.1	6
4	Application of transcranial magnetic stimulation in treatment of drug-resistant major depression--report of two cases. <i>Human Psychopharmacology</i> , 1993 , 8, 361-365	2.3	138
3	Recurrent brief depression and its relationship to seasonal affective disorder. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1992 , 242, 20-6	5.1	25
2	Electroencephalographic Predictors of Psychosis. <i>Key Issues in Mental Health</i> , 133-141		
1	Dysregulated lipid metabolism precedes onset of psychosis in people at clinical high risk		1