## Ting-Yun Lin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5285237/publications.pdf

Version: 2024-02-01

840776 713466 21 641 11 21 citations h-index g-index papers 21 21 21 1101 all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	Volume overload correlates with cardiovascular risk factors in patients with chronic kidney disease. Kidney International, 2014, 85, 703-709.	5.2	194
2	Indoxyl Sulfate: A Novel Cardiovascular Risk Factor in Chronic Kidney Disease. Journal of the American Heart Association, 2017, $6$ , .	3.7	127
3	Body composition is associated with clinical outcomes in patients with non–dialysis-dependent chronic kidney disease. Kidney International, 2018, 93, 733-740.	5.2	56
4	Impact of Misclassification of Obesity by Body Mass Index on Mortality in Patients With CKD. Kidney International Reports, 2018, 3, 447-455.	0.8	40
5	Factors Associated with Decreased Lean Tissue Index in Patients with Chronic Kidney Disease. Nutrients, 2017, 9, 434.	4.1	27
6	Gut dysbiosis and mortality in hemodialysis patients. Npj Biofilms and Microbiomes, 2021, 7, 20.	6.4	26
7	Obesity and risk of end-stage renal disease in patients with chronic kidney disease: a cohort study. American Journal of Clinical Nutrition, 2018, 108, 1145-1153.	4.7	24
8	Development and validation of a multifrequency bioimpedance spectroscopy equation to predict appendicular skeletal muscle mass in hemodialysis patients. Clinical Nutrition, 2021, 40, 3288-3295.	5.0	18
9	Comparative Gut Microbiome Differences between Ferric Citrate and Calcium Carbonate Phosphate Binders in Patients with End-Stage Kidney Disease. Microorganisms, 2020, 8, 2040.	3.6	17
10	Nephroprotective Role of Chrysophanol in Hypoxia/Reoxygenation-Induced Renal Cell Damage via Apoptosis, ER Stress, and Ferroptosis. Biomedicines, 2021, 9, 1283.	3.2	17
11	Geriatric Nutritional Risk Index Is Associated with Unique Health Conditions and Clinical Outcomes in Chronic Kidney Disease Patients. Nutrients, 2019, 11, 2769.	4.1	16
12	Association of subjective global assessment of nutritional status with gut microbiota in hemodialysis patients: a case–control study. Nephrology Dialysis Transplantation, 2021, 36, 1104-1111.	0.7	15
13	Normal-weight obesity and clinical outcomes in nondiabetic chronic kidney disease patients: a cohort study. American Journal of Clinical Nutrition, 2018, 107, 664-672.	4.7	13
14	Anti-Acid Drug Treatment Induces Changes in the Gut Microbiome Composition of Hemodialysis Patients. Microorganisms, 2021, 9, 286.	3.6	10
15	A Comparison of SARC-F, Calf Circumference, and Their Combination for Sarcopenia Screening among Patients Undergoing Peritoneal Dialysis. Nutrients, 2022, 14, 923.	4.1	10
16	Indoxyl Sulfate and Incident Peripheral Artery Disease in Hemodialysis Patients. Toxins, 2020, 12, 696.	3.4	8
17	Characterization of Gut Microbiota Composition in Hemodialysis Patients With Normal Weight Obesity. Journal of Clinical Endocrinology and Metabolism, 2020, 105, 2006-2014.	3.6	8
18	Loss of Function of von Hippel-Lindau Trigger Lipocalin 2-Dependent Inflammatory Responses in Cultured and Primary Renal Tubular Cells. Oxidative Medicine and Cellular Longevity, 2021, 2021, 1-10.	4.0	6

## Ting-Yun Lin

#	Article	IF	CITATION
19	Malnutrition is Associated with Increased Morbidity and Death in Dialysis Patients Undergoing Endovascular Therapy for Peripheral Artery Disease. European Journal of Vascular and Endovascular Surgery, 2022, 64, 225-233.	1.5	4
20	Differences in the Microbial Composition of Hemodialysis Patients Treated with and without $\hat{l}^2$ -Blockers. Journal of Personalized Medicine, 2021, 11, 198.	2.5	3
21	Central obesity and incident atherosclerotic cardiovascular disease events in hemodialysis patients. Nutrition, Metabolism and Cardiovascular Diseases, 2020, 30, 500-507.	2.6	2