

Atsunori Kashiwagi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5291009/publications.pdf>

Version: 2024-02-01

37
papers

1,919
citations

361413
20
h-index

315739
38
g-index

40
all docs

40
docs citations

40
times ranked

2161
citing authors

#	ARTICLE	IF	CITATIONS
1	Reduction in cardiovascular disease events in patients with type 2 diabetes mellitus treated with a sodium-glucose cotransporter 2 inhibitor versus a dipeptidyl peptidase 4 inhibitor: A real-world retrospective administrative database analysis in Japan. <i>Journal of Diabetes Investigation</i> , 2022, 13, 1175-1189.	2.4	5
2	Sodium-glucose cotransporter 2 inhibitors represent a paradigm shift in the prevention of heart failure in type 2 diabetes patients. <i>Journal of Diabetes Investigation</i> , 2021, 12, 6-20.	2.4	17
3	A Prospective, Open-Label Short-Term Pilot Study on Modification of the Skin Hydration Status During Treatment With a Sodium-Glucose Cotransporter-2 Inhibitor. <i>Diabetes Therapy</i> , 2021, 12, 431-440.	2.5	2
4	<i>Porphyromonas gingivalis</i> induces entero-hepatic metabolic derangements with alteration of gut microbiota in a type 2 diabetes mouse model. <i>Scientific Reports</i> , 2021, 11, 18398.	3.3	19
5	Safety of Ipragliflozin in Patients with Type 2 Diabetes Mellitus: Pooled Analysis of Phase II/III/IV Clinical Trials. <i>Diabetes Therapy</i> , 2019, 10, 2201-2217.	2.5	11
6	A new door opens, but it is essential to accumulate further clinical evidence to control heart failure in diabetes with preserved ejection fraction. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1145-1147.	2.4	2
7	Preserving β -cell function is the major determinant of diabetes remission following laparoscopic sleeve gastrectomy in Japanese obese diabetic patients. <i>Endocrine Journal</i> , 2019, 66, 817-826.	1.6	6
8	Achieving LDL cholesterol target levels <1.81 mmol/L may provide extra cardiovascular protection in patients at high risk: Exploratory analysis of the Standard Versus Intensive Statin Therapy for Patients with Hypercholesterolemia and Diabetic Retinopathy study. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 791-800.	4.4	15
9	Intensive Treat-to-Target Statin Therapy in High-Risk Japanese Patients With Hypercholesterolemia and Diabetic Retinopathy: Report of a Randomized Study. <i>Diabetes Care</i> , 2018, 41, 1275-1284.	8.6	43
10	Impact of obesity on annual medical expenditures and diabetes care in Japanese patients with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2018, 9, 776-781.	2.4	10
11	Definitive diagnosis of mandibular hypoplasia, deafness, progeroid features and lipodystrophy (MDPL) syndrome caused by a recurrent <i>de novo</i> mutation in the <i>POLD1</i> gene. <i>Endocrine Journal</i> , 2018, 65, 227-238.	1.6	42
12	Improved cardiometabolic risk factors in Japanese patients with type 2 diabetes treated with ipragliflozin: a pooled analysis of six randomized, placebo-controlled trials. <i>Endocrine Journal</i> , 2018, 65, 693-705.	1.6	15
13	Effects of ipragliflozin, a selective sodium-glucose co-transporter 2 inhibitor, on blood pressure in Japanese patients with type 2 diabetes mellitus: a pooled analysis of six randomized, placebo-controlled clinical trials. <i>Diabetology International</i> , 2017, 8, 76-86.	1.4	12
14	Metabolic and hemodynamic effects of sodium-dependent glucose cotransporter 2 inhibitors on cardiovascular protection in the treatment of patients with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2017, 8, 416-427.	2.4	59
15	Evaluation of a Novel Glucose Area Under the Curve (AUC) Monitoring System: Comparison with the AUC by Continuous Glucose Monitoring. <i>Diabetes and Metabolism Journal</i> , 2016, 40, 326.	4.7	7
16	Comparative Effects of Direct Renin Inhibitor and Angiotensin Receptor Blocker on Albuminuria in Hypertensive Patients with Type 2 Diabetes. A Randomized Controlled Trial. <i>PLoS ONE</i> , 2016, 11, e0164936.	2.5	11
17	Relationship between the efficacy of oral antidiabetic drugs and clinical features in type 2 diabetic patients (JDDM38). <i>Journal of Diabetes Investigation</i> , 2016, 7, 386-395.	2.4	15
18	Efficacy and safety of ipragliflozin in Japanese patients with type 2 diabetes stratified by body mass index: A subgroup analysis of five randomized clinical trials. <i>Journal of Diabetes Investigation</i> , 2016, 7, 544-554.	2.4	29

#	ARTICLE	IF	CITATIONS
19	Efficacy and safety of 40Âmg or 60Âmg duloxetine in Japanese adults with diabetic neuropathic pain: Results from a randomized, 52-week, open-label study. <i>Journal of Diabetes Investigation</i> , 2016, 7, 100-108.	2.4	23
20	Comparison of clinical characteristics in patients with type 2 diabetes among whom different antihyperglycemic agents were prescribed as monotherapy or combination therapy by diabetes specialists. <i>Journal of Diabetes Investigation</i> , 2016, 7, 260-269.	2.4	13
21	Glucose area under the curve during oral glucose tolerance test as an index of glucose intolerance. <i>Diabetology International</i> , 2016, 7, 53-58.	1.4	81
22	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , 2016, 246, 141-147.	0.8	48
23	Duality of n-3 Polyunsaturated Fatty Acids on Mcp-1 Expression in Vascular Smooth Muscle: A Potential Role of 4-Hydroxy Hexenal. <i>Nutrients</i> , 2015, 7, 8112-8126.	4.1	7
24	Efficacy and safety of ipragliflozin as an add-on to a sulfonylurea in Japanese patients with inadequately controlled type 2 diabetes: results of the randomized, placebo-controlled, double-blind, phase III EMIT study. <i>Diabetology International</i> , 2015, 6, 125-138.	1.4	47
25	Efficacy and safety of ipragliflozin as an add-on to pioglitazone in Japanese patients with inadequately controlled type 2 diabetes: a randomized, double-blind, placebo-controlled study (the SPOTLIGHT) <i>Tj ETQq1 1 0.784314 rgBT4/Overlook</i>	1.4	47
26	Ipragliflozin improves glycemic control in Japanese patients with type 2 diabetes mellitus: the BRIGHTEN study. <i>Diabetology International</i> , 2015, 6, 8-18.	1.4	66
27	Randomized, placebo-controlled, double-blind glycemic control trial of novel sodium-dependent glucose cotransporter 2 inhibitor ipragliflozin in Japanese patients with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2014, 5, 382-391.	2.4	90
28	A fish-based diet intervention improves endothelial function in postmenopausal women with type 2 diabetes mellitus: A randomized crossover trial. <i>Metabolism: Clinical and Experimental</i> , 2014, 63, 930-940.	3.4	43
29	Pharmacokinetic and pharmacodynamic study of ipragliflozin in Japanese patients with type 2 diabetes mellitus: A randomized, double-blind, placebo-controlled study. <i>Diabetes Research and Clinical Practice</i> , 2014, 106, 50-56.	2.8	32
30	Liraglutide versus Sitagliptin in a 24-week, Multicenter, Open-label, Randomized, Parallel-group Study in Japanese Type 2 Diabetes Mellitus Patients Responding Inadequately to a Sulfonylurea and/or One or Two Other Oral Antidiabetic Drugs (JDDM 33). <i>Japanese Clinical Medicine</i> , 2014, 5, JCM.S16585.	1.9	6
31	Omega-3 polyunsaturated fatty acid has an anti-oxidant effect via the Nrf-2/HO-1 pathway in 3T3-L1 adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2013, 430, 225-230.	2.1	81
32	4-Hydroxy Hexenal Derived from Docosahexaenoic Acid Protects Endothelial Cells via Nrf2 Activation. <i>PLoS ONE</i> , 2013, 8, e69415.	2.5	69
33	International clinical harmonization of glycated hemoglobin in Japan: From Japan Diabetes Society to National Glycohemoglobin Standardization Program values. <i>Journal of Diabetes Investigation</i> , 2012, 3, 39-40.	2.4	731
34	Low concentration of 4-hydroxy hexenal increases heme oxygenase-1 expression through activation of Nrf2 and antioxidative activity in vascular endothelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2010, 402, 99-104.	2.1	65
35	Coronary endothelial dysfunction in the insulin-resistant state is linked to abnormal pteridine metabolism and vascular oxidative stress. <i>Journal of the American College of Cardiology</i> , 2001, 38, 1821-1828.	2.8	68
36	Free radical production in endothelial cells as a pathogenetic factor for vascular dysfunction in the insulin resistance state. <i>Diabetes Research and Clinical Practice</i> , 1999, 45, 199-203.	2.8	49

#	ARTICLE	IF	CITATIONS
37	Insulin Signaling and Its Regulation of System A Amino Acid Uptake in Cultured Rat Vascular Smooth Muscle Cells. <i>Circulation Research</i> , 1996, 79, 1167-1176.	4.5	33