

Joaquin Sanchiz

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5293615/joaquin-sanchiz-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

93
papers

4,209
citations

41
h-index

63
g-index

98
ext. papers

4,380
ext. citations

4.3
avg, IF

5.11
L-index

| # | Paper | IF | Citations |
|----|--|------|-----------|
| 93 | Synthesis, structure and magnetic properties of a cobalt(II) mesoxalate 1D coordination polymer. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2021 , 647, 485-489 | 1.3 | 0 |
| 92 | Bio-inspired Ni dinuclear complexes as heterogeneous catalysts for hydrogen evolution. <i>Chemical Engineering Journal</i> , 2021 , 420, 130342 | 14.7 | 4 |
| 91 | Crystal structure and magnetic interactions of a new alkoxido and azido bridged 1D copper(II) coordination polymer. <i>Journal of Solid State Chemistry</i> , 2021 , 303, 122484 | 3.3 | 2 |
| 90 | Evolution of the external quantum efficiency of Si-based PV minimodules with encapsulated down-shifters and aged under UV radiation. <i>Materials Science and Engineering B: Solid-State Materials for Advanced Technology</i> , 2020 , 261, 114763 | 3.1 | 1 |
| 89 | Synthesis, crystal structure and magnetic properties of a pentanuclear Mn(III) cluster with 1,2,4-triazole based Schiff base ligand. <i>Inorganica Chimica Acta</i> , 2020 , 505, 119461 | 2.7 | 15 |
| 88 | Visible and NIR emitting Yb(iii) and Er(iii) complexes sensitized by β -diketonates and phenanthroline derivatives.. <i>RSC Advances</i> , 2020 , 10, 27815-27823 | 3.7 | 5 |
| 87 | The effect of the orientation of the Jahn-Teller distortion on the magnetic interactions of trinuclear mixed-valence Mn(ii)/Mn(iii) complexes. <i>Dalton Transactions</i> , 2019 , 48, 13799-13812 | 4.3 | 16 |
| 86 | Crystal structure and magneto-structural investigation of alkoxido bridged dinuclear Fe(III) complexes with 1,3-oxazolidine ligands. <i>Polyhedron</i> , 2019 , 162, 20-29 | 2.7 | 13 |
| 85 | Downshifting maximization procedure applied to [Eu(bphen)(tta)3] at different concentrations applied to a photovoltaic device and covered with a hemispherical reflector. <i>Sensors and Actuators A: Physical</i> , 2018 , 271, 60-65 | 3.9 | 7 |
| 84 | Highly luminescent film as enhancer of photovoltaic devices. <i>Journal of Luminescence</i> , 2018 , 201, 148-155 | 5.8 | 10 |
| 83 | Structure and magnetic behavior of unpredictable EE-azide bridged tetranuclear Mn(II) complex with ONO-donor hydrazone ligand and its transformation to dinuclear Mn(III) complex. <i>Polyhedron</i> , 2018 , 147, 142-151 | 2.7 | 28 |
| 82 | Durability analysis of the [Eu(bphen)(tta)3] down-shifter on Si-based PV modules exposed to extreme outdoor conditions. <i>Sensors and Actuators A: Physical</i> , 2018 , 276, 312-319 | 3.9 | 3 |
| 81 | Magnetostructural relationships in polymorphic ethylmalonate-containing copper(II) coordination polymers. <i>CrystEngComm</i> , 2018 , 20, 7464-7472 | 3.3 | 2 |
| 80 | Alternative and fully experimental procedure for characterizing down-shifters placed on photovoltaic devices. <i>Solar Energy Materials and Solar Cells</i> , 2018 , 185, 312-317 | 6.4 | 4 |
| 79 | Effect of the apical ligand on the geometry and magnetic properties of copper(ii)/mesoxalate trinuclear units. <i>Dalton Transactions</i> , 2017 , 46, 5260-5268 | 4.3 | 4 |
| 78 | H-bonding directed formation of 1D-single chains, 2D-sheets, and 3D structures in magnetically coupled tetranuclear nickel(II) complexes with incomplete double cubane core. <i>Polyhedron</i> , 2017 , 123, 361-375 | 2.7 | 11 |
| 77 | Optical properties of Nd ³⁺ -doped Tutton salts crystals. <i>Journal of Luminescence</i> , 2017 , 192, 136-140 | 3.8 | 2 |

| | | | |
|----|--|-----|----|
| 76 | Synthesis, characterization and magnetic properties of a dinuclear oxidovanadium(IV) complex: Magneto-structural DFT studies on the effects of out-of-plane OCH_3 angle. <i>Polyhedron</i> , 2017 , 122, 194-202 | 2.7 | 24 |
| 75 | Mesoxalate as Cu(II)/Mn(III) linker in the construction of MOFs in DMSO/water medium. <i>CrystEngComm</i> , 2015 , 17, 6555-6565 | 3.3 | 5 |
| 74 | Antiferromagnetically Coupled Dimeric Dodecacopper Supramolecular Architectures of Macrocyclic Ligands with a Symmetrical β -BO ₃ (3-) Central Moiety. <i>Inorganic Chemistry</i> , 2015 , 54, 6873-84 | 5.1 | 9 |
| 73 | Synthesis, structure, magnetic properties and EPR spectroscopy of a copper(II) coordination polymer with a ditopic hydrazone ligand and acetate bridges. <i>Dalton Transactions</i> , 2015 , 44, 1782-9 | 4.3 | 42 |
| 72 | Proton conduction and long-range ferrimagnetic ordering in two isostructural Copper(II) mesoxalate metal-organic frameworks. <i>Inorganic Chemistry</i> , 2015 , 54, 1597-605 | 5.1 | 42 |
| 71 | A new cost-effective polymeric film containing an Eu(III) complex acting as UV protector and down-converter for Si-based solar cells and modules. <i>Solar Energy Materials and Solar Cells</i> , 2015 , 136, 187-192 | 6.4 | 27 |
| 70 | Luminescent polymeric film containing an Eu(III) complex acting as UV protector and down-converter for Si-based solar cells and modules. <i>Surface and Coatings Technology</i> , 2015 , 271, 106-114 | 4.4 | 22 |
| 69 | Synthesis, crystal structure and magnetic properties of a trinuclear phenolate bridged manganese complex containing Mn(II)/Mn(III) ions. <i>RSC Advances</i> , 2014 , 4, 36175 | 3.7 | 7 |
| 68 | 1,3,5,7-Tetrakis(tetrazol-5-yl)-adamantane: the smallest tetrahedral tetrazole-functionalized ligand and its complexes formed by reaction with anhydrous M(II)Cl ₂ (M = Mn, Cu, Zn, Cd). <i>Dalton Transactions</i> , 2014 , 43, 12590-605 | 4.3 | 20 |
| 67 | Heteronuclear, mixed-metal Ag(I)-Mn(II) coordination polymers with bridging N-pyridinylisonicotinohydrazide ligands: synthesis, crystal structures, magnetic and photoluminescence properties. <i>Dalton Transactions</i> , 2014 , 43, 11925-35 | 4.3 | 36 |
| 66 | Influence of the coligand in the magnetic properties of a series of copper(II)/phenylmalonate complexes. <i>CrystEngComm</i> , 2014 , 16, 8106-8118 | 3.3 | 13 |
| 65 | Syntheses, structures and magnetic properties of azido- and phenoxo-bridged complexes of manganese containing tridentate arylhydrazone based ligands. <i>Polyhedron</i> , 2013 , 61, 45-55 | 2.7 | 46 |
| 64 | Bifunctional pyrazolate/carboxylate ligands for isorecticular cobalt and zinc MOF-5 analogs with magnetic analysis of the {Co ₄ (μ -O)} node. <i>CrystEngComm</i> , 2013 , 15, 9757 | 3.3 | 82 |
| 63 | A rare alb-4,8-Cmce metal coordination network based on tetrazolate and phosphonate functionalized 1,3,5,7-tetraphenyladamantane. <i>CrystEngComm</i> , 2013 , 15, 1235 | 3.3 | 37 |
| 62 | Cation effect on the crystal structure of polynuclear complexes with 2,2'-oxydiacetate as bridging ligand. <i>Inorganica Chimica Acta</i> , 2013 , 394, 196-202 | 2.7 | 5 |
| 61 | Copper(II) complexes with a new carboxylic-functionalized arylhydrazone of β -diketone as effective catalysts for acid-free oxidations. <i>New Journal of Chemistry</i> , 2012 , 36, 1646 | 3.6 | 48 |
| 60 | Hydrophobic-exterior layer structures and magnetic properties of trinuclear copper complexes with chiral amino alcoholate ligands. <i>New Journal of Chemistry</i> , 2012 , 36, 1596 | 3.6 | 37 |
| 59 | Bis(benzotriazol-1-yl)methane as a linker in the assembly of new copper(II) coordination polymers: Synthesis, structure and investigations. <i>Polyhedron</i> , 2012 , 48, 253-263 | 2.7 | 8 |

| | | | |
|----|---|-----|-----|
| 58 | Homochiral lanthanoid(III) mesoxalate metal-organic frameworks: synthesis, crystal growth, chirality, magnetic and luminescent properties. <i>CrystEngComm</i> , 2012 , 14, 2635 | 3.3 | 71 |
| 57 | Structures and magnetic properties of an antiferromagnetically coupled polymeric copper(II) complex and ferromagnetically coupled hexanuclear nickel(II) clusters. <i>Inorganic Chemistry</i> , 2012 , 51, 3270-82 | 5.1 | 41 |
| 56 | Two-dimensional (6,3) networks obtained with the {Cu ₃ (Hmesox) ₃ } ₃ secondary building unit (H ₄ mesox = mesoxalic acid). <i>CrystEngComm</i> , 2012 , 14, 4289 | 3.3 | 10 |
| 55 | Copper(II)-phenylmalonate complexes with the bifunctional ligands nicotinamide and isonicotinamide. <i>Polyhedron</i> , 2011 , 30, 2451-2458 | 2.7 | 10 |
| 54 | A ferromagnetically coupled copper(II) trinuclear secondary building unit as precursor for the preparation of molecule-based magnets. <i>Inorganica Chimica Acta</i> , 2011 , 371, 47-52 | 2.7 | 9 |
| 53 | [Cu ₃ (Hmesox) ₃] ₃ : a precursor for the rational design of chiral molecule-based magnets (H ₄ mesox = 2-dihydroxymalonic acid). <i>Inorganic Chemistry</i> , 2010 , 49, 7880-9 | 5.1 | 17 |
| 52 | Magnetic ordering in two molecule-based (10,3)-a nets prepared from a copper(II) trinuclear secondary building unit. <i>Inorganic Chemistry</i> , 2010 , 49, 7478-90 | 5.1 | 60 |
| 51 | Building-block process for the synthesis of new chromium(III) malonate complexes. <i>CrystEngComm</i> , 2010 , 12, 2711 | 3.3 | 19 |
| 50 | Spontaneous resolution upon crystallization of chiral La(III) and Gd(III) MOFs from achiral dihydroxymalonnate. <i>Chemical Communications</i> , 2010 , 46, 8270-2 | 5.8 | 108 |
| 49 | Self-assembled 3D heterometallic Cu(II)/Fe(II) coordination polymers with octahedral net skeletons: structural features, molecular magnetism, thermal and oxidation catalytic properties. <i>Inorganic Chemistry</i> , 2010 , 49, 11096-105 | 5.1 | 70 |
| 48 | Iron, copper and zinc ammonium-1-hydroxyalkylidene-diphosphonates with zero-, one- and two-dimensional covalent metal-ligand structures extended into three-dimensional supramolecular networks by charge-assisted hydrogen-bonding. <i>Polyhedron</i> , 2010 , 29, 2537-2545 | 2.7 | 44 |
| 47 | Synthesis, molecular structure and magnetic properties of a rhenium(IV) compound with catechol. <i>Journal of Molecular Structure</i> , 2009 , 921, 80-84 | 3.4 | 9 |
| 46 | Magnetic and luminescence properties of Cu(II), Cu(II)4O ₄ core, and Cd(II) mixed-ligand metal-organic frameworks constructed from 1,2-bis(1,2,4-triazol-4-yl)ethane and benzene-1,3,5-tricarboxylate. <i>Inorganica Chimica Acta</i> , 2009 , 362, 2452-2460 | 2.7 | 149 |
| 45 | Structure and magnetic properties of a tetranuclear Cu ₄ O ₄ open-cubane in [Cu(L)] ₄ ·4H ₂ O with L ₂ = (E)-N ⁺ -(2-oxy-3-methoxybenzylidene)benzohydrazide. <i>Inorganica Chimica Acta</i> , 2009 , 362, 3791-3795 | 2.7 | 54 |
| 44 | Copper(II)-methylmalonate complexes with unidentate N-donor ligands: Syntheses, structural characterization and magnetic properties. <i>Polyhedron</i> , 2009 , 28, 1802-1807 | 2.7 | 13 |
| 43 | Mono-, di- and polynuclear copper(II) compounds derived from N-butyl-diethanolamine: structural features, magnetism and catalytic activity for the mild peroxidative oxidation of cyclohexane. <i>Dalton Transactions</i> , 2009 , 2109-20 | 4.3 | 100 |
| 42 | Mixed-ligand coordination polymers from 1,2-bis(1,2,4-triazol-4-yl)ethane and benzene-1,3,5-tricarboxylate: trinuclear nickel or zinc secondary building units for three-dimensional networks with crystal-to-crystal transformation upon dehydration. <i>Dalton Transactions</i> , 2008 , 1734-44 | 4.3 | 246 |
| 41 | (infinity) ₂ [Cu ₂ (mu ₅ -btb)(mu-OH)(mu-H ₂ O)]: a two-dimensional coordination polymer built from ferromagnetically coupled Cu ₂ units (btb = benzene-1,2,3-tricarboxylate). <i>Dalton Transactions</i> , 2008 , 4877-84 | 4.3 | 76 |

| | | | |
|----|--|-----|-----|
| 40 | Multicopper(II) Pyromellitate Compounds: Self-Assembly Synthesis, Structural Topologies, and Magnetic Features. <i>Crystal Growth and Design</i> , 2008 , 8, 4100-4108 | 3-5 | 64 |
| 39 | Self-assembled copper(II) coordination polymers derived from aminopolyalcohols and benzenepolycarboxylates: structural and magnetic properties. <i>Inorganic Chemistry</i> , 2008 , 47, 162-75 | 5-1 | 109 |
| 38 | Crystal engineering of 3-D coordination polymers by pillaring ferromagnetic copper(II)-methylmalonate layers. <i>CrystEngComm</i> , 2007 , 9, 478-487 | 3-3 | 88 |
| 37 | Triangular oxalate clusters [W(3)(mu(3)-S)(mu(2)-S(2))(3)(C(2)O(4))(3)](2)(-) as building blocks for coordination polymers and nanosized complexes. <i>Inorganic Chemistry</i> , 2007 , 46, 2115-23 | 5-1 | 21 |
| 36 | Versatile supramolecular self-assembly. <i>CrystEngComm</i> , 2006 , 8, 530-544 | 3-3 | 35 |
| 35 | Metamagnetism in hydrophobically induced carboxylate (phenylmalonate)-bridged copper(II) layers. <i>Chemical Communications</i> , 2006 , 2857-9 | 5-8 | 31 |
| 34 | Versatile supramolecular self-assembly. Part I. Network formation and magnetic behaviour of the alkaline salts of the bis(malonate)cuprate(II) anion. <i>CrystEngComm</i> , 2006 , 8, 507-529 | 3-3 | 36 |
| 33 | Magnetic properties of copper(II) complexes containing peptides. Crystal structure of [Cu(phe-leu)]. <i>Journal of Molecular Structure</i> , 2006 , 797, 179-183 | 3-4 | 15 |
| 32 | Polymeric networks of copper(II) phenylmalonate with heteroaromatic n-donor ligands: synthesis, crystal structure, and magnetic properties. <i>Inorganic Chemistry</i> , 2005 , 44, 7794-801 | 5-1 | 49 |
| 31 | Synthesis, crystal structure and magnetic properties of two-dimensional malonato-bridged cobalt(II) and nickel(II) compounds. <i>CrystEngComm</i> , 2004 , 6, 106-111 | 3-3 | 97 |
| 30 | Phenylmalonate-Containing Copper(II) Complexes: Synthesis, Crystal Structure and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2004 , 2004, 4081-4090 | 2-3 | 56 |
| 29 | Protonated malonate: the influence of the hydrogen bonds on the magnetic behaviour. <i>CrystEngComm</i> , 2004 , 6, 443-450 | 3-3 | 30 |
| 28 | Design of high-dimensional copper(II) malonate complexes with exo-polydentate N-donor ligands. <i>Inorganic Chemistry</i> , 2003 , 42, 5938-48 | 5-1 | 111 |
| 27 | Interactions of nitric oxide with copper(II) dithiocarbamates in aqueous solution. <i>Journal of Inorganic Biochemistry</i> , 2003 , 95, 283-90 | 4-2 | 27 |
| 26 | Malonate-based copper(II) coordination compounds: ferromagnetic coupling controlled by dicarboxylates. <i>Polyhedron</i> , 2003 , 22, 2143-2153 | 2-7 | 101 |
| 25 | Malonic acid: a multi-modal bridging ligand for new architectures and properties on molecule-based magnets. <i>Polyhedron</i> , 2003 , 22, 2111-2123 | 2-7 | 74 |
| 24 | High-dimensional malonate-based materials: Synthesis, crystal structures and magnetic properties of [M2(mal)2(L)(H2O)2]n·h(H2O) M = Zn(II), Co(II); H2mal = malonic acid, L = pyrimidine, pyrazine. <i>CrystEngComm</i> , 2003 , 5, 280-284 | 3-3 | 51 |
| 23 | Crystal structures and magnetic properties of two- and three-dimensional malonato-bridged manganese(II) complexes. <i>Dalton Transactions</i> , 2003 , 2359-2365 | 4-3 | 65 |

| | | | |
|----|--|------|-----|
| 22 | {[Cu(H ₂ O) ₃][Cu(phmal) ₂]} _n : a new two-dimensional copper(II) complex with intralayer ferromagnetic interactions (phmal = phenylmalonate dianion). <i>New Journal of Chemistry</i> , 2003 , 27, 1557-1562 | 3.6 | 49 |
| 21 | Holo- and hemidirected lead(II) in the polymeric [Pb(4)(μ-3,4-TDTA) ₂ (H ₂ O) ₂]*4H ₂ O complex. N,N,N',N''-tetraacetate ligands derived from o-phenylenediamines as sequestering agents for lead(II). <i>Inorganic Chemistry</i> , 2002 , 41, 6048-55 | 5.1 | 40 |
| 20 | Structural versatility of the malonate ligand as a tool for crystal engineering in the design of molecular magnets. <i>CrystEngComm</i> , 2002 , 4, 522-535 | 3.3 | 130 |
| 19 | Ferromagnetic coupling in the malonato-bridged copper(II) chains [Cu(Im) ₂ (mal)] _n and [Cu(2-Melm) ₂ (mal)] _n (H ₂ mal = malonic acid, Im = imidazole and 2-Melm = 2-methylimidazole). <i>New Journal of Chemistry</i> , 2002 , 26, 1624-1628 | 3.6 | 77 |
| 18 | The flexibility of molecular components as a suitable tool in designing extended magnetic systems. <i>CrystEngComm</i> , 2002 , 4, 440-446 | 3.3 | 59 |
| 17 | Alternating cationic/anionic layers in the [MII(H ₂ O) ₆][CuII(mal) ₂ (H ₂ O)] complexes linked through hydrogen bonds (M = Mn, Co, Ni, Cu and Zn; H ₂ mal = malonic acid). <i>CrystEngComm</i> , 2002 , 4, 631-637 | 3.3 | 60 |
| 16 | A new eight-coordinate complex of manganese(II): synthesis, crystal structure, spectroscopy and magnetic properties of [Mn(Hoxam) ₂ (H ₂ O) ₄] (H ₂ oxam=oxamic acid). <i>Inorganica Chimica Acta</i> , 2001 , 315, 120-125 | 2.7 | 13 |
| 15 | Crystal structure and magnetic properties of the flexible self-assembled two-dimensional square network complex [Cu ₂ (mal) ₂ (H ₂ O) ₂ (4,4'-bpy)] (H ₂ mal=malonic acid and 4,4'-bpy=4,4'-bipyridine). <i>Inorganica Chimica Acta</i> , 2001 , 318, 159-165 | 2.7 | 125 |
| 14 | Synthesis, crystal structure and magnetic properties of [Cu(bpym)(mal)(H ₂ O)] ₂ ·H ₂ O and [Cu ₂ (bpym)(mal) ₂ (H ₂ O) ₂] ₂ ·4H ₂ O (bpym=2,2'-bipyrimidine, H ₂ mal=malonic acid). <i>Inorganica Chimica Acta</i> , 2001 , 326, 20-26 | 2.7 | 57 |
| 13 | Crystal structure of the 3-D complex [(H ₂ O)Cd(EB,4-TDTA)Cd(H ₂ O)]. Potentiometric and ¹¹³ Cd NMR studies in aqueous solution (3,4-TDTA = 3,4-toluenediamine-N,N,N',N''-tetraacetate). <i>Dalton Transactions RSC</i> , 2001 , 1559-1565 | | 4 |
| 12 | Synthesis, crystal structure and magnetic properties of the malonato-bridged bimetallic chain [Mn(II)Cu(II)(mal) ₂ (H ₂ O) ₄] ₂ ·H ₂ O. <i>Inorganica Chimica Acta</i> , 2000 , 298, 202-208 | 2.7 | 63 |
| 11 | Synthesis, crystal structure and magnetic properties of the three-dimensional compound [Na ₂ Ni(mal) ₂ (H ₂ O) ₆] _n (H ₂ mal=malonic acid). <i>Inorganica Chimica Acta</i> , 2000 , 298, 245-250 | 2.7 | 49 |
| 10 | Ferromagnetism in malonato-bridged copper(II) complexes. Synthesis, crystal structures, and magnetic properties of [[Cu(H ₂ O) ₃][Cu(mal) ₂ (H ₂ O)]] _n and [[Cu(H ₂ O) ₄][Cu(mal) ₂ (H ₂ O)]] _n and [[Cu(mal) ₂ (H ₂ O) ₂][Cu(mal) ₂ (H ₂ O) ₂]] _n (H ₂ mal = malonic acid). <i>Inorganic Chemistry</i> , 2000 , 39, 1363-70 | 5.1 | 202 |
| 9 | Solution studies of complexes of iron(III) with iminodiacetic, alkyl-substituted iminodiacetic and nitrilotriacetic acids by potentiometry and cyclic voltammetry. <i>Inorganica Chimica Acta</i> , 1999 , 291, 158-165 | 2.7 | 29 |
| 8 | Coordinating ability of ligands derived from phenylenediamines. <i>Coordination Chemistry Reviews</i> , 1999 , 193-195, 857-911 | 23.2 | 41 |
| 7 | Coordinating ability of phenylenediamines. <i>Coordination Chemistry Reviews</i> , 1999 , 193-195, 913-939 | 23.2 | 75 |
| 6 | Ferromagnetic Ordering, Anisotropy, and Spin Reorientation for the Cyano-Bridged Bimetallic Compound Mn ₂ (H ₂ O) ₅ Mo(CN) ₇ ·4H ₂ O (β-Phase). <i>Journal of the American Chemical Society</i> , 1998 , 120, 13088-13095 | 16.4 | 125 |
| 5 | Potentiometric studies on the formation and dissociation of the L-cysteine complexes of di-μ-sulfido and di-μ-oxo molybdenum(V) [Mo ₂ O ₂ (μ-S) ₂ (cys) ₂] ₂ and [Mo ₂ O ₂ (μ-O) ₂ (cys) ₂] ₂ . <i>Journal of the Chemical Society Dalton Transactions</i> , 1998 , 2723-2726 | | 3 |

| | | | |
|---|--|-----|-----|
| 4 | Crystal structure, ferromagnetic ordering and magnetic anisotropy for two cyano-bridged bimetallic compounds of formula $Mn_2(H_2O)_5Mo(CN)_7 \cdot H_2O$. <i>Chemical Communications</i> , 1998 , 953-954 | 5.8 | 58 |
| 3 | Magnetic Properties of the Two-Dimensional Bimetallic Compounds $(NBu_4)[MIRuIII(ox)_3]$ (NBu_4 = Tetra-n-butylammonium; M = Mn, Fe, Cu; ox = Oxalate). <i>Inorganic Chemistry</i> , 1998 , 37, 679-684 | 5.1 | 114 |
| 2 | Tetramethyl Carboxylic Acids Derived from o-Phenylenediamines as Sequestering Agents for Iron(III): Thermodynamic Studies. X-ray Crystal Structure of Sodium Aqua(4-chloro-1,2-phenylenediamine-N,N,N',N')tetraacetato)ferrate(III)Water (1/1.5). <i>Inorganic Chemistry</i> , 1997 , 36, 4108-4114 | 5.1 | 32 |
| 1 | Potassium Aqua(3,4-toluenediamine-N,N,N',N')tetraacetato)ferrate(III)Water (1/1.5). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 1996 , 52, 1618-1621 | | 1 |