

Sandra Ribeiro

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5296451/publications.pdf>

Version: 2024-02-01

23
papers

285
citations

933264

10
h-index

887953

17
g-index

23
all docs

23
docs citations

23
times ranked

484
citing authors

#	ARTICLE	IF	CITATIONS
1	Hepcidin in chronic kidney disease anemia. <i>Vitamins and Hormones</i> , 2019, 110, 243-264.	0.7	14
2	Influence of the 6-month physical activity programs on renal function in obese boys. <i>Pediatric Research</i> , 2018, 83, 1011-1015.	1.1	3
3	The HIF System Response to ESA Therapy in CKD Anemia. , 2017, , .		0
4	Resistance to Recombinant Human Erythropoietin Therapy in a Rat Model of Chronic Kidney Disease Associated Anemia. <i>International Journal of Molecular Sciences</i> , 2016, 17, 28.	1.8	11
5	SP313 LIVER IRON IS A MAJOR REGULATOR OF HEPCIDIN GENE EXPRESSION VIA BMP/SMAD PATHWAY IN A RAT MODEL OF CHRONIC RENAL FAILURE UNDER TREATMENT WITH HIGH rHuEPO DOSES. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i194-i194.	0.4	1
6	Renal risk-benefit determinants of recombinant human erythropoietin therapy in the remnant kidney rat model "hypertension, anaemia, inflammation and drug dose. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2016, 43, 343-354.	0.9	10
7	Pathological and molecular mechanisms underlying resistance to recombinant human erythropoietin therapy in the remnant kidney rat model of chronic kidney disease associated anemia. <i>Biochimie</i> , 2016, 125, 150-162.	1.3	11
8	Impaired renal endothelial nitric oxide synthase and reticulocyte production as modulators of hypertension induced by rHuEPO in the rat. <i>Life Sciences</i> , 2016, 151, 147-156.	2.0	4
9	Recombinant human erythropoietin-induced erythropoiesis regulates hepcidin expression over iron status in the rat. <i>Blood Cells, Molecules, and Diseases</i> , 2016, 59, 63-70.	0.6	6
10	Iron therapy in chronic kidney disease: Recent changes, benefits and risks. <i>Blood Reviews</i> , 2016, 30, 65-72.	2.8	28
11	Liver iron is a major regulator of hepcidin gene expression via <sc>BMP/SMAD</sc> pathway in a rat model of chronic renal failure under treatment with high r<sc>H</sc>u<sc>EPO</sc> doses. <i>BioFactors</i> , 2016, 42, 296-306.	2.6	8
12	Iron-Hepcidin Dysmetabolism, Anemia and Renal Hypoxia, Inflammation and Fibrosis in the Remnant Kidney Rat Model. <i>PLoS ONE</i> , 2015, 10, e0124048.	1.1	33
13	Potential Cardiovascular Risk Protection of Bilirubin in End-Stage Renal Disease Patients under Hemodialysis. <i>BioMed Research International</i> , 2014, 2014, 1-9.	0.9	12
14	Risk Factors for Mortality in Hemodialysis Patients: Two-Year Follow-Up Study. <i>Disease Markers</i> , 2013, 35, 791-798.	0.6	45
15	Markers of Increased Cardiovascular Risk in Postmenopausal Women: Focus on Oxidized-LDL and HDL Subpopulations. <i>Disease Markers</i> , 2013, 35, 85-96.	0.6	32
16	Circulating cell-free DNA levels in hemodialysis patients and its association with inflammation, iron metabolism, and rhEPO doses. <i>Hemodialysis International</i> , 2013, 17, n/a-n/a.	0.4	11
17	Body mass index and resistance to recombinant human erythropoietin therapy in maintenance hemodialysis patients. <i>Renal Failure</i> , 2013, 35, 1392-1398.	0.8	10
18	Vascular Access versus the Effect of Statins on Inflammation and Fibrinolysis in Renal Dialysis Patients. <i>Journal of Vascular Access</i> , 2013, 14, 335-341.	0.5	3

#	ARTICLE	IF	CITATIONS
19	Comparison of Bio-Plex measurements with standard techniques. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012, 50, 399-402.	1.4	1
20	Major Determinants of BMP-2 Serum Levels in Hemodialysis Patients. <i>Renal Failure</i> , 2012, 34, 1355-1358.	0.8	4
21	Main Determinants of PON1 Activity in Hemodialysis Patients. <i>American Journal of Nephrology</i> , 2012, 36, 317-323.	1.4	16
22	Adiponectin is an independent predictor of tissue plasminogen activator levels in patients under haemodialysis. <i>Scandinavian Journal of Urology and Nephrology</i> , 2012, 46, 461-465.	1.4	1
23	Oxidized low-density lipoprotein and lipoprotein(a) levels in chronic kidney disease patients under hemodialysis: Influence of adiponectin and of a polymorphism in the apolipoprotein(a) gene. <i>Hemodialysis International</i> , 2012, 16, 481-490.	0.4	21