

Lise Madsen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/530215/publications.pdf>

Version: 2024-02-01

109
papers

13,415
citations

50170

46
h-index

24179

110
g-index

112
all docs

112
docs citations

112
times ranked

19864
citing authors

#	ARTICLE	IF	CITATIONS
1	Dynamics and Stabilization of the Human Gut Microbiome during the First Year of Life. <i>Cell Host and Microbe</i> , 2015, 17, 690-703.	5.1	2,276
2	Gut microbiome and serum metabolome alterations in obesity and after weight-loss intervention. <i>Nature Medicine</i> , 2017, 23, 859-868.	15.2	1,074
3	The gut microbiome in atherosclerotic cardiovascular disease. <i>Nature Communications</i> , 2017, 8, 845.	5.8	1,029
4	Metagenomic analysis of faecal microbiome as a tool towards targeted non-invasive biomarkers for colorectal cancer. <i>Gut</i> , 2017, 66, 70-78.	6.1	865
5	The microbiota continuum along the female reproductive tract and its relation to uterine-related diseases. <i>Nature Communications</i> , 2017, 8, 875.	5.8	572
6	Peroxisome Proliferator-activated Receptor δ Activators Improve Insulin Sensitivity and Reduce Adiposity. <i>Journal of Biological Chemistry</i> , 2000, 275, 16638-16642.	1.6	554
7	A catalog of the mouse gut metagenome. <i>Nature Biotechnology</i> , 2015, 33, 1103-1108.	9.4	422
8	A reference gene catalogue of the pig gut microbiome. <i>Nature Microbiology</i> , 2016, 1, 16161.	5.9	416
9	Persistent Organic Pollutant Exposure Leads to Insulin Resistance Syndrome. <i>Environmental Health Perspectives</i> , 2010, 118, 465-471.	2.8	326
10	Analyses of gut microbiota and plasma bile acids enable stratification of patients for antidiabetic treatment. <i>Nature Communications</i> , 2017, 8, 1785.	5.8	312
11	Microarray Analyses during Adipogenesis: Understanding the Effects of Wnt Signaling on Adipogenesis and the Roles of Liver X Receptor δ in Adipocyte Metabolism. <i>Molecular and Cellular Biology</i> , 2002, 22, 5989-5999.	1.1	227
12	Regulation of adipocyte differentiation and function by polyunsaturated fatty acids. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2005, 1740, 266-286.	1.8	218
13	Dietary Linoleic Acid Elevates Endogenous 2-AG and Anandamide and Induces Obesity. <i>Obesity</i> , 2012, 20, 1984-1994.	1.5	200
14	Chronic <i>Trichuris muris</i> Infection Decreases Diversity of the Intestinal Microbiota and Concomitantly Increases the Abundance of Lactobacilli. <i>PLoS ONE</i> , 2015, 10, e0125495.	1.1	190
15	UCP1 Induction during Recruitment of Brown Adipocytes in White Adipose Tissue Is Dependent on Cyclooxygenase Activity. <i>PLoS ONE</i> , 2010, 5, e11391.	1.1	174
16	Interplay between food and gut microbiota in health and disease. <i>Food Research International</i> , 2019, 115, 23-31.	2.9	168
17	Eicosapentaenoic and docosahexaenoic acid affect mitochondrial and peroxisomal fatty acid oxidation in relation to substrate preference. <i>Lipids</i> , 1999, 34, 951-963.	0.7	160
18	Cyclic AMP (cAMP)-Mediated Stimulation of Adipocyte Differentiation Requires the Synergistic Action of Epac- and cAMP-Dependent Protein Kinase-Dependent Processes. <i>Molecular and Cellular Biology</i> , 2008, 28, 3804-3816.	1.1	136

#	ARTICLE	IF	CITATIONS
19	Chronic Consumption of Farmed Salmon Containing Persistent Organic Pollutants Causes Insulin Resistance and Obesity in Mice. PLoS ONE, 2011, 6, e25170.	1.1	133
20	High-fat feeding rather than obesity drives taxonomical and functional changes in the gut microbiota in mice. Microbiome, 2017, 5, 43.	4.9	132
21	Adipocyte differentiation of 3T3-L1 preadipocytes is dependent on lipoxygenase activity during the initial stages of the differentiation process. Biochemical Journal, 2003, 375, 539-549.	1.7	118
22	Transcriptome profiling of brown adipose tissue during cold exposure reveals extensive regulation of glucose metabolism. American Journal of Physiology - Endocrinology and Metabolism, 2015, 308, E380-E392.	1.8	105
23	Fish protein hydrolysate elevates plasma bile acids and reduces visceral adipose tissue mass in rats. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2009, 1791, 254-262.	1.2	98
24	Circulating FGF21 in humans is potently induced by short term overfeeding of carbohydrates. Molecular Metabolism, 2017, 6, 22-29.	3.0	95
25	Links between Dietary Protein Sources, the Gut Microbiota, and Obesity. Frontiers in Physiology, 2017, 8, 1047.	1.3	83
26	Lower levels of Persistent Organic Pollutants, metals and the marine omega 3-fatty acid DHA in farmed compared to wild Atlantic salmon (<i>Salmo salar</i>). Environmental Research, 2017, 155, 49-59.	3.7	76
27	Two distinct metacommunities characterize the gut microbiota in Crohn's disease patients. GigaScience, 2017, 6, 1-11.	3.3	75
28	All-Trans Retinoic Acid Increases Oxidative Metabolism in Mature Adipocytes. Cellular Physiology and Biochemistry, 2007, 20, 1061-1072.	1.1	72
29	cAMP-dependent Signaling Regulates the Adipogenic Effect of n-6 Polyunsaturated Fatty Acids. Journal of Biological Chemistry, 2008, 283, 7196-7205.	1.6	72
30	Dietary Linoleic Acid Elevates the Endocannabinoids 2-AG and Anandamide and Promotes Weight Gain in Mice Fed a Low Fat Diet. Lipids, 2014, 49, 59-69.	0.7	70
31	The metagenome of the female upper reproductive tract. GigaScience, 2018, 7, .	3.3	68
32	Carnitine palmitoyltransferase I, carnitine palmitoyltransferase II, and Acyl-CoA oxidase activities in atlantic salmon (<i>Salmo salar</i>). Lipids, 1998, 33, 923-930.	0.7	66
33	Dietary linoleic acid elevates endogenous 2-arachidonoylglycerol and anandamide in Atlantic salmon (<i>Salmo salar</i>) and mice, and induces weight gain and inflammation in mice. British Journal of Nutrition, 2013, 109, 1508-1517.	1.2	66
34	Effects of geography and species variation on selenium and mercury molar ratios in Northeast Atlantic marine fish communities. Science of the Total Environment, 2019, 652, 1482-1496.	3.9	65
35	Arachidonic acid-dependent inhibition of adipocyte differentiation requires PKA activity and is associated with sustained expression of cyclooxygenases. Journal of Lipid Research, 2003, 44, 2320-2330.	2.0	61
36	Contaminant levels in Norwegian farmed Atlantic salmon (<i>Salmo salar</i>) in the 13-year period from 1999 to 2011. Environment International, 2015, 74, 274-280.	4.8	61

#	ARTICLE	IF	CITATIONS
37	Tetradecylthioacetic acid inhibits growth of rat glioma cells ex vivo and in vivo via PPAR-dependent and PPAR-independent pathways. <i>Carcinogenesis</i> , 2001, 22, 1747-1755.	1.3	59
38	The protein source determines the potential of high protein diets to attenuate obesity development in C57BL/6J mice. <i>Adipocyte</i> , 2016, 5, 196-211.	1.3	59
39	The tumor suppressors pRB and p53 as regulators of adipocyte differentiation and function. <i>Expert Opinion on Therapeutic Targets</i> , 2009, 13, 235-246.	1.5	56
40	Nutritional Regulation of Bile Acid Metabolism Is Associated with Improved Pathological Characteristics of the Metabolic Syndrome. <i>Journal of Biological Chemistry</i> , 2011, 286, 28382-28395.	1.6	55
41	Effect of a long-term high-protein diet on survival, obesity development, and gut microbiota in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 310, E886-E899.	1.8	55
42	Proliferation of mitochondria and gene expression of carnitine palmitoyltransferase and fatty acyl-CoA oxidase in rat skeletal muscle, heart and liver by hypolipidemic fatty acids. <i>Biology of the Cell</i> , 2000, 92, 317-329.	0.7	53
43	Establishment of a <i>Macaca fascicularis</i> gut microbiome gene catalog and comparison with the human, pig, and mouse gut microbiomes. <i>GigaScience</i> , 2018, 7, .	3.3	53
44	Hydrolyzed Casein Reduces Diet-Induced Obesity in Male C57BL/6J Mice. <i>Journal of Nutrition</i> , 2013, 143, 1367-1375.	1.3	50
45	Mechanisms Preserving Insulin Action during High Dietary Fat Intake. <i>Cell Metabolism</i> , 2019, 29, 50-63.e4.	7.2	50
46	Lipidomic profiling reveals distinct differences in plasma lipid composition in healthy, prediabetic, and type 2 diabetic individuals. <i>GigaScience</i> , 2017, 6, 1-12.	3.3	49
47	In contrast with docosahexaenoic acid, eicosapentaenoic acid and hypolipidaemic derivatives decrease hepatic synthesis and secretion of triacylglycerol by decreased diacylglycerol acyltransferase activity and stimulation of fatty acid oxidation. <i>Biochemical Journal</i> , 1999, 343, 191.	1.7	47
48	Sucrose Counteracts the Anti-Inflammatory Effect of Fish Oil in Adipose Tissue and Increases Obesity Development in Mice. <i>PLoS ONE</i> , 2011, 6, e21647.	1.1	47
49	Opposite Regulation of Insulin Sensitivity by Dietary Lipid Versus Carbohydrate Excess. <i>Diabetes</i> , 2017, 66, 2583-2595.	0.3	46
50	The importance of dietary modulation of cAMP and insulin signaling in adipose tissue and the development of obesity. <i>Annals of the New York Academy of Sciences</i> , 2010, 1190, 1-14.	1.8	45
51	Epidermis-Type Lipoxygenase 3 Regulates Adipocyte Differentiation and Peroxisome Proliferator-Activated Receptor β Activity. <i>Molecular and Cellular Biology</i> , 2010, 30, 4077-4091.	1.1	45
52	CRISPR/Cas9-Mediated Genome Editing-Challenges and Opportunities. <i>Frontiers in Genetics</i> , 2018, 9, 240.	1.1	45
53	A comparative analysis of the intestinal metagenomes present in guinea pigs (<i>Cavia porcellus</i>) and humans (<i>Homo sapiens</i>). <i>BMC Genomics</i> , 2012, 13, 514.	1.2	43
54	Urinary Loss of Tricarboxylic Acid Cycle Intermediates As Revealed by Metabolomics Studies: An Underlying Mechanism to Reduce Lipid Accretion by Whey Protein Ingestion?. <i>Journal of Proteome Research</i> , 2014, 13, 2560-2570.	1.8	42

#	ARTICLE	IF	CITATIONS
55	Haploinsufficiency of the retinoblastoma protein gene reduces diet-induced obesity, insulin resistance, and hepatosteatosis in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2009, 297, E184-E193.	1.8	41
56	Scallop protein with endogenous high taurine and glycine content prevents high-fat, high-sucrose-induced obesity and improves plasma lipid profile in male C57BL/6J mice. <i>Amino Acids</i> , 2014, 46, 1659-1671.	1.2	41
57	FFAR4 (GPR120) Signaling Is Not Required for Anti-Inflammatory and Insulin-Sensitizing Effects of Omega-3 Fatty Acids. <i>Mediators of Inflammation</i> , 2016, 2016, 1-12.	1.4	40
58	Dietary eicosapentaenoic acid supplementation accentuates hepatic triglyceride accumulation in mice with impaired fatty acid oxidation capacity. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2013, 1831, 291-299.	1.2	39
59	High-glycemic index carbohydrates abrogate the antiobesity effect of fish oil in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012, 302, E1097-E1112.	1.8	37
60	Intake of Farmed Atlantic Salmon Fed Soybean Oil Increases Insulin Resistance and Hepatic Lipid Accumulation in Mice. <i>PLoS ONE</i> , 2013, 8, e53094.	1.1	37
61	Impact of mitochondrial β -oxidation in fatty acid-mediated inhibition of glioma cell proliferation. <i>Journal of Lipid Research</i> , 2003, 44, 118-127.	2.0	36
62	Dynamic Regulation of Genes Involved in Mitochondrial DNA Replication and Transcription during Mouse Brown Fat Cell Differentiation and Recruitment. <i>PLoS ONE</i> , 2009, 4, e8458.	1.1	36
63	Visualization and Quantification of Browning Using a <i>Ucp1</i> -2A-Luciferase Knock-in Mouse Model. <i>Diabetes</i> , 2017, 66, 407-417.	0.3	35
64	A catalog of microbial genes from the bovine rumen unveils a specialized and diverse biomass-degrading environment. <i>GigaScience</i> , 2020, 9, .	3.3	35
65	Obesity is associated with depot-specific alterations in adipocyte DNA methylation and gene expression. <i>Adipocyte</i> , 2017, 6, 124-133.	1.3	34
66	Indomethacin Treatment Prevents High Fat Diet-induced Obesity and Insulin Resistance but Not Glucose Intolerance in C57BL/6J Mice. <i>Journal of Biological Chemistry</i> , 2014, 289, 16032-16045.	1.6	33
67	Oceans and Human Health (OHH): a European Perspective from the Marine Board of the European Science Foundation (Marine Board-ESF). <i>Microbial Ecology</i> , 2013, 65, 889-900.	1.4	32
68	Intake of a Western diet containing cod instead of pork alters fatty acid composition in tissue phospholipids and attenuates obesity and hepatic lipid accumulation in mice. <i>Journal of Nutritional Biochemistry</i> , 2016, 33, 119-127.	1.9	32
69	Diet-induced obesity, energy metabolism and gut microbiota in C57BL/6J mice fed Western diets based on lean seafood or lean meat mixtures. <i>Journal of Nutritional Biochemistry</i> , 2016, 31, 127-136.	1.9	32
70	The elusive endogenous adipogenic PPAR β agonists: Lining up the suspects. <i>Progress in Lipid Research</i> , 2016, 61, 149-162.	5.3	32
71	Mitochondrial 3-hydroxy-3-methylglutaryl coenzyme A synthase and carnitine palmitoyltransferase II as potential control sites for ketogenesis during mitochondrion and peroxisome proliferation. <i>Biochemical Pharmacology</i> , 1999, 57, 1011-1019.	2.0	30
72	Intake of farmed Atlantic salmon fed soybean oil increases hepatic levels of arachidonic acid-derived oxylipins and ceramides in mice. <i>Journal of Nutritional Biochemistry</i> , 2015, 26, 585-595.	1.9	30

#	ARTICLE	IF	CITATIONS
73	ERR β enhances UCP1 expression and fatty acid oxidation in brown adipocytes. <i>Obesity</i> , 2013, 21, 516-524.	1.5	29
74	Factors influencing risk assessments of brominated flame-retardants; evidence based on seafood from the North East Atlantic Ocean. <i>Environment International</i> , 2018, 119, 544-557.	4.8	28
75	Depot-Dependent Effects of Adipose Tissue Explants on Co-Cultured Hepatocytes. <i>PLoS ONE</i> , 2011, 6, e20917.	1.1	27
76	Low doses of eicosapentaenoic acid, docosahexaenoic acid, and hypolipidemic eicosapentaenoic acid derivatives have no effect on lipid peroxidation in plasma. <i>Lipids</i> , 1998, 33, 1131-1137.	0.7	26
77	p53 regulates expression of uncoupling protein 1 through binding and repression of PPAR β coactivator-1 α . <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 310, E116-E128.	1.8	26
78	Iodine Status and Thyroid Function in a Group of Seaweed Consumers in Norway. <i>Nutrients</i> , 2020, 12, 3483.	1.7	26
79	β -oxidation modulates metabolic competition between eicosapentaenoic acid and arachidonic acid regulating prostaglandin E2 synthesis in rat hepatocytes/Kupffer cells. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2010, 1801, 526-536.	1.2	25
80	Glucose-induced lipogenesis in pancreatic β -cells is dependent on SREBP-1. <i>Molecular and Cellular Endocrinology</i> , 2005, 240, 94-106.	1.6	23
81	Dietary fat drives whole-body insulin resistance and promotes intestinal inflammation independent of body weight gain. <i>Metabolism: Clinical and Experimental</i> , 2016, 65, 1706-1719.	1.5	22
82	ADD1/SREBP1c activates the PGC1- α promoter in brown adipocytes. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2010, 1801, 421-429.	1.2	20
83	Early modulation of genes encoding peroxisomal and mitochondrial β -oxidation enzymes by 3-thia fatty acids. <i>Biochemical Pharmacology</i> , 1998, 56, 1571-1582.	2.0	19
84	PPAR β ligand production is tightly linked to clonal expansion during initiation of adipocyte differentiation. <i>Journal of Lipid Research</i> , 2014, 55, 2491-2500.	2.0	19
85	Lean Seafood Intake Reduces Postprandial C-peptide and Lactate Concentrations in Healthy Adults in a Randomized Controlled Trial with a Crossover Design. <i>Journal of Nutrition</i> , 2016, 146, 1027-1034.	1.3	19
86	A safflower oil based high-fat/high-sucrose diet modulates the gut microbiota and liver phospholipid profiles associated with early glucose intolerance in the absence of tissue inflammation. <i>Molecular Nutrition and Food Research</i> , 2017, 61, 1600528.	1.5	19
87	Growth reduction in glioma cells after treatment with tetradecylthioacetic acid: changes in fatty acid metabolism and oxidative status. <i>Biochemical Pharmacology</i> , 2001, 61, 639-649.	2.0	18
88	3-thia fatty acid treatment, in contrast to eicosapentaenoic acid and starvation, induces gene expression of carnitine palmitoyltransferase-II in rat liver. <i>Lipids</i> , 1999, 34, 447-456.	0.7	17
89	Multi-block PCA and multi-compartmental study of the metabolic responses to intake of hydrolysed versus intact casein in C57BL/6J mice by NMR-based metabolomics. <i>Metabolomics</i> , 2014, 10, 938-949.	1.4	14
90	Marine fatty acids aggravate hepatotoxicity of β -HBCD in juvenile female BALB/c mice. <i>Food and Chemical Toxicology</i> , 2016, 97, 411-423.	1.8	14

#	ARTICLE	IF	CITATIONS
91	Macronutrient composition determines accumulation of persistent organic pollutants from dietary exposure in adipose tissue of mice. <i>Journal of Nutritional Biochemistry</i> , 2016, 27, 307-316.	1.9	14
92	The Impact of Different Animal-Derived Protein Sources on Adiposity and Glucose Homeostasis during Ad Libitum Feeding and Energy Restriction in Already Obese Mice. <i>Nutrients</i> , 2019, 11, 1153.	1.7	14
93	An Expanded Gene Catalog of Mouse Gut Metagenomes. <i>MSphere</i> , 2021, 6, .	1.3	13
94	Dietary Proteins, Brown Fat, and Adiposity. <i>Frontiers in Physiology</i> , 2018, 9, 1792.	1.3	11
95	Methylated eicosapentaenoic acid and tetradecylthioacetic acid: effects on fatty acid metabolism. <i>Biochemical Pharmacology</i> , 1999, 58, 1133-1143.	2.0	10
96	Intake of Hydrolyzed Casein is Associated with Reduced Body Fat Accretion and Enhanced Phase II Metabolism in Obesity Prone C57BL/6J Mice. <i>PLoS ONE</i> , 2015, 10, e0118895.	1.1	10
97	Subchronic dietary exposure to ethoxyquin dimer induces microvesicular steatosis in male BALB/c mice. <i>Food and Chemical Toxicology</i> , 2018, 118, 608-625.	1.8	9
98	Tissue Inhibitor Of Matrix Metalloproteinase-1 Is Required for High-Fat Diet-Induced Glucose Intolerance and Hepatic Steatosis in Mice. <i>PLoS ONE</i> , 2015, 10, e0132910.	1.1	9
99	Fatty fish, hair mercury and cognitive function in Norwegian preschool children: Results from the randomized controlled trial FINS-KIDS. <i>Environment International</i> , 2018, 121, 1098-1105.	4.8	8
100	Dietary intake and adipose tissue content of long-chain n-3 PUFAs and subsequent 5-y change in body weight and waist circumference. <i>American Journal of Clinical Nutrition</i> , 2017, 105, 1148-1157.	2.2	7
101	Effects of Frozen Storage on Phospholipid Content in Atlantic Cod Fillets and the Influence on Diet-Induced Obesity in Mice. <i>Nutrients</i> , 2018, 10, 695.	1.7	7
102	Macronutrients and obesity: views, news and reviews. <i>Future Lipidology</i> , 2008, 3, 43-74.	0.5	6
103	Proteomic analysis of cAMP-mediated signaling during differentiation of 3 T3-L1 preadipocytes. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2014, 1844, 2096-2107.	1.1	6
104	Meals based on cod or veal in combination with high or low glycemic index carbohydrates did not affect diet-induced thermogenesis, appetite sensations, or subsequent energy intake differently. <i>Appetite</i> , 2018, 130, 199-208.	1.8	6
105	Activation of Liver X Receptors Prevents Statin-induced Death of 3T3-L1 Preadipocytes. <i>Journal of Biological Chemistry</i> , 2008, 283, 22723-22736.	1.6	4
106	Long Term Treatment with Tetradecylthioacetic Acid Improves the Antioxidant Status in Obese Zucker (fa/fa) Rats. <i>Drug Metabolism Letters</i> , 2008, 2, 138-145.	0.5	4
107	Dietary n-6 PUFA, carbohydrate:protein ratio and change in body weight and waist circumference: a follow-up study. <i>Public Health Nutrition</i> , 2015, 18, 1317-1323.	1.1	4
108	Salmon in Combination with High Glycemic Index Carbohydrates Increases Diet-Induced Thermogenesis Compared with Salmon with Low Glycemic Index Carbohydrates—An Acute Randomized Cross-Over Meal Test Study. <i>Nutrients</i> , 2019, 11, 365.	1.7	3

#	ARTICLE	IF	CITATIONS
109	The Anti-Obesogenic Effect of Lean Fish Species Is Influenced by the Fatty Acid Composition in Fish Fillets. <i>Nutrients</i> , 2020, 12, 3038.	1.7	0