

Li Jia

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5304093/publications.pdf>

Version: 2024-02-01

56
papers

2,249
citations

236925
25
h-index

214800
47
g-index

57
all docs

57
docs citations

57
times ranked

3275
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Inhibition of autophagy abrogates tumour necrosis factor α induced apoptosis in human T α lymphoblastic leukaemic cells. British Journal of Haematology, 1997, 98, 673-685. | 2.5 | 221 |
| 2 | Apaf-1 protein deficiency confers resistance to cytochrome c dependent apoptosis in human leukemic cells. Blood, 2001, 98, 414-421. | 1.4 | 136 |
| 3 | Inhibition of HIF-1 α by PX-478 enhances the anti-tumor effect of gemcitabine by inducing immunogenic cell death in pancreatic ductal adenocarcinoma. Oncotarget, 2015, 6, 2250-2262. | 1.8 | 110 |
| 4 | Extracellular HMGB1 promotes differentiation of nurse-like cells in chronic lymphocytic leukemia. Blood, 2014, 123, 1709-1719. | 1.4 | 95 |
| 5 | Role of DNA methylation in the suppression of Apaf-1 protein in human leukaemia. Oncogene, 2003, 22, 451-455. | 5.9 | 87 |
| 6 | Blocking Autophagy Prevents Bortezomib-Induced NF- κ B Activation by Reducing I- κ B α Degradation in Lymphoma Cells. PLoS ONE, 2012, 7, e32584. | 2.5 | 87 |
| 7 | Quantitative determination of apoptosis on leukemia cells by infrared spectroscopy. Apoptosis: an International Journal on Programmed Cell Death, 2001, 6, 269-278. | 4.9 | 85 |
| 8 | Dietary flavonoids inhibit the anticancer effects of the proteasome inhibitor bortezomib. Blood, 2008, 112, 3835-3846. | 1.4 | 83 |
| 9 | Subcellular Distribution and Redistribution of Bcl-2 Family Proteins in Human Leukemia Cells Undergoing Apoptosis. Blood, 1999, 93, 2353-2359. | 1.4 | 80 |
| 10 | Bcl-2 Inhibitors Sensitize Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Induced Apoptosis by Uncoupling of Mitochondrial Respiration in Human Leukemic CEM Cells. Cancer Research, 2004, 64, 3607-3616. | 0.9 | 79 |
| 11 | Bortezomib blocks Bax degradation in malignant B cells during treatment with TRAIL. Blood, 2008, 111, 2797-2805. | 1.4 | 79 |
| 12 | Dynamin-related protein Drp1 is required for Bax translocation to mitochondria in response to irradiation-induced apoptosis. Oncotarget, 2015, 6, 22598-22612. | 1.8 | 74 |
| 13 | Bax translocation is crucial for the sensitivity of leukaemic cells to etoposide-induced apoptosis. Oncogene, 2001, 20, 4817-4826. | 5.9 | 73 |
| 14 | Bax conformational change is a crucial step for PUMA-mediated apoptosis in human leukemia. Biochemical and Biophysical Research Communications, 2003, 310, 956-962. | 2.1 | 67 |
| 15 | Novel HDAC inhibitor Chidamide synergizes with Rituximab to inhibit diffuse large B-cell lymphoma tumour growth by upregulating CD20. Cell Death and Disease, 2020, 11, 20. | 6.3 | 62 |
| 16 | Role of Smac in human leukaemic cell apoptosis and proliferation. Oncogene, 2003, 22, 1589-1599. | 5.9 | 54 |
| 17 | CD160 signaling mediates PI3K-dependent survival and growth signals in chronic lymphocytic leukemia. Blood, 2010, 115, 3079-3088. | 1.4 | 48 |
| 18 | Periostin and CA242 as potential diagnostic serum biomarkers complementing CA19.9 in detecting pancreatic cancer. Cancer Science, 2018, 109, 2841-2851. | 3.9 | 47 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Mitochondrial electron transport chain activity, but not ATP synthesis, is required for drug-induced apoptosis in human leukaemic cells: a possible novel mechanism of regulating drug resistance. <i>British Journal of Haematology</i> , 1997, 98, 686-698. | 2.5 | 42 |
| 20 | Single nucleotide polymorphism in the microRNA-199a binding site of HIF1A gene is associated with pancreatic ductal adenocarcinoma risk and worse clinical outcomes. <i>Oncotarget</i> , 2016, 7, 13717-13729. | 1.8 | 40 |
| 21 | Mitochondrial ultracondensation, but not swelling, is involved in TNF α -induced apoptosis in human T-lymphoblastic leukaemic cells. <i>Leukemia Research</i> , 1997, 21, 973-983. | 0.8 | 34 |
| 22 | Blockade of HMGB1 signaling pathway by ethyl pyruvate inhibits tumor growth in diffuse large B-cell lymphoma. <i>Cell Death and Disease</i> , 2019, 10, 330. | 6.3 | 29 |
| 23 | Liposomal encapsulation diminishes daunorubicin-induced generation of reactive oxygen species, depletion of ATP and necrotic cell death in human leukaemic cells. <i>British Journal of Haematology</i> , 2002, 117, 333-342. | 2.5 | 28 |
| 24 | Increase in the ratio of mitochondrial Bax/Bcl-XL induces Bax activation in human leukemic K562 cell line. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2004, 9, 377-384. | 4.9 | 28 |
| 25 | Increased proteasomal degradation of Bax is a common feature of poor prognosis chronic lymphocytic leukemia. <i>Blood</i> , 2008, 111, 2790-2796. | 1.4 | 28 |
| 26 | HIF-2-dependent expression of stem cell factor promotes metastasis in hepatocellular carcinoma. <i>Cancer Letters</i> , 2017, 393, 113-124. | 7.2 | 26 |
| 27 | Ethyl pyruvate suppresses the growth, invasion and migration and induces the apoptosis of non-small cell lung cancer cells via the HMGB1/RAGE axis and the NF κ B/STAT3 pathway. <i>Oncology Reports</i> , 2019, 42, 817-825. | 2.6 | 26 |
| 28 | BH3-domain mimetic compound BH3I-2 α induces rapid damage to the inner mitochondrial membrane prior to the cytochrome c release from mitochondria. <i>British Journal of Haematology</i> , 2003, 121, 332-340. | 2.5 | 25 |
| 29 | STAT3 and NF κ B cooperatively control <i>in vitro</i> spontaneous apoptosis and poor chemo-responsiveness in patients with chronic lymphocytic leukemia. <i>Oncotarget</i> , 2016, 7, 32031-32045. | 1.8 | 24 |
| 30 | Stem cell factor is a novel independent prognostic biomarker for hepatocellular carcinoma after curative resection. <i>Carcinogenesis</i> , 2014, 35, 2283-2290. | 2.8 | 23 |
| 31 | Dysregulation of autophagy in human follicular lymphoma is independent of overexpression of BCL-2. <i>Oncotarget</i> , 2014, 5, 11653-11668. | 1.8 | 22 |
| 32 | TRAIL-Induced Apoptosis in Type I Leukemic Cells Is Not Enhanced by Overexpression of Bax. <i>Biochemical and Biophysical Research Communications</i> , 2001, 283, 1037-1045. | 2.1 | 21 |
| 33 | Serum level of ANGPTL4 as a potential biomarker in renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017, 35, 279-285. | 1.6 | 21 |
| 34 | Increased autocrine interleukin-6 production is significantly associated with worse clinical outcome in patients with chronic lymphocytic leukemia. <i>Journal of Cellular Physiology</i> , 2019, 234, 13994-14006. | 4.1 | 21 |
| 35 | Comprehensive Analysis of lncRNA-Mediated ceRNA Crosstalk and Identification of Prognostic Biomarkers in Wilms' Tumor. <i>BioMed Research International</i> , 2020, 2020, 1-13. | 1.9 | 20 |
| 36 | Rituximab-induced HMGB1 release is associated with inhibition of STAT3 activity in human diffuse large B-cell lymphoma. <i>Oncotarget</i> , 2015, 6, 27816-27831. | 1.8 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Why bortezomib cannot go with 'green'?. Cancer Biology and Medicine, 2013, 10, 206-13. | 3.0 | 19 |
| 38 | c-IAP1 Blocks TNF α -Mediated Cytotoxicity Upstream of Caspase-Dependent and -Independent Mitochondrial Events in Human Leukemic Cells. Biochemical and Biophysical Research Communications, 2001, 287, 181-189. | 2.1 | 18 |
| 39 | Lower expression of Bax predicts poor clinical outcome in patients with glioma after curative resection and radiotherapy/chemotherapy. Journal of Neuro-Oncology, 2019, 141, 71-81. | 2.9 | 18 |
| 40 | CD126 and Targeted Therapy with Tocilizumab in Chronic Lymphocytic Leukemia. Clinical Cancer Research, 2016, 22, 2462-2469. | 7.0 | 17 |
| 41 | 8-Cl-adenosine mediated cytotoxicity and sensitization of T-lymphoblastic leukemia cells to TNF α -induced apoptosis is via inactivation of NF- κ B. Leukemia Research, 2001, 25, 423-431. | 0.8 | 15 |
| 42 | Constitutive levels of cAMP-dependent protein kinase activity determine sensitivity of human multidrug-resistant leukaemic cell lines to growth inhibition and apoptosis by forskolin and tumour necrosis factor alpha. British Journal of Haematology, 2000, 108, 565-573. | 2.5 | 14 |
| 43 | Modulation of surface TNF expression by human leukaemic cells alters their sensitivity to exogenous TNF. Leukemia Research, 1996, 20, 47-55. | 0.8 | 13 |
| 44 | Pgp-positive leukaemic cells have increased mtDNA but no increased rate of proliferation. British Journal of Haematology, 1999, 107, 861-869. | 2.5 | 13 |
| 45 | UNC5D , suppressed by promoter hypermethylation, inhibits cell metastasis by activating death-associated protein kinase 1 in prostate cancer. Cancer Science, 2019, 110, 1244-1255. | 3.9 | 12 |
| 46 | BCR signaling contributes to autophagy regulation in chronic lymphocytic leukemia. Leukemia, 2020, 34, 640-644. | 7.2 | 12 |
| 47 | TNF-mediated killing of human leukaemic cells: Effects of endogenous antioxidant levels and TNF α expression in leukaemic cell lines. Leukemia Research, 1995, 19, 187-194. | 0.8 | 10 |
| 48 | Apaf-1XL Is an Inactive Isoform Compared with Apaf-1L. Biochemical and Biophysical Research Communications, 2001, 282, 268-272. | 2.1 | 10 |
| 49 | Generation of reactive oxygen species is not involved in idarubicin-induced apoptosis in human leukaemic cells. British Journal of Haematology, 2001, 115, 817-825. | 2.5 | 7 |
| 50 | The alpha-5 helix of Bax is sensitive to ubiquitin-dependent degradation. Biochemical and Biophysical Research Communications, 2008, 371, 10-15. | 2.1 | 6 |
| 51 | Subcellular Distribution and Redistribution of Bcl-2 Family Proteins in Human Leukemia Cells Undergoing Apoptosis. Blood, 1999, 93, 2353-2359. | 1.4 | 6 |
| 52 | Dangerous power: mitochondria in CLL cells. Blood, 2014, 123, 2596-2597. | 1.4 | 5 |
| 53 | Methylseleninic acid antagonizes the cytotoxic effect of bortezomib in mantle cell lymphoma cell lines through modulation of Bcl-2 family proteins. British Journal of Haematology, 2012, 156, 286-289. | 2.5 | 3 |
| 54 | Activation of Mitochondrial STAT3 Increases Mitochondrial Respiration and Inhibits Oxidative Stress in Chronic Lymphocytic Leukemic Cells. Blood, 2011, 118, 287-287. | 1.4 | 2 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | HMGB1 Activates TLR9/RAGE Signalling Pathway and Sustains Chronic Lymphocytic Leukemic Cell in Vitro Survival. Blood, 2012, 120, 3860-3860. | 1.4 | 1 |
| 56 | Overexpression of HMGB1 Receptor RAGE Is Associated with Worse Clinical Outcome in Patients with Chronic Lymphocytic Leukemia. Blood, 2015, 126, 617-617. | 1.4 | 0 |