


List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5304093/publications.pdf Version: 2024-02-01



#	ARTICLE	IF	CITATIONS
1	BCR signaling contributes to autophagy regulation in chronic lymphocytic leukemia. Leukemia, 2020, 34, 640-644.	7.2	12
2	Novel HDAC inhibitor Chidamide synergizes with Rituximab to inhibit diffuse large B-cell lymphoma tumour growth by upregulating CD20. Cell Death and Disease, 2020, 11, 20.	6.3	62
3	Comprehensive Analysis of IncRNA-Mediated ceRNA Crosstalk and Identification of Prognostic Biomarkers in Wilms' Tumor. BioMed Research International, 2020, 2020, 1-13.	1.9	20
4	Blockade of HMGB1 signaling pathway by ethyl pyruvate inhibits tumor growth in diffuse large B-cell lymphoma. Cell Death and Disease, 2019, 10, 330.	6.3	29
5	UNC5D , suppressed by promoter hypermethylation, inhibits cell metastasis by activating deathâ€associated protein kinase 1 in prostate cancer. Cancer Science, 2019, 110, 1244-1255.	3.9	12
6	Ethyl pyruvate suppresses the growth, invasion and migration and induces the apoptosis of non‑small cell lung cancer cells via the HMGB1/RAGE axis and the NFâ€îºB/STAT3 pathway. Oncology Reports, 2019, 42, 817-825.	2.6	26
7	Lower expression of Bax predicts poor clinical outcome in patients with glioma after curative resection and radiotherapy/chemotherapy. Journal of Neuro-Oncology, 2019, 141, 71-81.	2.9	18
8	Increased autocrine interleukinâ€6 production is significantly associated with worse clinical outcome in patients with chronic lymphocytic leukemia. Journal of Cellular Physiology, 2019, 234, 13994-14006.	4.1	21
9	Periostin and CA242 as potential diagnostic serum biomarkers complementing CA19.9 in detecting pancreatic cancer. Cancer Science, 2018, 109, 2841-2851.	3.9	47
10	HIF-2-dependent expression of stem cell factor promotes metastasis in hepatocellular carcinoma. Cancer Letters, 2017, 393, 113-124.	7.2	26
11	Serum level of ANGPTL4 as a potential biomarker in renal cell carcinoma. Urologic Oncology: Seminars and Original Investigations, 2017, 35, 279-285.	1.6	21
12	Single nucleotide polymorphism in the microRNA-199a binding site of HIF1A gene is associated with pancreatic ductal adenocarcinoma risk and worse clinical outcomes. Oncotarget, 2016, 7, 13717-13729.	1.8	40
13	CD126 and Targeted Therapy with Tocilizumab in Chronic Lymphocytic Leukemia. Clinical Cancer Research, 2016, 22, 2462-2469.	7.0	17
14	STAT3 and NF-κB cooperatively control <i>in vitro</i> spontaneous apoptosis and poor chemo-responsiveness in patients with chronic lymphocytic leukemia. Oncotarget, 2016, 7, 32031-32045.	1.8	24
15	Inhibition of HIF-1α by PX-478 enhances the anti-tumor effect of gemcitabine by inducing immunogenic cell death in pancreatic ductal adenocarcinoma. Oncotarget, 2015, 6, 2250-2262.	1.8	110
16	Dynamin-related protein Drp1 is required for Bax translocation to mitochondria in response to irradiation-induced apoptosis. Oncotarget, 2015, 6, 22598-22612.	1.8	74
17	Rituximab-induced HMGB1 release is associated with inhibition of STAT3 activity in human diffuse large B-cell lymphoma. Oncotarget, 2015, 6, 27816-27831.	1.8	20
18	Overexpression of HMGB1 Receptor RAGE Is Associated with Worse Clinical Outcome in Patients with Chronic Lymphocytic Leukemia. Blood, 2015, 126, 617-617.	1.4	0

Li Jia

#	Article	IF	CITATIONS
19	Extracellular HMGB1 promotes differentiation of nurse-like cells in chronic lymphocytic leukemia. Blood, 2014, 123, 1709-1719.	1.4	95
20	Stem cell factor is a novel independent prognostic biomarker for hepatocellular carcinoma after curative resection. Carcinogenesis, 2014, 35, 2283-2290.	2.8	23
21	Dangerous power: mitochondria in CLL cells. Blood, 2014, 123, 2596-2597.	1.4	5
22	Dysregulation of autophagy in human follicular lymphoma is independent of overexpression of BCL-2. Oncotarget, 2014, 5, 11653-11668.	1.8	22
23	Why bortezomib cannot go with 'green'?. Cancer Biology and Medicine, 2013, 10, 206-13.	3.0	19
24	Blocking Autophagy Prevents Bortezomib-Induced NF-κB Activation by Reducing I-κBα Degradation in Lymphoma Cells. PLoS ONE, 2012, 7, e32584.	2.5	87
25	Methylseleninic acid antagonizes the cytotoxic effect of bortezomib in mantle cell lymphoma cell lines through modulation of Bclâ€⊋ family proteins. British Journal of Haematology, 2012, 156, 286-289.	2.5	3
26	HMGB1 Activates TLR9/RAGE Signalling Pathway and Sustains Chronic Lymphocytic Leukemic Cell in Vitro Survival. Blood, 2012, 120, 3860-3860.	1.4	1
27	Activation of Mitochondrial STAT3 Increases Mitochondrial Respiration and Inhibits Oxidative Stress in Chronic Lymphocytic Leukemic Cells. Blood, 2011, 118, 287-287.	1.4	2
28	CD160 signaling mediates PI3K-dependent survival and growth signals in chronic lymphocytic leukemia. Blood, 2010, 115, 3079-3088.	1.4	48
29	The alpha-5 helix of Bax is sensitive to ubiquitin-dependent degradation. Biochemical and Biophysical Research Communications, 2008, 371, 10-15.	2.1	6
30	Bortezomib blocks Bax degradation in malignant B cells during treatment with TRAIL. Blood, 2008, 111, 2797-2805.	1.4	79
31	Increased proteasomal degradation of Bax is a common feature of poor prognosis chronic lymphocytic leukemia. Blood, 2008, 111, 2790-2796.	1.4	28
32	Dietary flavonoids inhibit the anticancer effects of the proteasome inhibitor bortezomib. Blood, 2008, 112, 3835-3846.	1.4	83
33	Bcl-2 Inhibitors Sensitize Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Induced Apoptosis by Uncoupling of Mitochondrial Respiration in Human Leukemic CEM Cells. Cancer Research, 2004, 64, 3607-3616.	0.9	79
34	Increase in the ratio of mitochondrial Bax/Bcl-XL induces Bax activation in human leukemic K562 cell line. Apoptosis: an International Journal on Programmed Cell Death, 2004, 9, 377-384.	4.9	28
35	BH3-domain mimetic compound BH3I-2′ induces rapid damage to the inner mitochondrial membrane prior to the cytochrome c release from mitochondria. British Journal of Haematology, 2003, 121, 332-340.	2.5	25
36	Role of DNA methylation in the suppression of Apaf-1 protein in human leukaemia. Oncogene, 2003, 22, 451-455.	5.9	87

Li Jia

#	Article	IF	CITATIONS
37	Role of Smac in human leukaemic cell apoptosis and proliferation. Oncogene, 2003, 22, 1589-1599.	5.9	54
38	Bax conformational change is a crucial step for PUMA-mediated apoptosis in human leukemia. Biochemical and Biophysical Research Communications, 2003, 310, 956-962.	2.1	67
39	Liposomal encapsulation diminishes daunorubicin-induced generation of reactive oxygen species, depletion of ATP and necrotic cell death in human leukaemic cells. British Journal of Haematology, 2002, 117, 333-342.	2.5	28
40	Apaf-1XL Is an Inactive Isoform Compared with Apaf-1L. Biochemical and Biophysical Research Communications, 2001, 282, 268-272.	2.1	10
41	TRAIL-Induced Apoptosis in Type I Leukemic Cells Is Not Enhanced by Overexpression of Bax. Biochemical and Biophysical Research Communications, 2001, 283, 1037-1045.	2.1	21
42	c-IAP1 Blocks TNFα-Mediated Cytotoxicity Upstream of Caspase-Dependent and -Independent Mitochondrial Events in Human Leukemic Cells. Biochemical and Biophysical Research Communications, 2001, 287, 181-189.	2.1	18
43	Apaf-1 protein deficiency confers resistance to cytochromec–dependent apoptosis in human leukemic cells. Blood, 2001, 98, 414-421.	1.4	136
44	Generation of reactive oxygen species is not involved in idarubicin-induced apoptosis in human leukaemic cells. British Journal of Haematology, 2001, 115, 817-825.	2.5	7
45	Quantitative determination of apoptosis on leukemia cells by infrared spectroscopy. Apoptosis: an International Journal on Programmed Cell Death, 2001, 6, 269-278.	4.9	85
46	Bax translocation is crucial for the sensitivity of leukaemic cells to etoposide-induced apoptosis. Oncogene, 2001, 20, 4817-4826.	5.9	73
47	8-Cl-adenosine mediated cytotoxicity and sensitization of T-lymphoblastic leukemia cells to TNFα-induced apoptosis is via inactivation of NF-κB. Leukemia Research, 2001, 25, 423-431.	0.8	15
48	Constitutive levels of cAMP-dependent protein kinase activity determine sensitivity of human multidrug-resistant leukaemic cell lines to growth inhibition and apoptosis by forskolin and tumour necrosis factor alpha. British Journal of Haematology, 2000, 108, 565-573.	2.5	14
49	Subcellular Distribution and Redistribution of Bcl-2 Family Proteins in Human Leukemia Cells Undergoing Apoptosis. Blood, 1999, 93, 2353-2359.	1.4	80
50	Pgp-positive leukaemic cells have increased mtDNA but no increased rate of proliferation. British Journal of Haematology, 1999, 107, 861-869.	2.5	13
51	Subcellular Distribution and Redistribution of Bcl-2 Family Proteins in Human Leukemia Cells Undergoing Apoptosis. Blood, 1999, 93, 2353-2359.	1.4	6
52	Mitochondrial ultracondensation, but not swelling, is involved in TNFα-induced apoptosis in human T-lymphoblastic leukaemic cells. Leukemia Research, 1997, 21, 973-983.	0.8	34
53	Inhibition of autophagy abrogates tumour necrosis factor α induced apoptosis in human T″ymphoblastic leukaemic cells. British Journal of Haematology, 1997, 98, 673-685.	2.5	221
54	Mitochondrial electron transport chain activity, but not ATP synthesis, is required for drugâ€induced apoptosis in human leukaemic cells: a possible novel mechanism of regulating drug resistance. British Journal of Haematology, 1997, 98, 686-698.	2.5	42

Li Jia

#	Article	IF	CITATIONS
55	Modulation of surface TNF expression by human leukaemic cells alters their sensitivity to exogenous TNF. Leukemia Research, 1996, 20, 47-55.	0.8	13
56	TNF-mediated killing of human leukaemic cells: Effects of endogenous antioxidant levels and TNFα expression in leukaemic cell lines. Leukemia Research, 1995, 19, 187-194.	0.8	10