

# Li Jia

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5304093/publications.pdf>

Version: 2024-02-01

56  
papers

2,249  
citations

236612

25  
h-index

214527

47  
g-index

57  
all docs

57  
docs citations

57  
times ranked

3275  
citing authors

#	ARTICLE	IF	CITATIONS
1	BCR signaling contributes to autophagy regulation in chronic lymphocytic leukemia. <i>Leukemia</i> , 2020, 34, 640-644.	3.3	12
2	Novel HDAC inhibitor Chidamide synergizes with Rituximab to inhibit diffuse large B-cell lymphoma tumour growth by upregulating CD20. <i>Cell Death and Disease</i> , 2020, 11, 20.	2.7	62
3	Comprehensive Analysis of lncRNA-Mediated ceRNA Crosstalk and Identification of Prognostic Biomarkers in Wilms's Tumor. <i>BioMed Research International</i> , 2020, 2020, 1-13.	0.9	20
4	Blockade of HMGB1 signaling pathway by ethyl pyruvate inhibits tumor growth in diffuse large B-cell lymphoma. <i>Cell Death and Disease</i> , 2019, 10, 330.	2.7	29
5	UNC5D, suppressed by promoter hypermethylation, inhibits cell metastasis by activating death-associated protein kinase 1 in prostate cancer. <i>Cancer Science</i> , 2019, 110, 1244-1255.	1.7	12
6	Ethyl pyruvate suppresses the growth, invasion and migration and induces the apoptosis of non-small cell lung cancer cells via the HMGB1/RAGE axis and the NF- $\kappa$ B/STAT3 pathway. <i>Oncology Reports</i> , 2019, 42, 817-825.	1.2	26
7	Lower expression of Bax predicts poor clinical outcome in patients with glioma after curative resection and radiotherapy/chemotherapy. <i>Journal of Neuro-Oncology</i> , 2019, 141, 71-81.	1.4	18
8	Increased autocrine interleukin-6 production is significantly associated with worse clinical outcome in patients with chronic lymphocytic leukemia. <i>Journal of Cellular Physiology</i> , 2019, 234, 13994-14006.	2.0	21
9	Periostin and CA242 as potential diagnostic serum biomarkers complementing CA19.9 in detecting pancreatic cancer. <i>Cancer Science</i> , 2018, 109, 2841-2851.	1.7	47
10	HIF-2-dependent expression of stem cell factor promotes metastasis in hepatocellular carcinoma. <i>Cancer Letters</i> , 2017, 393, 113-124.	3.2	26
11	Serum level of ANGPTL4 as a potential biomarker in renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017, 35, 279-285.	0.8	21
12	Single nucleotide polymorphism in the microRNA-199a binding site of HIF1A gene is associated with pancreatic ductal adenocarcinoma risk and worse clinical outcomes. <i>Oncotarget</i> , 2016, 7, 13717-13729.	0.8	40
13	CD126 and Targeted Therapy with Tocilizumab in Chronic Lymphocytic Leukemia. <i>Clinical Cancer Research</i> , 2016, 22, 2462-2469.	3.2	17
14	STAT3 and NF- $\kappa$ B cooperatively control <i>in vitro</i> spontaneous apoptosis and poor chemo-responsiveness in patients with chronic lymphocytic leukemia. <i>Oncotarget</i> , 2016, 7, 32031-32045.	0.8	24
15	Inhibition of HIF-1 $\alpha$ by PX-478 enhances the anti-tumor effect of gemcitabine by inducing immunogenic cell death in pancreatic ductal adenocarcinoma. <i>Oncotarget</i> , 2015, 6, 2250-2262.	0.8	110
16	Dynamin-related protein Drp1 is required for Bax translocation to mitochondria in response to irradiation-induced apoptosis. <i>Oncotarget</i> , 2015, 6, 22598-22612.	0.8	74
17	Rituximab-induced HMGB1 release is associated with inhibition of STAT3 activity in human diffuse large B-cell lymphoma. <i>Oncotarget</i> , 2015, 6, 27816-27831.	0.8	20
18	Overexpression of HMGB1 Receptor RAGE Is Associated with Worse Clinical Outcome in Patients with Chronic Lymphocytic Leukemia. <i>Blood</i> , 2015, 126, 617-617.	0.6	0

#	ARTICLE	IF	CITATIONS
19	Extracellular HMGB1 promotes differentiation of nurse-like cells in chronic lymphocytic leukemia. <i>Blood</i> , 2014, 123, 1709-1719.	0.6	95
20	Stem cell factor is a novel independent prognostic biomarker for hepatocellular carcinoma after curative resection. <i>Carcinogenesis</i> , 2014, 35, 2283-2290.	1.3	23
21	Dangerous power: mitochondria in CLL cells. <i>Blood</i> , 2014, 123, 2596-2597.	0.6	5
22	Dysregulation of autophagy in human follicular lymphoma is independent of overexpression of BCL-2. <i>Oncotarget</i> , 2014, 5, 11653-11668.	0.8	22
23	Why bortezomib cannot go with 'green'?. <i>Cancer Biology and Medicine</i> , 2013, 10, 206-13.	1.4	19
24	Blocking Autophagy Prevents Bortezomib-Induced NF- $\kappa$ B Activation by Reducing I- $\kappa$ B $\alpha$ Degradation in Lymphoma Cells. <i>PLoS ONE</i> , 2012, 7, e32584.	1.1	87
25	Methylseleninic acid antagonizes the cytotoxic effect of bortezomib in mantle cell lymphoma cell lines through modulation of Bcl-2 family proteins. <i>British Journal of Haematology</i> , 2012, 156, 286-289.	1.2	3
26	HMGB1 Activates TLR9/RAGE Signalling Pathway and Sustains Chronic Lymphocytic Leukemic Cell in Vitro Survival. <i>Blood</i> , 2012, 120, 3860-3860.	0.6	1
27	Activation of Mitochondrial STAT3 Increases Mitochondrial Respiration and Inhibits Oxidative Stress in Chronic Lymphocytic Leukemic Cells. <i>Blood</i> , 2011, 118, 287-287.	0.6	2
28	CD160 signaling mediates PI3K-dependent survival and growth signals in chronic lymphocytic leukemia. <i>Blood</i> , 2010, 115, 3079-3088.	0.6	48
29	The alpha-5 helix of Bax is sensitive to ubiquitin-dependent degradation. <i>Biochemical and Biophysical Research Communications</i> , 2008, 371, 10-15.	1.0	6
30	Bortezomib blocks Bax degradation in malignant B cells during treatment with TRAIL. <i>Blood</i> , 2008, 111, 2797-2805.	0.6	79
31	Increased proteasomal degradation of Bax is a common feature of poor prognosis chronic lymphocytic leukemia. <i>Blood</i> , 2008, 111, 2790-2796.	0.6	28
32	Dietary flavonoids inhibit the anticancer effects of the proteasome inhibitor bortezomib. <i>Blood</i> , 2008, 112, 3835-3846.	0.6	83
33	Bcl-2 Inhibitors Sensitize Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Induced Apoptosis by Uncoupling of Mitochondrial Respiration in Human Leukemic CEM Cells. <i>Cancer Research</i> , 2004, 64, 3607-3616.	0.4	79
34	Increase in the ratio of mitochondrial Bax/Bcl-XL induces Bax activation in human leukemic K562 cell line. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2004, 9, 377-384.	2.2	28
35	BH3-domain mimetic compound BH3I-2 induces rapid damage to the inner mitochondrial membrane prior to the cytochrome c release from mitochondria. <i>British Journal of Haematology</i> , 2003, 121, 332-340.	1.2	25
36	Role of DNA methylation in the suppression of Apaf-1 protein in human leukaemia. <i>Oncogene</i> , 2003, 22, 451-455.	2.6	87

#	ARTICLE	IF	CITATIONS
37	Role of Smac in human leukaemic cell apoptosis and proliferation. <i>Oncogene</i> , 2003, 22, 1589-1599.	2.6	54
38	Bax conformational change is a crucial step for PUMA-mediated apoptosis in human leukemia. <i>Biochemical and Biophysical Research Communications</i> , 2003, 310, 956-962.	1.0	67
39	Liposomal encapsulation diminishes daunorubicin-induced generation of reactive oxygen species, depletion of ATP and necrotic cell death in human leukaemic cells. <i>British Journal of Haematology</i> , 2002, 117, 333-342.	1.2	28
40	Apaf-1XL Is an Inactive Isoform Compared with Apaf-1L. <i>Biochemical and Biophysical Research Communications</i> , 2001, 282, 268-272.	1.0	10
41	TRAIL-Induced Apoptosis in Type I Leukemic Cells Is Not Enhanced by Overexpression of Bax. <i>Biochemical and Biophysical Research Communications</i> , 2001, 283, 1037-1045.	1.0	21
42	c-IAP1 Blocks TNF $\alpha$ -Mediated Cytotoxicity Upstream of Caspase-Dependent and -Independent Mitochondrial Events in Human Leukemic Cells. <i>Biochemical and Biophysical Research Communications</i> , 2001, 287, 181-189.	1.0	18
43	Apaf-1 protein deficiency confers resistance to cytochrome c-dependent apoptosis in human leukemic cells. <i>Blood</i> , 2001, 98, 414-421.	0.6	136
44	Generation of reactive oxygen species is not involved in idarubicin-induced apoptosis in human leukaemic cells. <i>British Journal of Haematology</i> , 2001, 115, 817-825.	1.2	7
45	Quantitative determination of apoptosis on leukemia cells by infrared spectroscopy. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2001, 6, 269-278.	2.2	85
46	Bax translocation is crucial for the sensitivity of leukaemic cells to etoposide-induced apoptosis. <i>Oncogene</i> , 2001, 20, 4817-4826.	2.6	73
47	8-Cl-adenosine mediated cytotoxicity and sensitization of T-lymphoblastic leukemia cells to TNF $\alpha$ -induced apoptosis is via inactivation of NF- $\kappa$ B. <i>Leukemia Research</i> , 2001, 25, 423-431.	0.4	15
48	Constitutive levels of cAMP-dependent protein kinase activity determine sensitivity of human multidrug-resistant leukaemic cell lines to growth inhibition and apoptosis by forskolin and tumour necrosis factor alpha. <i>British Journal of Haematology</i> , 2000, 108, 565-573.	1.2	14
49	Subcellular Distribution and Redistribution of Bcl-2 Family Proteins in Human Leukemia Cells Undergoing Apoptosis. <i>Blood</i> , 1999, 93, 2353-2359.	0.6	80
50	Pgp-positive leukaemic cells have increased mtDNA but no increased rate of proliferation. <i>British Journal of Haematology</i> , 1999, 107, 861-869.	1.2	13
51	Subcellular Distribution and Redistribution of Bcl-2 Family Proteins in Human Leukemia Cells Undergoing Apoptosis. <i>Blood</i> , 1999, 93, 2353-2359.	0.6	6
52	Mitochondrial ultracondensation, but not swelling, is involved in TNF $\alpha$ -induced apoptosis in human T-lymphoblastic leukaemic cells. <i>Leukemia Research</i> , 1997, 21, 973-983.	0.4	34
53	Inhibition of autophagy abrogates tumour necrosis factor $\alpha$ induced apoptosis in human T-lymphoblastic leukaemic cells. <i>British Journal of Haematology</i> , 1997, 98, 673-685.	1.2	221
54	Mitochondrial electron transport chain activity, but not ATP synthesis, is required for drug-induced apoptosis in human leukaemic cells: a possible novel mechanism of regulating drug resistance. <i>British Journal of Haematology</i> , 1997, 98, 686-698.	1.2	42

#	ARTICLE	IF	CITATIONS
55	Modulation of surface TNF expression by human leukaemic cells alters their sensitivity to exogenous TNF. <i>Leukemia Research</i> , 1996, 20, 47-55.	0.4	13
56	TNF-mediated killing of human leukaemic cells: Effects of endogenous antioxidant levels and TNF $\alpha$ expression in leukaemic cell lines. <i>Leukemia Research</i> , 1995, 19, 187-194.	0.4	10