

Kylie Hill

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5306859/kylie-hill-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

134
papers

4,710
citations

30
h-index

67
g-index

142
ext. papers

5,785
ext. citations

3.4
avg, IF

5.17
L-index

#	Paper	IF	Citations
134	Patterns of Change in Device-Based Physical Activity and Sedentary Time Following Bariatric Surgery: a Longitudinal Observational Study. <i>Obesity Surgery</i> , 2021 , 31, 3015-3025	3.7	0
133	In People With COPD, There Is Limited Evidence That Exercise Training Reduces Sedentary Time, and Behavior Change Techniques Are Poorly Reported: Systematic Review and Meta-Analysis. <i>Physical Therapy</i> , 2021 , 101,	3.3	4
132	Endurance cycle ergometry tests performed at a sub-maximal work rate elicit peak physiological and symptom responses in adults with cystic fibrosis. <i>Internal Medicine Journal</i> , 2021 , 51, 1168-1172	1.6	
131	Use of supplemental oxygen during exercise testing and training for people with chronic obstructive pulmonary disease: a survey of Australian pulmonary rehabilitation programs. <i>Brazilian Journal of Physical Therapy</i> , 2021 , 25, 97-102	3.7	2
130	Exercise & Sports Science Australia (ESSA) position statement on exercise and chronic obstructive pulmonary disease. <i>Journal of Science and Medicine in Sport</i> , 2021 , 24, 52-59	4.4	5
129	Berg Balance Scale Score as a Predictor of Independent Walking at Discharge among Adult Stroke Survivors. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2021 , 73, 252-256	0.8	
128	A Behavior Change Intervention Aimed at Increasing Physical Activity Improves Clinical Control in Adults With Asthma: A Randomized Controlled Trial. <i>Chest</i> , 2021 , 159, 46-57	5.3	9
127	Aquatic high intensity interval training to improve aerobic capacity is feasible in adolescents with cerebral palsy: pilot randomised controlled trial. <i>Clinical Rehabilitation</i> , 2021 , 35, 222-231	3.3	2
126	Reducing sedentary behavior in individuals with COPD: healthcare professionals' perspectives. <i>Physiotherapy Theory and Practice</i> , 2021 , 1-12	1.5	
125	Feasibility, tolerance and effects of adding impact loading exercise to pulmonary rehabilitation in people with chronic obstructive pulmonary disease: study protocol for a pilot randomised controlled trial. <i>Pilot and Feasibility Studies</i> , 2021 , 7, 151	1.9	
124	Exercise prescription for people with stable COPD as part of a pulmonary rehabilitation programme 2021 , 53-66		1
123	Initiating exercise training early during the hospitalisation for an exacerbation of chronic obstructive pulmonary disease improves exercise capacity and quadriceps strength: A randomised controlled trial. <i>Respiratory Medicine: X</i> , 2020 , 2, 100024	1.6	0
122	Modified Chester Step Test in a Healthy Adult Population: Measurement Properties and Development of a Regression Equation to Estimate Test Duration. <i>Physical Therapy</i> , 2020 , 100, 1411-1418	3.3	1
121	Determinants of Sedentary Behaviour in Individuals with COPD: A Qualitative Exploration Guided by the Theoretical Domains Framework. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2020 , 17, 65-73	2	7
120	Effect of Using a Wheeled Walker on Physical Activity and Sedentary Time in People with Chronic Obstructive Pulmonary Disease: A Randomised Cross-Over Trial. <i>Lung</i> , 2020 , 198, 213-219	2.9	1
119	High-Intensity Interval Training Is Effective at Increasing Exercise Endurance Capacity and Is Well Tolerated by Adults with Cystic Fibrosis. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	3
118	OPTimising the implementation of pulMonyary rehAbiLitation in people with chronic obstructive pulmonary disease (the OPTIMAL study): mixed methods study protocol. <i>BMC Pulmonary Medicine</i> , 2020 , 20, 286	3.5	0

117	Survivors of Acute Lung Injury Have Greater Impairments in Strength and Exercise Capacity Than Survivors of Other Critical Illnesses as Measured Shortly After ICU Discharge. <i>Journal of Intensive Care Medicine</i> , 2020 , 885066620981899	3.3	1
116	Reduced Step Count and Clinical Frailty in Hospitalized Adults With Community-Acquired Pneumonia. <i>Respiratory Care</i> , 2020 , 65, 455-463	2.1	4
115	A smartphone application for reporting symptoms in adults with cystic fibrosis improves the detection of exacerbations: Results of a randomised controlled trial. <i>Journal of Cystic Fibrosis</i> , 2020 , 19, 271-276	4.1	12
114	Exercise training for adults hospitalized with an acute respiratory condition: a systematic scoping review. <i>Clinical Rehabilitation</i> , 2020 , 34, 45-55	3.3	6
113	Exercise testing and exercise training within cystic fibrosis centres across Australia and New Zealand: what is considered important and what is current practice?. <i>Internal Medicine Journal</i> , 2020 , 50, 1091-1099	1.6	3
112	Effects of high intensity interval training on exercise capacity in people with chronic pulmonary conditions: a narrative review. <i>BMC Sports Science, Medicine and Rehabilitation</i> , 2020 , 12, 22	2.4	15
111	Effects of a behaviour change intervention aimed at increasing physical activity on clinical control of adults with asthma: study protocol for a randomised controlled trial. <i>BMC Sports Science, Medicine and Rehabilitation</i> , 2019 , 11, 16	2.4	10
110	Reliability and Validity of the Four-Point Pusher Score: An Assessment Tool for Measuring Lateropulsion and Pusher Behaviour in Adults after Stroke. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2019 , 71, 34-42	0.8	6
109	Higher Levels of Education Are Associated With Full-Time Work in Adults With Cystic Fibrosis. <i>Respiratory Care</i> , 2019 , 64, 1116-1122	2.1	1
108	Effects of Ongoing Feedback During a 12-Month Maintenance Walking Program on Daily Physical Activity in People with COPD. <i>Lung</i> , 2019 , 197, 315-319	2.9	3
107	Oxygen compared to air during exercise training in COPD with exercise-induced desaturation. <i>European Respiratory Journal</i> , 2019 , 53,	13.6	24
106	The effect of aquatic high-intensity interval training on aerobic performance, strength and body composition in a non-athletic population: systematic review and meta-analysis. <i>Clinical Rehabilitation</i> , 2019 , 33, 157-170	3.3	8
105	Exercise training undertaken by people within 12 months of lung resection for non-small cell lung cancer. <i>The Cochrane Library</i> , 2019 , 6, CD009955	5.2	25
104	Exercise training in COPD with exercise-induced desaturation does improve exercise capacity, irrespective of whether supplemental oxygen or air is provided during training. <i>European Respiratory Journal</i> , 2019 , 54,	13.6	2
103	People With COPD Who Respond to Ground-Based Walking Training Are Characterized by Lower Pre-training Exercise Capacity and Better Lung Function and Have Greater Progression in Walking Training Distance. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2019 , 39, 338-343	3.6	3
102	Abnormal Exercise Responses in Survivors of Acute Lung Injury During Cardiopulmonary Exercise Testing: AN OBSERVATIONAL STUDY. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2019 , 39, E16-E22	3.6	1
101	The minimal detectable difference for endurance shuttle walk test performance in people with COPD on completion of a program of high-intensity ground-based walking. <i>Respiratory Medicine</i> , 2019 , 146, 18-22	4.6	0
100	Critically appraised paper: A 12-week pedometer-based intervention, delivered in primary care, produces long-term gains in physical activity [synopsis]. <i>Journal of Physiotherapy</i> , 2019 , 65, 54	2.9	

99	Behaviour change techniques to optimise participation in physical activity or exercise in adolescents and young adults with chronic cardiorespiratory conditions: a systematic review. <i>Internal Medicine Journal</i> , 2019 , 49, 1209-1220	1.6	8
98	Regression equations to estimate the 2-min walk distance in an adult Asian population aged 40-75 years. <i>Respirology</i> , 2018 , 23, 674-680	3.6	4
97	A smartphone application for reporting symptoms in adults with cystic fibrosis: protocol of a randomised controlled trial. <i>BMJ Open</i> , 2018 , 8, e021136	3	6
96	Critically Appraised Papers: In people with chronic obstructive pulmonary disease, home-based pulmonary rehabilitation produces similar results to a hospital-based outpatient program [synopsis]. <i>Journal of Physiotherapy</i> , 2018 , 64, 122	2.9	1
95	Shuttle walk tests in people with COPD who demonstrate exercise-induced oxygen desaturation: An analysis of test repeatability and cardiorespiratory responses. <i>Chronic Respiratory Disease</i> , 2018 , 15, 131-137	3	2
94	Do factors related to participation in physical activity change following restrictive bariatric surgery? A qualitative study. <i>Obesity Research and Clinical Practice</i> , 2018 , 12, 307-316	5.4	18
93	Effect on health-related quality of life of ongoing feedback during a 12-month maintenance walking programme in patients with COPD: a randomized controlled trial. <i>Respirology</i> , 2018 , 23, 60-67	3.6	10
92	High usability of a smartphone application for reporting symptoms in adults with cystic fibrosis. <i>Journal of Telemedicine and Telecare</i> , 2018 , 24, 547-552	6.8	6
91	Walking-based activity and sedentary behavior in Saudi males with chronic obstructive pulmonary disease. <i>Journal of King Abdulaziz University, Islamic Economics</i> , 2018 , 39, 506-513	1.1	3
90	Effects of high intensity interval training on exercise capacity in people with cystic fibrosis: study protocol for a randomised controlled trial. <i>BMC Sports Science, Medicine and Rehabilitation</i> , 2018 , 10, 19	2.4	2
89	Neuromuscular electrostimulation for adults with chronic obstructive pulmonary disease. <i>The Cochrane Library</i> , 2018 , 5, CD010821	5.2	11
88	The impact of cystic fibrosis on work attendance and performance in adults living in rural and remote Western Australia. <i>Journal of Cystic Fibrosis</i> , 2017 , 16, e1-e2	4.1	2
87	Exercise training for people following curative intent treatment for non-small cell lung cancer: a randomized controlled trial. <i>Brazilian Journal of Physical Therapy</i> , 2017 , 21, 58-68	3.7	24
86	Analysis of nocturnal actigraphic sleep measures in patients with COPD and their association with daytime physical activity. <i>Thorax</i> , 2017 , 72, 694-701	7.3	32
85	Quantification of walking-based physical activity and sedentary time in individuals with Rett syndrome. <i>Developmental Medicine and Child Neurology</i> , 2017 , 59, 605-611	3.3	21
84	Approaches and adjuncts used by physiotherapists when suctioning adult patients who are intubated and ventilated in intensive care units in Australia and New Zealand: A cross-sectional survey. <i>Australian Critical Care</i> , 2017 , 30, 307-313	2.9	5
83	Telehealth clinics increase access to care for adults with cystic fibrosis living in rural and remote Western Australia. <i>Journal of Telemedicine and Telecare</i> , 2017 , 23, 673-679	6.8	35
82	Effects of ground-based walking training on daily physical activity in people with COPD: A randomised controlled trial. <i>Respiratory Medicine</i> , 2017 , 132, 139-145	4.6	16

81	Measurement of Sedentary Behaviors or "Downtime" in Rett Syndrome. <i>Journal of Child Neurology</i> , 2017 , 32, 1009-1013	2.5	6
80	Should resistance training be targeted to a specific subgroup of patients with non-small cell lung cancer?. <i>Respirology</i> , 2017 , 22, 1473	3.6	0
79	Physical activity patterns and clusters in 1001 patients with COPD. <i>Chronic Respiratory Disease</i> , 2017 , 14, 256-269	3	36
78	Beliefs, Barriers and Facilitators to Physical Activity in Bariatric Surgery Candidates. <i>Obesity Surgery</i> , 2016 , 26, 1097-109	3.7	35
77	Patterns of sedentary behaviour and physical activity in people following curative intent treatment for non-small cell lung cancer. <i>Chronic Respiratory Disease</i> , 2016 , 13, 82-5	3	20
76	Comparison of the six-minute walk test with a cycle-based cardiopulmonary exercise test in people following curative intent treatment for non-small cell lung cancer. <i>Chronic Respiratory Disease</i> , 2016 , 13, 118-27	3	7
75	Supervised walking training improves health-related quality of life and exercise endurance in people with chronic obstructive pulmonary disease [synopsis]. <i>Journal of Physiotherapy</i> , 2016 , 62, 50	2.9	1
74	Letter to the Editor: "Risk factors for postoperative pneumonia after lung cancer surgery and impact of pneumonia on survival". <i>Respiratory Medicine</i> , 2016 , 112, 132	4.6	
73	Pulmonary Rehabilitation Exercise Prescription in Chronic Obstructive Pulmonary Disease: Review of Selected Guidelines: AN OFFICIAL STATEMENT FROM THE AMERICAN ASSOCIATION OF CARDIOVASCULAR AND PULMONARY REHABILITATION. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2016 , 36, 75-83	3.6	92
72	A randomised controlled trial of supplemental oxygen versus medical air during exercise training in people with chronic obstructive pulmonary disease: supplemental oxygen in pulmonary rehabilitation trial (SuppORT) (Protocol). <i>BMC Pulmonary Medicine</i> , 2016 , 16, 25	3.5	13
71	Early goal-directed mobilisation in the intensive care unit is feasible and safe, and increases both the level and duration of activity [synopsis]. <i>Journal of Physiotherapy</i> , 2016 , 62, 225	2.9	
70	Changing physical activity and sedentary behaviour in people with COPD. <i>Respirology</i> , 2016 , 21, 419-26	3.6	41
69	Discordance Between Distance Ambulated as Part of Usual Care and Functional Exercise Capacity in Survivors of Critical Illness Upon Intensive Care Discharge: Observational Study. <i>Physical Therapy</i> , 2015 , 95, 1254-63	3.3	6
68	Rett syndrome: establishing a novel outcome measure for walking activity in an era of clinical trials for rare disorders. <i>Disability and Rehabilitation</i> , 2015 , 37, 1992-6	2.4	21
67	Impairments after curative intent treatment for non-small cell lung cancer: a comparison with age and gender-matched healthy controls. <i>Respiratory Medicine</i> , 2015 , 109, 1332-9	4.6	23
66	Physical activity and sedentary behaviour: applying lessons to chronic obstructive pulmonary disease. <i>Internal Medicine Journal</i> , 2015 , 45, 474-82	1.6	64
65	A validation study of a modified Bouchard activity record that extends the concept of Riptime to Rett syndrome. <i>Developmental Medicine and Child Neurology</i> , 2015 , 57, 1137-42	3.3	9
64	Is there an association between symptoms of anxiety and depression and quality of life in patients with chronic obstructive pulmonary disease?. <i>Canadian Respiratory Journal</i> , 2015 , 22, 37-41	2.1	8

63	An Official American Thoracic Society/European Respiratory Society Policy Statement: Enhancing Implementation, Use, and Delivery of Pulmonary Rehabilitation. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2015 , 192, 1373-86	10.2	350
62	The use of positive expiratory pressure therapy does not appear to be effective in people hospitalised with an acute exacerbation of chronic obstructive pulmonary disease (AECOPD). <i>Journal of Physiotherapy</i> , 2015 , 61, 43	2.9	3
61	Reliability and validity of an arabic version of the dyspnea-12 questionnaire for Saudi nationals with chronic obstructive pulmonary disease. <i>Annals of Thoracic Medicine</i> , 2015 , 10, 112-7	2.2	16
60	Patterns of physical activity and sedentary behavior after bariatric surgery: an observational study. <i>Surgery for Obesity and Related Diseases</i> , 2014 , 10, 524-30	3	20
59	Ground-based walking training improves quality of life and exercise capacity in COPD. <i>European Respiratory Journal</i> , 2014 , 44, 885-94	13.6	45
58	Strategies to enhance the benefits of exercise training in the respiratory patient. <i>Clinics in Chest Medicine</i> , 2014 , 35, 323-36	5.3	12
57	Balance-specific training embedded within a pulmonary rehabilitation program may reduce falls risk in people with COPD. <i>Journal of Physiotherapy</i> , 2014 , 60, 111	2.9	
56	Comorbidities and medication burden in patients with chronic obstructive pulmonary disease attending pulmonary rehabilitation. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2014 , 34, 75-9	3.6	10
55	Estimating endurance shuttle walk test speed using the six-minute walk test in people with chronic obstructive pulmonary disease. <i>Chronic Respiratory Disease</i> , 2014 , 11, 89-94	3	3
54	Exercise training for people following lung resection for non-small cell lung cancer - a Cochrane systematic review. <i>Cancer Treatment Reviews</i> , 2014 , 40, 585-94	14.4	58
53	Exercise training undertaken by people within 12 months of lung resection for non-small cell lung cancer. <i>The Cochrane Library</i> , 2013 , CD009955	5.2	32
52	An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2013 , 188, e13-64	10.2	1863
51	Relationships between mortality, morbidity, and physical function in adults who survived a period of prolonged mechanical ventilation. <i>Journal of Critical Care</i> , 2013 , 28, 427-32	4	8
50	Comparing finger and forehead sensors to measure oxygen saturation in people with chronic obstructive pulmonary disease. <i>Respirology</i> , 2013 , 18, 1143-7	3.6	5
49	The use of regression equations to estimate peak work rate in people with COPD. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2013 , 10, 118-9	2	1
48	Xbox Kinect represents high intensity exercise for adults with cystic fibrosis. <i>Journal of Cystic Fibrosis</i> , 2013 , 12, 604-8	4.1	21
47	Physiotherapy practice patterns for patients undergoing surgery for lung cancer: a survey of hospitals in Australia and New Zealand. <i>Internal Medicine Journal</i> , 2013 , 43, 394-401	1.6	23
46	The importance of components of pulmonary rehabilitation, other than exercise training, in COPD. <i>European Respiratory Review</i> , 2013 , 22, 405-13	9.8	42

45	Relationship and responsiveness of three upper-limb tests in patients with chronic obstructive pulmonary disease. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2013 , 65, 40-3	0.8	24
44	Arm elevation and coordinated breathing strategies in patients with COPD. <i>Chest</i> , 2013 , 144, 128-135	5.3	11
43	Response. <i>Chest</i> , 2013 , 143, 281	5.3	
42	Neuromuscular electrostimulation for chronic obstructive pulmonary disease. <i>The Cochrane Library</i> , 2013 ,	5.2	1
41	International COPD Coalition Column: pulmonary rehabilitation-reaching out to our international community. <i>Journal of Thoracic Disease</i> , 2013 , 5, 343-8	2.6	2
40	Comparing peak and submaximal cardiorespiratory responses during field walking tests with incremental cycle ergometry in COPD. <i>Respirology</i> , 2012 , 17, 278-84	3.6	53
39	Pulmonary rehabilitation: a review of the recent literature. <i>Chest</i> , 2012 , 142, 738-749	5.3	34
38	Accuracy and responsiveness of the stepwatch activity monitor and ActivPAL in patients with COPD when walking with and without a rollator. <i>Disability and Rehabilitation</i> , 2012 , 34, 1317-22	2.4	44
37	Initial assessment of the StepWatch Activity Monitor to measure walking activity in Rett syndrome. <i>Disability and Rehabilitation</i> , 2012 , 34, 1010-5	2.4	14
36	Rollator use does not consistently change the metabolic cost of walking in people with chronic obstructive pulmonary disease. <i>Archives of Physical Medicine and Rehabilitation</i> , 2012 , 93, 1077-80	2.8	10
35	Does exercise training change physical activity in people with COPD? A systematic review and meta-analysis. <i>Chronic Respiratory Disease</i> , 2012 , 9, 17-26	3	150
34	Exploring the capacity to ambulate after a period of prolonged mechanical ventilation. <i>Journal of Critical Care</i> , 2012 , 27, 542-8	4	4
33	A simple method to derive speed for the endurance shuttle walk test. <i>Respiratory Medicine</i> , 2012 , 106, 1665-70	4.6	17
32	Learning from the Learning Effect in the Six-Minute-Walk Test. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2012 , 185, 684-685	10.2	1
31	Has My Patient Responded?. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2012 , 185, 895-896.2		
30	Maximum voluntary ventilation is more strongly associated with energy expenditure during simple activities of daily living than measures of airflow obstruction or respiratory muscle strength in patients with COPD. <i>Chronic Respiratory Disease</i> , 2012 , 9, 239-40	3	2
29	Defining the relationship between average daily energy expenditure and field-based walking tests and aerobic reserve in COPD. <i>Chest</i> , 2012 , 141, 406-412	5.3	17
28	The effect of pulmonary rehabilitation on critical walk speed in patients with COPD: a comparison with self-paced walks. <i>Chest</i> , 2012 , 141, 413-419	5.3	12

27	The 6-min walk test: responses in healthy Canadians aged 45 to 85 years. <i>Applied Physiology, Nutrition and Metabolism</i> , 2011 , 36, 643-9	3	43
26	The incidence of falls in intensive care survivors. <i>Australian Critical Care</i> , 2011 , 24, 167-74	2.9	14
25	Effects of loading on upper airway and respiratory pump muscle motoneurons. <i>Respiratory Physiology and Neurobiology</i> , 2011 , 179, 64-70	2.8	4
24	Has my patient responded? Interpreting clinical measurements such as the 6-minute-walk test. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2011 , 184, 642-6	10.2	48
23	Effects of high-intensity inspiratory muscle training following a near-fatal gunshot wound. <i>Physical Therapy</i> , 2011 , 91, 1377-84	3.3	
22	Resistance arm training in patients with COPD: A Randomized Controlled Trial. <i>Chest</i> , 2011 , 139, 151-8	5.3	66
21	Properties of self-paced walking in chronic respiratory disease: a patient goal-oriented assessment. <i>Chest</i> , 2011 , 140, 737-743	5.3	11
20	Identifying adults at risk of COPD who need confirmatory spirometry in primary care: Do symptom-based questions help?. <i>Canadian Family Physician</i> , 2011 , 57, e51-7	0.9	9
19	State of the art: how to set up a pulmonary rehabilitation program. <i>Respirology</i> , 2010 , 15, 1157-73	3.6	63
18	Measurement properties of the SenseWear armband in adults with chronic obstructive pulmonary disease. <i>Thorax</i> , 2010 , 65, 486-91	7.3	99
17	Effect of airway clearance techniques in patients experiencing an acute exacerbation of chronic obstructive pulmonary disease: a systematic review. <i>Chronic Respiratory Disease</i> , 2010 , 7, 9-17	3	31
16	Interval versus continuous training in individuals with chronic obstructive pulmonary disease--a systematic review. <i>Thorax</i> , 2010 , 65, 157-64	7.3	123
15	Prevalence and underdiagnosis of chronic obstructive pulmonary disease among patients at risk in primary care. <i>Cmaj</i> , 2010 , 182, 673-8	3.5	167
14	Inspiratory muscle training for patients with chronic obstructive pulmonary disease: a practical guide for clinicians. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010 , 91, 1466-70	2.8	59
13	Characteristics of people with chronic lung disease who rest during the six-minute walk test. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010 , 91, 1765-9	2.8	3
12	A description of weekend physiotherapy services in three tertiary hospitals in the greater Toronto area. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2010 , 62, 155-62	0.8	8
11	The provision of weekend physiotherapy services in tertiary-care hospitals in Canada. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2010 , 62, 347-54	0.8	26
10	Disease-specific education in the primary care setting increases the knowledge of people with chronic obstructive pulmonary disease: a randomized controlled trial. <i>Patient Education and Counseling</i> , 2010 , 81, 14-8	3.1	39

9	Workshops to disseminate the Canadian Thoracic Society guidelines for chronic obstructive pulmonary disease to health care professionals in Ontario: impact on knowledge, perceived health care practices and participant satisfaction. <i>Canadian Respiratory Journal</i> , 2009 , 16, 81-5	2.1	2
8	Physical therapy practice patterns in acute exacerbations of chronic obstructive pulmonary disease. <i>Canadian Respiratory Journal</i> , 2009 , 16, 86-92	2.1	20
7	Arm exercise training in patients with chronic obstructive pulmonary disease: a systematic review. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2009 , 29, 277-83	3.6	46
6	Daily utility and satisfaction with rollators among persons with chronic obstructive pulmonary disease. <i>Archives of Physical Medicine and Rehabilitation</i> , 2008 , 89, 1108-13	2.8	29
5	Estimating maximum work rate during incremental cycle ergometry testing from six-minute walk distance in patients with chronic obstructive pulmonary disease. <i>Archives of Physical Medicine and Rehabilitation</i> , 2008 , 89, 1782-7	2.8	53
4	Modifying track layout from straight to circular has a modest effect on the 6-min walk distance. <i>Chest</i> , 2008 , 133, 1155-60	5.3	35
3	Repeat pulmonary rehabilitation programs confer similar increases in functional exercise capacity to initial programs. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2008 , 28, 410-4	3.6	16
2	Limited functional performance in chronic obstructive pulmonary disease: nature, causes and measurement. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2007 , 4, 257-61	2	9
1	Dyspnoea in COPD: can inspiratory muscle training help?. <i>Australian Journal of Physiotherapy</i> , 2004 , 50, 169-80		30