

# Arja Jukkola-Vuorinen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5318613/publications.pdf>

Version: 2024-02-01

56  
papers

4,947  
citations

236925

25  
h-index

168389

53  
g-index

58  
all docs

58  
docs citations

58  
times ranked

8892  
citing authors

#	ARTICLE	IF	CITATIONS
1	How breast cancer recurrences are found – a real-world, prospective cohort study. <i>Acta Oncologica</i> , 2022, 61, 417-424.	1.8	0
2	Adjuvant Capecitabine for Early Breast Cancer: 15-Year Overall Survival Results From a Randomized Trial. <i>Journal of Clinical Oncology</i> , 2022, , JCO2102054.	1.6	14
3	Association of Metformin, Other Antidiabetic Medications, and Statins With Incidence of Colon Cancer in Patients With Type 2 Diabetes. <i>Clinical Colorectal Cancer</i> , 2021, 20, e113-e119.	2.3	5
4	Real-world, single-centre prospective data of age at breast cancer onset: focus on survival and reproductive history. <i>BMJ Open</i> , 2021, 11, e041706.	1.9	2
5	Early-Life Risk Factors for Breast Cancer – Prospective Follow-up in the Northern Finland Birth Cohort 1966. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021, 30, 616-622.	2.5	0
6	CD73 facilitates EMT progression and promotes lung metastases in triple-negative breast cancer. <i>Scientific Reports</i> , 2021, 11, 6035.	3.3	42
7	Association of antidiabetic medication and statins with survival from ductal and lobular breast carcinoma in women with type 2 diabetes. <i>Scientific Reports</i> , 2021, 11, 10445.	3.3	5
8	Radiological and pathological assessment of response to neoadjuvant CDK4/6 inhibitor and endocrine treatments in a real-life setting – initial results. <i>Acta Radiologica Open</i> , 2021, 10, 205846012110306.	0.6	1
9	Fine-mapping of 150 breast cancer risk regions identifies 191 likely target genes. <i>Nature Genetics</i> , 2020, 52, 56-73.	21.4	120
10	Prognostic factors in metastatic breast cancer: a prospective single-centre cohort study in a Finnish University Hospital. <i>BMJ Open</i> , 2020, 10, e038798.	1.9	12
11	High Parity Predicts Poor Outcomes in Patients With Luminal B-Like (HER2 Negative) Early Breast Cancer: A Prospective Finnish Single-Center Study. <i>Frontiers in Oncology</i> , 2020, 10, 1470.	2.8	10
12	Early progression of breast cancer during neoadjuvant chemotherapy may predict poorer prognoses. <i>Acta Oncologica</i> , 2020, 59, 1036-1042.	1.8	2
13	Survival after breast cancer in women with type 2 diabetes using antidiabetic medication and statins: a retrospective cohort study. <i>Acta Oncologica</i> , 2020, 59, 1110-1117.	1.8	12
14	Elevated preoperative serum levels of collagen I carboxyterminal telopeptide predict better outcome in early-stage luminal-B-like (HER2-negative) and triple-negative subtypes of breast cancer. <i>Tumor Biology</i> , 2019, 41, 101042831984708.	1.8	5
15	High baseline Tie1 level predicts poor survival in metastatic breast cancer. <i>BMC Cancer</i> , 2019, 19, 732.	2.6	12
16	Cytoplasmic Mineralocorticoid Receptor Expression Predicts Dismal Local Relapse-free Survival in Non-triple-negative Breast Cancer. <i>Anticancer Research</i> , 2019, 39, 5879-5890.	1.1	6
17	Low Plasma IL-8 Levels During Chemotherapy Are Predictive of Excellent Long-Term Survival in Metastatic Breast Cancer. <i>Clinical Breast Cancer</i> , 2019, 19, e522-e533.	2.4	27
18	Association of antidiabetic medication and statins with breast cancer incidence in women with type 2 diabetes. <i>Breast Cancer Research and Treatment</i> , 2019, 175, 741-748.	2.5	9

#	ARTICLE	IF	CITATIONS
19	Genome-wide association study of germline variants and breast cancer-specific mortality. <i>British Journal of Cancer</i> , 2019, 120, 647-657.	6.4	52
20	Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer Subtypes. <i>American Journal of Human Genetics</i> , 2019, 104, 21-34.	6.2	711
21	Neuroendocrine Breast Carcinomas Share Prognostic Factors with Gastroenteropancreatic Neuroendocrine Tumors: A Putative Prognostic Role of Menin, p27, and SSTR-2A. <i>Oncology</i> , 2019, 96, 147-155.	1.9	7
22	Rare missense mutations in <i>RECQL</i> and <i>POLG</i> associate with inherited predisposition to breast cancer. <i>International Journal of Cancer</i> , 2018, 142, 2286-2292.	5.1	15
23	High-level cytoplasmic claudin 3 expression is an independent predictor of poor survival in triple-negative breast cancer. <i>BMC Cancer</i> , 2018, 18, 223.	2.6	25
24	USP28 Deficiency Promotes Breast and Liver Carcinogenesis as well as Tumor Angiogenesis in a HIF-independent Manner. <i>Molecular Cancer Research</i> , 2018, 16, 1000-1012.	3.4	23
25	Long-term cardiovascular morbidity and mortality in patients treated for differentiated thyroid cancer. <i>Clinical Endocrinology</i> , 2018, 88, 303-310.	2.4	47
26	Primary neuroendocrine breast carcinomas are associated with poor local control despite favourable biological profile: a retrospective clinical study. <i>BMC Cancer</i> , 2017, 17, 72.	2.6	27
27	Adjuvant Capecitabine in Combination With Docetaxel, Epirubicin, and Cyclophosphamide for Early Breast Cancer. <i>JAMA Oncology</i> , 2017, 3, 793.	7.1	74
28	Case-control analysis of truncating mutations in DNA damage response genes connects TEX15 and FANCD2 with hereditary breast cancer susceptibility. <i>Scientific Reports</i> , 2017, 7, 681.	3.3	20
29	Association analysis identifies 65 new breast cancer risk loci. <i>Nature</i> , 2017, 551, 92-94.	27.8	1,099
30	Identification of ten variants associated with risk of estrogen-receptor-negative breast cancer. <i>Nature Genetics</i> , 2017, 49, 1767-1778.	21.4	289
31	Body mass index and breast cancer survival: a Mendelian randomization analysis. <i>International Journal of Epidemiology</i> , 2017, 46, 1814-1822.	1.9	45
32	CHEK2 c.1100delC mutation is associated with an increased risk for male breast cancer in Finnish patient population. <i>BMC Cancer</i> , 2017, 17, 620.	2.6	29
33	Targeted Next-Generation Sequencing Identifies a Recurrent Mutation in MCPH1 Associating with Hereditary Breast Cancer Susceptibility. <i>PLoS Genetics</i> , 2016, 12, e1005816.	3.5	22
34	Fine-Mapping of the 1p11.2 Breast Cancer Susceptibility Locus. <i>PLoS ONE</i> , 2016, 11, e0160316.	2.5	12
35	<i>PALB2</i> , <i>CHEK2</i> and <i>ATM</i> rare variants and cancer risk: data from COGS. <i>Journal of Medical Genetics</i> , 2016, 53, 800-811.	3.2	174
36	Increased risk of certain second primary malignancies in patients treated for well-differentiated thyroid cancer. <i>International Journal of Clinical Oncology</i> , 2016, 21, 231-239.	2.2	15

#	ARTICLE	IF	CITATIONS
37	No clinical utility of KRAS variant rs61764370 for ovarian or breast cancer. <i>Gynecologic Oncology</i> , 2016, 141, 386-401.	1.4	18
38	Adjuvant capecitabine in combination with docetaxel (T), epirubicin (E), and cyclophosphamide (C) in the treatment of early breast cancer (BC): 10-year survival results from the randomized FinXX trial.. <i>Journal of Clinical Oncology</i> , 2016, 34, 1001-1001.	1.6	3
39	Toll-like receptor 9 expression is associated with breast cancer sensitivity to the growth inhibitory effects of bisphosphonates <i>in vitro</i> and <i>in vivo</i> . <i>Oncotarget</i> , 2016, 7, 87373-87389.	1.8	11
40	Bevacizumab Combined with Docetaxel or Paclitaxel as First-line Treatment of HER2-negative Metastatic Breast Cancer. <i>Anticancer Research</i> , 2016, 36, 6431-6438.	1.1	21
41	Common germline polymorphisms associated with breast cancer-specific survival. <i>Breast Cancer Research</i> , 2015, 17, 58.	5.0	26
42	Fine-Scale Mapping of the 5q11.2 Breast Cancer Locus Reveals at Least Three Independent Risk Variants Regulating MAP3K1. <i>American Journal of Human Genetics</i> , 2015, 96, 5-20.	6.2	76
43	<i>KEAP1</i> Genetic Polymorphisms Associate with Breast Cancer Risk and Survival Outcomes. <i>Clinical Cancer Research</i> , 2015, 21, 1591-1601.	7.0	37
44	Genome-wide association analysis of more than 120,000 individuals identifies 15 new susceptibility loci for breast cancer. <i>Nature Genetics</i> , 2015, 47, 373-380.	21.4	513
45	Polymorphisms in a Putative Enhancer at the 10q21.2 Breast Cancer Risk Locus Regulate NRBF2 Expression. <i>American Journal of Human Genetics</i> , 2015, 97, 22-34.	6.2	37
46	MicroRNA Related Polymorphisms and Breast Cancer Risk. <i>PLoS ONE</i> , 2014, 9, e109973.	2.5	49
47	Genetic Predisposition to In Situ and Invasive Lobular Carcinoma of the Breast. <i>PLoS Genetics</i> , 2014, 10, e1004285.	3.5	39
48	2q36.3 is associated with prognosis for oestrogen receptor-negative breast cancer patients treated with chemotherapy. <i>Nature Communications</i> , 2014, 5, 4051.	12.8	16
49	Evidence that breast cancer risk at the 2q35 locus is mediated through IGFBP5 regulation. <i>Nature Communications</i> , 2014, 5, 4999.	12.8	105
50	Genetic variation at CYP3A is associated with age at menarche and breast cancer risk: a case-control study. <i>Breast Cancer Research</i> , 2014, 16, R51.	5.0	14
51	Multiple independent variants at the TERT locus are associated with telomere length and risks of breast and ovarian cancer. <i>Nature Genetics</i> , 2013, 45, 371-384.	21.4	493
52	Adjuvant Capecitabine, Docetaxel, Cyclophosphamide, and Epirubicin for Early Breast Cancer: Final Analysis of the Randomized FinXX Trial. <i>Journal of Clinical Oncology</i> , 2012, 30, 11-18.	1.6	114
53	Adjuvant capecitabine in combination with docetaxel and cyclophosphamide plus epirubicin for breast cancer: an open-label, randomised controlled trial. <i>Lancet Oncology</i> , The, 2009, 10, 1145-1151.	10.7	65
54	Prognostic factors in differentiated thyroid carcinomas and their implications for current staging classifications. <i>Endocrine-Related Cancer</i> , 2004, 11, 571-579.	3.1	122

#	ARTICLE	IF	CITATIONS
55	Aberrant type I and type III collagen gene expression in human breast cancer in vivo. Journal of Pathology, 1998, 186, 262-268.	4.5	216
56	Aberrant type I and type III collagen gene expression in human breast cancer in vivo. , 1998, 186, 262.		2