John G Kuruvilla

List of Publications by Citations

Source: https://exaly.com/author-pdf/5323376/john-g-kuruvilla-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

96 papers

4,140 citations

23 h-index

64 g-index

118 ext. papers

5,053 ext. citations

avg, IF

5.18 L-index

#	Paper	IF	Citations
96	Outcomes in refractory diffuse large B-cell lymphoma: results from the international SCHOLAR-1 study. <i>Blood</i> , 2017 , 130, 1800-1808	2.2	653
95	Nivolumab for classical HodgkinQ lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort, single-arm phase 2 trial. <i>Lancet Oncology, The</i> , 2016 , 17, 1283-94	21.7	643
94	Programmed Death-1 Blockade With Pembrolizumab in Patients With Classical Hodgkin Lymphoma After Brentuximab Vedotin Failure. <i>Journal of Clinical Oncology</i> , 2016 , 34, 3733-3739	2.2	486
93	Nivolumab for Relapsed/Refractory Classic Hodgkin Lymphoma After Failure of Autologous Hematopoietic Cell Transplantation: Extended Follow-Up of the Multicohort Single-Arm Phase II CheckMate 205 Trial. <i>Journal of Clinical Oncology</i> , 2018 , 36, 1428-1439	2.2	377
92	Randomized comparison of gemcitabine, dexamethasone, and cisplatin versus dexamethasone, cytarabine, and cisplatin chemotherapy before autologous stem-cell transplantation for relapsed and refractory aggressive lymphomas: NCIC-CTG LY.12. <i>Journal of Clinical Oncology</i> , 2014 , 32, 3490-6	2.2	251
91	Safety and tolerability of pembrolizumab in patients with relapsed/refractory primary mediastinal large B-cell lymphoma. <i>Blood</i> , 2017 , 130, 267-270	2.2	201
90	Pretreatment with anti-thymocyte globulin versus no anti-thymocyte globulin in patients with haematological malignancies undergoing haemopoietic cell transplantation from unrelated donors: a randomised, controlled, open-label, phase 3, multicentre trial. <i>Lancet Oncology, The</i> , 2016 , 17, 164-17	21.7 73	199
89	Major Histocompatibility Complex Class II and Programmed Death Ligand 1 Expression Predict Outcome After Programmed Death 1 Blockade in Classic Hodgkin Lymphoma. <i>Journal of Clinical Oncology</i> , 2018 , 36, 942-950	2.2	175
88	How I treat relapsed and refractory Hodgkin lymphoma. <i>Blood</i> , 2011 , 117, 4208-17	2.2	106
87	Single-agent ibrutinib in relapsed or refractory follicular lymphoma: a phase 2 consortium trial. <i>Blood</i> , 2018 , 131, 182-190	2.2	92
86	Selective inhibition of nuclear export with selinexor in patients with non-Hodgkin lymphoma. <i>Blood</i> , 2017 , 129, 3175-3183	2.2	88
85	Autologous and allogeneic stem-cell transplantation for transformed follicular lymphoma: a report of the Canadian blood and marrow transplant group. <i>Journal of Clinical Oncology</i> , 2013 , 31, 1164-71	2.2	80
84	Brentuximab vedotin plus bendamustine in relapsed or refractory Hodgkin@lymphoma: an international, multicentre, single-arm, phase 1-2 trial. <i>Lancet Oncology, The</i> , 2018 , 19, 257-266	21.7	67
83	Promising SINEs for embargoing nuclear-cytoplasmic export as an anticancer strategy. <i>Cancer Discovery</i> , 2014 , 4, 527-37	24.4	67
82	Autologous stem cell transplantation in follicular lymphoma: a systematic review and meta-analysis. <i>Journal of the National Cancer Institute</i> , 2012 , 104, 18-28	9.7	61
81	Standard therapy of advanced Hodgkin lymphoma. <i>Hematology American Society of Hematology Education Program</i> , 2009 , 497-506	3.1	54
80	Pembrolizumab versus brentuximab vedotin in relapsed or refractory classical Hodgkin lymphoma (KEYNOTE-204): an interim analysis of a multicentre, randomised, open-label, phase 3 study. <i>Lancet Oncology, The</i> , 2021 , 22, 512-524	21.7	36

Checkmate 205 Update with Minimum 12-Month Follow up: A Phase 2 Study of Nivolumab in Patients with Relapsed/Refractory Classical Hodgkin Lymphoma. <i>Blood</i> , 2016 , 128, 1110-1110	2.2	33
Addition of anti-thymocyte globulin to standard graft-versus-host disease prophylaxis versus standard treatment alone in patients with haematological malignancies undergoing transplantation from unrelated donors: final analysis of a randomised, open-label, multicentre,	14.6	32
Earlier Steroid Use with Axicabtagene Ciloleucel (Axi-Cel) in Patients with Relapsed/Refractory Large B Cell Lymphoma. <i>Blood</i> , 2019 , 134, 243-243	2.2	32
Validation of National Institutes of Health global scoring system for chronic graft-versus-host disease (GVHD) according to overall and GVHD-specific survival. <i>Biology of Blood and Marrow Transplantation</i> , 2014 , 20, 556-63	4.7	27
Salvage chemotherapy and autologous stem cell transplantation for transformed indolent lymphoma: a subset analysis of NCIC CTG LY12. <i>Blood</i> , 2015 , 126, 733-8	2.2	26
Chemoimmunotherapy resistant follicular lymphoma: predictors of resistance, association with transformation and prognosis. <i>Leukemia and Lymphoma</i> , 2014 , 55, 2502-7	1.9	25
The CDK inhibitor AT7519M in patients with relapsed or refractory chronic lymphocytic leukemia (CLL) and mantle cell lymphoma. A Phase II study of the Canadian Cancer Trials Group. <i>Leukemia and Lymphoma</i> , 2017 , 58, 1358-1365	1.9	23
Pembrolizumab in Patients with Classical Hodgkin Lymphoma after Brentuximab Vedotin Failure: Long-Term Efficacy from the Phase 1b Keynote-013 Study. <i>Blood</i> , 2016 , 128, 1108-1108	2.2	23
The role of autologous and allogeneic stem cell transplantation in the management of indolent B-cell lymphoma. <i>Blood</i> , 2016 , 127, 2093-100	2.2	22
Second-line salvage chemotherapy for transplant-eligible patients with Hodgkin@lymphoma resistant to platinum-containing first-line salvage chemotherapy. <i>Haematologica</i> , 2012 , 97, 751-7	6.6	21
A phase II study of thalidomide and vinblastine for palliative patients with Hodgkin@lymphoma. <i>Hematology</i> , 2006 , 11, 25-9	2.2	17
Different response to salvage chemotherapy but similar post-transplant outcomes in patients with relapsed and refractory Hodgkin@lymphoma. <i>Haematologica</i> , 2010 , 95, 1496-502	6.6	14
Incidence, risk factors, and long-term outcomes of sclerotic graft-versus-host disease after allogeneic hematopoietic cell transplantation. <i>Biology of Blood and Marrow Transplantation</i> , 2014 , 20, 1751-7	4.7	13
Phase 1b Study of PD-1 Blockade with Pembrolizumab in Patients with Relapsed/Refractory Primary Mediastinal Large B-Cell Lymphoma (PMBCL). <i>Blood</i> , 2015 , 126, 3986-3986	2.2	13
A Canadian evidence-based guideline for the first-line treatment of follicular lymphoma: joint consensus of the Lymphoma Canada Scientific Advisory Board. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2015 , 15, 59-74	2	12
Favorable outcomes from allogeneic and autologous stem cell transplantation for patients with transformed nonfollicular indolent lymphoma. <i>Biology of Blood and Marrow Transplantation</i> , 2014 , 20, 1813-8	4.7	12
A phase I study of romidepsin, gemcitabine, dexamethasone and cisplatin combination therapy in the treatment of peripheral T-cell and diffuse large B-cell lymphoma; the Canadian cancer trials group LY.15 study [] Leukemia and Lymphoma, 2019, 60, 912-919	1.9	12
Bleomycin pulmonary toxicity does not adversely affect the outcome of patients with Hodgkin lymphoma. <i>Leukemia and Lymphoma</i> , 2017 , 58, 2607-2614	1.9	10
	Patients with Relapsed/Refractory Classical Hodgkin Lymphoma. <i>Blood</i> , 2016, 128, 1110-1110 Addition of anti-thymocyte globulin to standard graft-wersus-host disease prophylaxis versus standard treatment alone in patients with haematological malignancies undergoing transplantation from unrelated donors: final analysis of a randomised, open-label, multicentre, Earlier Steroid Use with Axicabtagene Ciloleucel (Axi-Cel) in Patients with Relapsed/Refractory Large B Cell Lymphoma. <i>Blood</i> , 2019, 134, 243-243 Validation of National Institutes of Health global scoring system for chronic graft-versus-host disease (GVHD) according to overall and GVHD-specific survival. <i>Biology of Blood and Marrow Transplantation</i> , 2014, 20, 556-63 Salvage chemotherapy and autologous stem cell transplantation for transformed indolent lymphoma: a subset analysis of NCIC CTG LV12. <i>Blood</i> , 2015, 126, 733-8 Chemoimmunotherapy resistant follicular lymphoma: predictors of resistance, association with transformation and prognosis. <i>Leukemia and Lymphoma</i> , 2014, 55, 2502-7 The CDK inhibitor AT7519M in patients with relapsed or refractory chronic lymphocytic leukemia (CLL) and mantle cell lymphoma. A Phase II study of the Canadian Cancer Trials Group. <i>Leukemia and Lymphoma</i> , 2017, 58, 1358-1365 Pembrolizumab in Patients with Classical Hodgkin Lymphoma after Brentuximab Vedotin Failure: Long-Term Efficacy from the Phase 1b Keynote-013 Study. <i>Blood</i> , 2016, 128, 1108-1108 The role of autologous and allogeneic stem cell transplantation in the management of indolent B-cell lymphoma. <i>Blood</i> , 2016, 127, 2093-100 Second-line salvage chemotherapy for transplant-eligible patients with Hodgkin@lymphoma. <i>Hematology</i> , 2006, 11, 25-9 Different response to salvage chemotherapy but similar post-transplant outcomes in patients with relapsed and refractory Hodgkin@lymphoma. <i>Hematology</i> , 2006, 11, 25-9 Different response to salvage chemotherapy but similar post-transplant outcomes in patients with relapsed and refractory Hodgkin@lymphoma. <i>Hematolo</i>	Addition of anti-thymocyte globulin to standard graft-versus-host disease prophylaxis versus standard treatment alone in patients with haematological malignancies undergoing transplantation from unrelated donors: final analysis of a randomised, open-label, multicentre, Lariler Steroid Use with Axicabtagene Ciloleucel (Axi-Cel) in Patients with Relapsed/Refractory Large B Cell Lymphoma. Blood, 2019, 134, 243-243 2.2 Validation of National Institutes of Health global scoring system for chronic graft-versus-host disease (GVHD) according to overall and GVHD-specific survival. Biology of Blood and Marrow Transplantation, 2014, 20, 556-63 Salvage chemotherapy and autologous stem cell transplantation for transformed indolent lymphoma: a subset analysis of NCIC CTG LY12. Blood, 2015, 126, 733-8 Chemoimmunotherapy resistant follicular lymphoma: predictors of resistance, association with transformation and prognosis. Leukemia and Lymphoma, 2014, 55, 2502-7 The CDK inhibitor AT7519M in patients with relapsed or refractory chronic lymphocytic leukemia (CLL) and mantle cell lymphoma. A Phase II study of the Canadian Cancer Trials Group. Leukemia and Lymphoma, 2017, 58, 1358-1365 Pembrolizumab in Patients with Classical Hodgkin Lymphoma after Brentuximab Vedotin Failure: Long-Term Efficacy from the Phase 1b Keynote-013 Study. Blood, 2016, 128, 1108-1108 The role of autologous and allogeneic stem cell transplantation in the management of indolent B-cell lymphoma. Blood, 2016, 127, 2093-100 Different response to salvage chemotherapy for transplant-eligible patients with HodgkinQ lymphoma. Persistant to platinum-containing first-line salvage chemotherapy. Haematologica, 2012, 97, 751-7 A phase II study of thalidomide and vinblastine for palliative patients with HodgkinQ lymphoma. Persistant to platinum-containing first-line salvage chemotherapy. Haematologica, 2012, 97, 751-7 A phase II study of thalidomide and vinblastine for sell transplantation in patients with Relapsed/Refractory Primary Mediastinal Large B-Cell

61	Pembrolizumab for the treatment of diffuse large B-cell lymphoma. <i>Expert Opinion on Biological Therapy</i> , 2019 , 19, 1119-1126	5.4	10
60	Modified EBMT Pretransplant Risk Score Can Identify Favorable-risk Patients Undergoing Allogeneic Hematopoietic Cell Transplantation for AML, Not Identified by the HCT-CI Score. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2015 , 15, e73-81	2	10
59	Brentuximab vedotin and bendamustine produce high complete response rates in patients with chemotherapy refractory Hodgkin lymphoma. <i>British Journal of Haematology</i> , 2018 , 180, 757-760	4.5	10
58	Selinexor (KTP-330) - a selective inhibitor of nuclear export (SINE): anti-tumor activity in diffuse large B-cell lymphoma (DLBCL). <i>Expert Opinion on Investigational Drugs</i> , 2020 , 29, 15-21	5.9	10
57	Effect of the addition of rituximab to salvage chemotherapy prior to autologous stem cell transplant in aggressive CD20+ lymphoma: a cohort comparison from the NCIC Clinical Trials Group Study LY.12. <i>Leukemia and Lymphoma</i> , 2017 , 58, 64-69	1.9	9
56	Improved prognostic stratification power of CIBMTR risk score with the addition of absolute lymphocyte and eosinophil counts at the onset of chronic GVHD. <i>Annals of Hematology</i> , 2017 , 96, 805-8	13	8
55	Earlier corticosteroid use for adverse event management in patients receiving axicabtagene ciloleucel for large B-cell lymphoma. <i>British Journal of Haematology</i> , 2021 , 195, 388-398	4.5	8
54	Limited benefit of pentostatin salvage therapy for steroid-refractory grade III-IV acute graft-versus-host disease. <i>Clinical Transplantation</i> , 2013 , 27, 930-7	3.8	7
53	Novel Therapies for Chronic Lymphocytic Leukemia: A Canadian Perspective. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2015 , 15, 627-634.e5	2	6
52	Clinical Activity of Lucatumumab (HCD122) In Patients (pts) with Relapsed/Refractory Hodgkin or Non-Hodgkin Lymphoma Treated In a Phase Ia/II Clinical Trial (NCT00670592). <i>Blood</i> , 2010 , 116, 284-284	1 ^{2.2}	6
51	A Comparison of Two-Year Outcomes in ZUMA-1 (Axicabtagene Ciloleucel) and SCHOLAR-1 in Patients with Refractory Large B Cell Lymphoma. <i>Blood</i> , 2019 , 134, 4095-4095	2.2	5
50	Mocetinostat (MGCD0103), An Isotype-Selective Histone Deacetylase (HDAC) Inhibitor, Produces Clinical Responses In Relapsed/Refractory Hodgkin Lymphoma (HL): Update From a Phase II Clinical Study. <i>Blood</i> , 2010 , 116, 1763-1763	2.2	5
49	Anti-PD-1 Antibodies as a Therapeutic Strategy in Classical Hodgkin Lymphoma. <i>Drugs</i> , 2017 , 77, 1645-1	6:5: 51	4
48	Effect of Pembrolizumab (Pembro) Monotherapy Versus Brentuximab Vedotin (BV) on Patients (Pts) with Relapsed/Refractory Classical Hodgkin Lymphoma (R/R cHL): Exploratory Analysis of the Randomized, Phase 3 Keynote-204 Study By Prior Lines of Therapy. <i>Blood</i> , 2020 , 136, 12-12	2.2	4
47	Chromosome 9p24.1/PD-L1/PD-L2Alterations and PD-L1 Expression and Treatment Outcomes in Patients with Classical Hodgkin Lymphoma Treated with Nivolumab (PD-1 Blockade). <i>Blood</i> , 2016 , 128, 2923-2923	2.2	4
46	Treatment approaches in relapsed or refractory peripheral T-cell lymphomas. <i>F1000Research</i> , 2020 , 9,	3.6	4
45	Blood-based biomarkers of cancer-related cognitive impairment in non-central nervous system cancer: protocol for a scoping review. <i>BMJ Open</i> , 2018 , 8, e017578	3	3
44	Updated Results of a Phase I Study of Ibrutinib and Lenalidomide in Patients with Relapsed and Refractory B-Cell Non-Hodgkin@Lymphoma. <i>Blood</i> , 2015 , 126, 3983-3983	2.2	3

43	Computerized cognitive training in post-treatment hematological cancer survivors: a feasibility study. <i>Pilot and Feasibility Studies</i> , 2021 , 7, 36	1.9	3
42	Developing anti-CD19 antibody-based combinations in lymphoma. Lancet Oncology, The, 2020, 21, 870-	8 72 .7	2
41	Significance of treatment response when managing patients with primary central nervous system lymphoma. <i>Leukemia and Lymphoma</i> , 2019 , 60, 349-357	1.9	2
40	A risk model for relapsed/refractory aggressive NHL integrating clinical risk factors and pretransplant Deauville score. <i>Blood Advances</i> , 2020 , 4, 5762-5771	7.8	2
39	Efficacy and safety of stem cell mobilization following gemcitabine, dexamethasone, cisplatin (GDP) salvage chemotherapy in patients with relapsed or refractory lymphoma. <i>Leukemia and Lymphoma</i> , 2020 , 61, 2153-2160	1.9	1
38	Gamma Delta T Cell Subset V Gamma 2+ Expansion Associated with Longterm Infliximab Treatment in a Patient with Ankylosing Spondylitis. <i>Journal of Rheumatology</i> , 2016 , 43, 2079-2082	4.1	1
37	A phase IV, randomized, multicenter, open-label trial comparing efficacy and systemic exposure for a standard weight-based dose versus a fixed dose of plerixafor in combination with G-CSF in patients with Non-Hodgkin@lymphoma weighing 100 kg. Bone Marrow Transplantation, 2019, 54, 258-26	4·4 54	1
36	Phase Ib study of combinations of avadomide (CC-122), CC-223, CC-292, and rituximab in patients with relapsed/refractory diffuse large B-cell lymphoma. <i>EJHaem</i> , 2022 , 3, 139-153	0.9	1
35	The Association between Patient Characteristics and the Efficacy and Safety of Selinexor in Diffuse Large B-Cell Lymphoma in the SADAL Study <i>Cancers</i> , 2022 , 14,	6.6	1
34	Outcomes of Salvage Chemotherapy and Autologous Stem Cell Transplantation for Relapsed or Refractory Primary Mediastinal Large B Cell Lymphoma (PMLCL) Are Inferior to Diffuse Large B Cell Lymphoma (DLBCL) <i>Blood</i> , 2005 , 106, 2085-2085	2.2	1
33	Predictors of the Trajectory of Neurocognitive Functioning in the First Six Months after Allogeneic Hematopoietic Stem Cell Transplantation. <i>Blood</i> , 2015 , 126, 3289-3289	2.2	1
32	Quality-of-life analysis of pembrolizumab vs brentuximab vedotin for relapsed/refractory classical Hodgkin lymphoma. <i>Blood Advances</i> , 2021 ,	7.8	1
31	Evolving management strategies for lymphomas during the COVID-19 pandemic. <i>Leukemia and Lymphoma</i> , 2021 , 62, 1046-1056	1.9	1
30	Autologous Stem Cell Transplantion for Adolescents and Young Adults with Relapsed and Refractory Hodgkin Lymphoma. <i>Blood</i> , 2010 , 116, 2402-2402	2.2	1
29	Prolonged progression free survival in a subset of responders to the combination of brentuximab vedotin and bendamustine in heavily treated patients with relapsed or refractory Hodgkin lymphoma: updated results from an international multi-center phase I/II experience. Leukemia and	1.9	1
28	Lymphoma, 2020, 61, 3014-3017 Risk stratification for relapsed/refractory classical Hodgkin lymphoma integrating pre-transplant Deauville score and residual metabolic tumor volume American Journal of Hematology, 2022,	7.1	1
27	Clinical Activity of CC-99282, a Novel, Oral Small Molecule Cereblon E3 Ligase Modulator (CELMoD) Agent, in Patients (Pts) with Relapsed or Refractory Non-Hodgkin Lymphoma (R/R NHL) - First Results from a Phase 1, Open-Label Study. <i>Blood</i> , 2021 , 138, 3574-3574	2.2	0
26	Quality of Life and Caregiver Burden in Patients and Their Caregivers Undergoing Outpatient Autologous Stem Cell Transplantation Compared to Inpatient Transplantation. <i>Blood</i> , 2021 , 138, 3055-	3653	Ο

25	Functional Predictors of Chemotherapy Toxicity in Elderly Lymphoma Patients: A Prospective Pilot Study. <i>Blood</i> , 2018 , 132, 4843-4843	2.2	0
24	Failure to Achieve CR with First-Line Treatment Is the Primary Cause of Treatment Failure in T-Cell Lymphoma: The Princess Margaret Cancer Centre Experience. <i>Blood</i> , 2016 , 128, 3005-3005	2.2	O
23	Radiotherapy in mantle cell lymphoma: A literature review. Hematological Oncology, 2020, 38, 223-228	1.3	0
22	Cost-effectiveness analysis of allogeneic versus autologous stem cell transplant versus chemo-immunotherapy for early relapse of follicular lymphoma within 2 years of initial therapy. Bone Marrow Transplantation, 2021, 56, 2400-2409	4.4	O
21	Long-term follow up of relapsed/refractory non-Hodgkin lymphoma patients treated with single-agent selinexor - a retrospective, single center study <i>Leukemia and Lymphoma</i> , 2022 , 1-8	1.9	0
20	Outcomes of patients with relapsed low-grade lymphoma retreated with rituximab are similar to rituximab nate patients. <i>Leukemia and Lymphoma</i> , 2019 , 60, 2576-2579	1.9	
19	Effectiveness and tolerability of first-line autologous stem cell transplant and maintenance rituximab for mantle cell lymphoma. <i>Bone Marrow Transplantation</i> , 2018 , 53, 347-351	4.4	
18	Optimal management of mantle cell lymphoma in the primary setting. <i>Expert Review of Hematology</i> , 2019 , 12, 715-721	2.8	
17	Outcomes in Relapsed/Refractory Burkitt Lymphoma: A Multi-Centre Canadian Experience. <i>Blood</i> , 2021 , 138, 2525-2525	2.2	
16	Risk Stratification for Relapsed/Refractory Classical Hodgkin Lymphoma Integrating Pretransplant Deauville Score and Residual Metabolic Tumor Volume. <i>Blood</i> , 2021 , 138, 1383-1383	2.2	
15	Less Chronic Graft-Versus-Host Disease, Immunosuppressive Therapy and Better Survival after Anti-Thymocyte Globulin in Unrelated Donor Stem Cell Transplant Recipients: Longer Follow-up of a Multicentre Cell Therapy Transplant Canada Randomized Trial. <i>Blood</i> , 2019 , 134, 875-875	2.2	
14	The Impact of Age on the Outcome of Primary Treatment for Classical Hodgkin@Lymphoma: 70 Years of Age Is a Clinical Relevant Cutpoint and the Most Important Predictor of Overall Survival. <i>Blood</i> , 2019 , 134, 1559-1559	2.2	
13	Obinutuzumab Plus Gemcitabine, Dexamethasone and Cisplatin (O-GDP) As Salvage Chemotherapy Prior to Autologous Stem Cell Transplant in Aggressive B Cell Lymphoma. <i>Blood</i> , 2019 , 134, 2027-2027	2.2	
12	Enteropathy-Associated T-Cell Lymphoma: A Single Centre Experience. <i>Blood</i> , 2015 , 126, 5057-5057	2.2	
11	Is Prior Rituximab Exposure Associated with a Difference in Outcomes in Relapsed Low-Grade Lymphomas Retreated with Rituximab in Comparison to Rituximab NaWe Patients?. <i>Blood</i> , 2016 , 128, 1795-1795	2.2	
10	Impact of Wait Times for Autologous Stem Cell Transplantation in Patients with Aggressive Non-Hodgkin Lymphoma, a Subset Analysis of the Canadian Cancer Trials Group (CCTG) LY.12 Clinical Trial. <i>Blood</i> , 2016 , 128, 690-690	2.2	
9	A Single-Center Retrospective Study of 140 Patients with Post-Transplant Lymphoproliferative Disorder Following Solid Organ Transplantation. <i>Blood</i> , 2016 , 128, 4227-4227	2.2	
8	Preliminary Results of FDG-PET Scanning after GDP Chemotherapy Prior to Autologous Stem Cell Transplant (ASCT) for Relapsed/Refractory (RR) Lymphoma. <i>Blood</i> , 2016 , 128, 4645-4645	2.2	

LIST OF PUBLICATIONS

7	Treatment Outcome of North American Patients with Adult T-Cell Leukemia/Lymphoma: Princess Margaret Cancer Centre Experience. <i>Blood</i> , 2016 , 128, 3010-3010	2.2
6	A Phase 1b, Multi-Center, Open-Label Study of Novel Combinations of CC-122, CC-223, CC-292, and Rituximab in Diffuse Large B-Cell Lymphoma: CC-122-DLBCL-001. <i>Blood</i> , 2016 , 128, 1849-1849	2.2
5	Comparing MicroRNA Expression In Aggressive and Indolent Non-Hodgkin Lymphomas Identifies a Prognostic Signature for Mantle Cell Lymphoma. <i>Blood</i> , 2010 , 116, 800-800	2.2
4	Prognostic Power of Chronic Gvhd Risk Score Model by Ibmtr Can Be Improved with Addition of Absolute Lymphocyte Counts and Eosinophil Counts At the Onset of Chronic Gvhd. <i>Blood</i> , 2012 , 120, 4184-4184	2.2
3	Pembrolizumab for relapsed or refractory Hodgkin lymphoma - Author@reply. <i>Lancet Oncology, The</i> , 2021 , 22, e235	21.7
2	An evaluation of pembrolizumab for classical Hodgkin lymphoma <i>Expert Review of Hematology</i> , 2022 , 1-9	2.8
1	Transverse myelitis after anti-CD19 directed CAR T cell therapy for relapsed large B cell lymphoma. <i>EJHaem</i> , 2022 , 3, 223-227	0.9