Axel C Carlsson

List of Publications by Citations

Source: https://exaly.com/author-pdf/532624/axel-c-carlsson-publications-by-citations.pdf
Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

174 papers	8,749 citations	28 h-index	92 g-index
183 ext. papers	11,558 ext. citations	4.7 avg, IF	5.22 L-index

#	Paper	IF	Citations
174	Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128 million children, adolescents, and adults. *Lancet, The*, **2017**, 390, 2627-2642	40	2980
173	Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet, The*, **2016**, 387, 1513-1530	40	2039
172	Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19 million participants. *Lancet, The*, **2017**, 389, 37-55	40	1100
171	Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet, The*, **2021**, 398, 957-980	40	154
170	Higher fibroblast growth factor-23 increases the risk of all-cause and cardiovascular mortality in the community. *Kidney International*, **2013**, 83, 160-6	9.9	112
169	High prevalence of diagnosis of diabetes, depression, anxiety, hypertension, asthma and COPD in the total population of Stockholm, Sweden - a challenge for public health. *BMC Public Health*, **2013**, 13, 670	4.1	86
168	Serum FGF23 and risk of cardiovascular events in relation to mineral metabolism and cardiovascular pathology. *Clinical Journal of the American Society of Nephrology: CJASN*, **2013**, 8, 781-6	6.9	86
167	Height and body-mass index trajectories of school-aged children and adolescents from 1985 to 2019 in 200 countries and territories: a pooled analysis of 2181 population-based studies with 65 million participants. *Lancet, The*, **2020**, 396, 1511-1524	40	73
166	Seven modifiable lifestyle factors predict reduced risk for ischemic cardiovascular disease and all-cause mortality regardless of body mass index: a cohort study. *International Journal of Cardiology*, **2013**, 168, 946-52	3.2	70
165	Most common diseases diagnosed in primary care in Stockholm, Sweden, in 2011. *Family Practice*, **2013**, 30, 506-13	1.9	52
164	Association of soluble tumor necrosis factor receptors 1 and 2 with nephropathy, cardiovascular events, and total mortality in type 2 diabetes. *Cardiovascular Diabetology*, **2016**, 15, 40	8.7	50
163	Gender differences and time trends in incidence and prevalence of type 2 diabetes in Sweden--a model explaining the diabetes epidemic worldwide today?. *Diabetes Research and Clinical Practice*, **2014**, 106, e90-2	7.4	42
162	Obesity, metabolic syndrome and risk of atrial fibrillation: a Swedish, prospective cohort study. *PLoS ONE*, **2015**, 10, e0127111	3.7	41
161	Contributions of mean and shape of blood pressure distribution to worldwide trends and variations in raised blood pressure: a pooled analysis of 1018 population-based measurement studies with 88.6 million participants. *International Journal of Epidemiology*, **2018**, 47, 872-883i	7.8	40
160	Time trends and gender differences in incidence and prevalence of type 1 diabetes in Sweden. *Current Diabetes Reviews*, **2013**, 9, 342-9	2.7	40
159	Association between circulating endostatin, hypertension duration, and hypertensive target-organ damage. *Hypertension*, **2013**, 62, 1146-51	8.5	37
158	Relationship of proximal tubular injury to chronic kidney disease as assessed by urinary kidney injury molecule-1 in five cohort studies. *Nephrology Dialysis Transplantation*, **2016**, 31, 1460-70	4.3	35

157	Use of Proteomics To Investigate Kidney Function Decline over 5 Years. *Clinical Journal of the American Society of Nephrology: CJASN*, **2017**, 12, 1226-1235	6.9	35
156	Endostatin level is associated with kidney injury in the elderly: findings from two community-based cohorts. *American Journal of Nephrology*, **2014**, 40, 417-24	4.6	34
155	Risk factors associated with newly diagnosed high blood pressure in men and women. *American Journal of Hypertension*, **2008**, 21, 771-7	2.3	32
154	Psychiatric comorbidities in patients with hypertension--a study of registered diagnoses 2009-2013 in the total population in Stockholm County, Sweden. *Journal of Hypertension*, **2016**, 34, 414-20; discussion 420	1.9	32
153	Barriers and facilitators to participation in a health check for cardiometabolic diseases in primary care: A systematic review. *European Journal of Preventive Cardiology*, **2018**, 25, 1326-1340	3.9	31
152	Soluble TNF receptors and kidney dysfunction in the elderly. *Journal of the American Society of Nephrology: JASN*, **2014**, 25, 1313-20	12.7	31
151	Prevalence of hypertension in immigrants and Swedish-born individuals, a cross-sectional study of 60-year-old men and women in Sweden. *Journal of Hypertension*, **2008**, 26, 2295-302	1.9	31
150	Financial stress in late adulthood and diverse risks of incident cardiovascular disease and all-cause mortality in women and men. *BMC Public Health*, **2014**, 14, 17	4.1	30
149	Barriers and facilitators among health professionals in primary care to prevention of cardiometabolic diseases: A systematic review. *Family Practice*, **2018**, 35, 383-398	1.9	29
148	Urinary kidney injury molecule 1 and incidence of heart failure in elderly men. *European Journal of Heart Failure*, **2013**, 15, 441-6	12.3	29
147	Neighbourhood socioeconomic status and coronary heart disease in individuals between 40 and 50 years. *Heart*, **2016**, 102, 775-82	5.1	28
146	Diabetes and psychiatric illness in the total population of Stockholm. *Journal of Psychosomatic Research*, **2014**, 77, 169-73	4.1	28
145	Multiplex proteomics for prediction of major cardiovascular events in type 2 diabetes. *Diabetologia*, **2018**, 61, 1748-1757	10.3	28
144	Hypertriglyceridemic waist phenotype is associated with decreased insulin sensitivity and incident diabetes in elderly men. *Obesity*, **2014**, 22, 526-9	8	27
143	The association between endostatin and kidney disease and mortality in patients with type 2 diabetes. *Diabetes and Metabolism*, **2016**, 42, 351-357	5.4	26
142	Country of birth-specific and gender differences in prevalence of diabetes in Sweden. *Diabetes Research and Clinical Practice*, **2013**, 100, 404-8	7.4	26
141	Prevalence of blood lipid disturbances in Swedish and foreign-born 60-year-old men and women in Stockholm, Sweden. *Nutrition, Metabolism and Cardiovascular Diseases*, **2011**, 21, 173-81	4.5	26
140	Differences and time trends in drug treatment of atrial fibrillation in men and women and doctors' adherence to warfarin therapy recommendations: a Swedish study of prescribed drugs in primary care in 2002 and 2007. *European Journal of Clinical Pharmacology*, **2013**, 69, 245-53	2.8	25

139	Physical activity, obesity and risk of cardiovascular disease in middle-aged men during a median of 30 years of follow-up. *European Journal of Preventive Cardiology*, **2016**, 23, 359-65	3.9	24
138	Soluble tumor necrosis factor receptor 1 (sTNFR1) is associated with increased total mortality due to cancer and cardiovascular causes - findings from two community based cohorts of elderly. *Atherosclerosis*, **2014**, 237, 236-42	3.1	24
137	Urinary neutrophil gelatinase-associated lipocalin (NGAL) is associated with mortality in a community-based cohort of older Swedish men. *Atherosclerosis*, **2013**, 227, 408-13	3.1	24
136	Physicians' gender is associated with risk factor control in patients on antihypertensive and lipid lowering treatment. *Blood Pressure*, **2010**, 19, 240-8	1.7	24
135	Neighbourhood socio-economic status and all-cause mortality in adults with atrial fibrillation: A cohort study of patients treated in primary care in Sweden. *International Journal of Cardiology*, **2016**, 202, 776-81	3.2	24
134	Circulating proteins as predictors of cardiovascular mortality in end-stage renal disease. *Journal of Nephrology*, **2019**, 32, 111-119	4.8	24
133	Gout and its comorbidities in the total population of Stockholm. *Preventive Medicine*, **2015**, 81, 387-91	4.3	23
132	HIV-infection and psychiatric illnesses - A double edged sword that threatens the vision of a contained epidemic: The Greater Stockholm HIV Cohort Study. *Journal of Infection*, **2017**, 74, 22-28	18.9	23
131	Serum Fatty Acids, Desaturase Activities and Abdominal Obesity - A Population-Based Study of 60-Year Old Men and Women. *PLoS ONE*, **2017**, 12, e0170684	3.7	22
130	The association between relevant co-morbidities and prevalent as well as incident heart failure in patients with atrial fibrillation. *Journal of Cardiology*, **2018**, 72, 26-32	3	21
129	High-sensitivity cardiac troponin T levels in the emergency department in patients with chest pain but no myocardial infarction. *International Journal of Cardiology*, **2017**, 228, 253-259	3.2	21
128	Is a unified definition of metabolic syndrome needed? Comparison of three definitions of metabolic syndrome in 60-year-old men and women. *Metabolic Syndrome and Related Disorders*, **2009**, 7, 231-41	2.6	21
127	Urinary kidney injury molecule-1 and the risk of cardiovascular mortality in elderly men. *Clinical Journal of the American Society of Nephrology: CJASN*, **2014**, 9, 1393-401	6.9	20
126	District nurses' preventive home visits to 75-year-olds: an opportunity to identify factors related to unsafe medication management. *Scandinavian Journal of Public Health*, **2014**, 42, 786-94	3	20
125	Waist/Hip Ratio Better Predicts Development of Severe Liver Disease Within 20 Years Than Body Mass Index: A Population-based Cohort Study. *Clinical Gastroenterology and Hepatology*, **2017**, 15, 1294-1301.e2	6.9	19
124	Factors associated with uncontrolled hypertension and cardiovascular risk in hypertensive 60-year-old men and women--a population-based study. *Hypertension Research*, **2009**, 32, 780-5	4.7	19
123	Investigations, findings, and follow-up in patients with chest pain and elevated high-sensitivity cardiac troponin T levels but no myocardial infarction. *International Journal of Cardiology*, **2017**, 232, 111-116	3.2	18
122	Neighborhood socioeconomic status at the age of 40 years and ischemic stroke before the age of 50 years: A nationwide cohort study from Sweden. *International Journal of Stroke*, **2017**, 12, 815-826	6.3	18

121	Neighborhood deprivation and warfarin, aspirin and statin prescription - A cohort study of men and women treated for atrial fibrillation in Swedish primary care. *International Journal of Cardiology*, **2015**, 187, 547-52	3.2	18
120	Depression or anxiety and all-cause mortality in adults with atrial fibrillation--A cohort study in Swedish primary care. *Annals of Medicine*, **2016**, 48, 59-66	1.5	18
119	Socioeconomic factors and mortality in patients with atrial fibrillation-a cohort study in Swedish primary care. *European Journal of Public Health*, **2018**, 28, 1103-1109	2.1	18
118	Soluble Tumor Necrosis Factor Receptor 1 Is Associated with Glomerular Filtration Rate Progression and Incidence of Chronic Kidney Disease in Two Community-Based Cohorts of Elderly Individuals. *CardioRenal Medicine*, **2015**, 5, 278-88	2.8	18
117	Health factors and longevity in men and women: a 26-year follow-up study. *European Journal of Epidemiology*, **2010**, 25, 547-51	12.1	17
116	Prevalence, incidence, mortality and co-morbidities amongst human immunodeficiency virus (HIV) patients in Stockholm County, Sweden - the Greater Stockholm HIV Cohort Study. *AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV*, **2015**, 27, 142-9	2.2	16
115	Mortality trends and cause of death in patients with new-onset type 2 diabetes and controls: A 24-year follow-up prospective cohort study. *Diabetes Research and Clinical Practice*, **2018**, 138, 81-89	7.4	16
114	10-Year Associations Between Tumor Necrosis Factor Receptors 1 and 2 and Cardiovascular Events in Patients With Stable Coronary Heart Disease: A CLARICOR (Effect of Clarithromycin on Mortality and Morbidity in Patients With Ischemic Heart Disease) Trial Substudy. *Journal of the American Heart Association*, **2018**, 7,	6	16
113	Kidney injury molecule (KIM)-1 is associated with insulin resistance: results from two community-based studies of elderly individuals. *Diabetes Research and Clinical Practice*, **2014**, 103, 516-21	7.4	16
112	Prediction of cardiovascular disease by abdominal obesity measures is dependent on body weight and sex--results from two community based cohort studies. *Nutrition, Metabolism and Cardiovascular Diseases*, **2014**, 24, 891-9	4.5	16
111	Measuring possible effect on health-related quality of life by tactile massage or relaxation in patients with type 2 diabetes. *Complementary Therapies in Medicine*, **2012**, 20, 8-15	3.5	16
110	Poor self-rated health is not associated with a high total allostatic load in type 2 diabetic patients--but high blood pressure is. *Diabetes and Metabolism*, **2011**, 37, 446-51	5.4	16
109	Elevated ApoB serum levels strongly predict early cardiovascular events. *Heart*, **2012**, 98, 1242-5	5.1	15
108	Growth differentiation factor 15 (GDF-15) is a potential biomarker of both diabetic kidney disease and future cardiovascular events in cohorts of individuals with type 2 diabetes: a proteomics approach. *Upsala Journal of Medical Sciences*, **2020**, 125, 37-43	2.8	15
107	Cathepsin B and S as markers for cardiovascular risk and all-cause mortality in patients with stable coronary heart disease during 10 years: a CLARICOR trial sub-study. *Atherosclerosis*, **2018**, 278, 97-102	3.1	15
106	Endostatin: a promising biomarker in the cardiovascular continuum?. *Biomarkers in Medicine*, **2017**, 11, 905-916	2.3	14
105	Cardiovascular and total mortality in men and women with different blood pressure levels--A 26-year follow-up. *Blood Pressure*, **2009**, 18, 105-10	1.7	14
104	The association between relevant comorbidities and dementia in patients with atrial fibrillation. *GeroScience*, **2018**, 40, 317	8.9	14

103	Chronic kidney disease and 10-year risk of cardiovascular death. *European Journal of Preventive Cardiology*, **2016**, 23, 1187-94	3.9	13
102	Effect of cardiovascular drug classes on all-cause mortality among atrial fibrillation patients treated in primary care in Sweden: a cohort study. *European Journal of Clinical Pharmacology*, **2013**, 69, 279-87	2.8	13
101	High levels of soluble tumor necrosis factor receptors 1 and 2 and their association with mortality in patients undergoing hemodialysis. *CardioRenal Medicine*, **2015**, 5, 89-95	2.8	12
100	Selective prevention of cardiometabolic diseases: activities and attitudes of general practitioners across Europe. *European Journal of Public Health*, **2019**, 29, 88-93	2.1	12
99	Dementia in immigrant groups: A cohort study of all adults 45 years of age and older in Sweden. *Archives of Gerontology and Geriatrics*, **2019**, 82, 251-258	4	11
98	Use of a proximity extension assay proteomics chip to discover new biomarkers associated with albuminuria. *European Journal of Preventive Cardiology*, **2017**, 24, 340-348	3.9	11
97	Endostatin, Cathepsin S, and Cathepsin L, and Their Association with Inflammatory Markers and Mortality in Patients Undergoing Hemodialysis. *Blood Purification*, **2015**, 39, 259-65	3.1	11
96	Design and methodology of the screening for CKD among older patients across Europe (SCOPE) study: a multicenter cohort observational study. *BMC Nephrology*, **2018**, 19, 260	2.7	11
95	Circulating cathepsin-S levels correlate with GFR decline and sTNFR1 and sTNFR2 levels in mice and humans. *Scientific Reports*, **2017**, 7, 43538	4.9	10
94	Levels of soluble tumor necrosis factor receptor 1 and 2, gender, and risk of myocardial infarction in Northern Sweden. *Atherosclerosis*, **2018**, 272, 41-46	3.1	10
93	Antihypertensive drugs and relevant cardiovascular pharmacotherapies and the risk of incident dementia in patients with atrial fibrillation. *International Journal of Cardiology*, **2018**, 272, 149-154	3.2	10
92	Effects of prescribed antihypertensives and other cardiovascular drugs on mortality in patients with atrial fibrillation and hypertension: a cohort study from Sweden. *Hypertension Research*, **2014**, 37, 553-9	4.7	10
91	Effects of tactile massage on metabolic biomarkers in patients with type 2 diabetes. *Diabetes and Metabolism*, **2013**, 39, 411-7	5.4	10
90	Atrial fibrillation in immigrant groups: a cohort study of all adults 45 years of age and older in Sweden. *European Journal of Epidemiology*, **2017**, 32, 785-796	12.1	10
89	Estimated glomerular filtration rate and functional status among older people: A systematic review. *European Journal of Internal Medicine*, **2018**, 56, 39-48	3.9	10
88	Diagnostic errors reported in primary healthcare and emergency departments: A retrospective and descriptive cohort study of 4830 reported cases of preventable harm in Sweden. *European Journal of General Practice*, **2019**, 25, 128-135	2.8	9
87	Differences in anthropometric measures in immigrants and Swedish-born individuals: results from two community-based cohort studies. *Preventive Medicine*, **2014**, 69, 151-6	4.3	9
86	Increased urinary cystatin C indicated higher risk of cardiovascular death in a community cohort. *Atherosclerosis*, **2014**, 234, 108-13	3.1	9

85	Urinary Osteopontin Predicts Incident Chronic Kidney Disease, while Plasma Osteopontin Predicts Cardiovascular Death in Elderly Men. *CardioRenal Medicine*, **2017**, 7, 245-254	2.8	9
84	Total mortality among levothyroxine-treated women with atrial fibrillation in Swedish primary health care. *International Journal of Cardiology*, **2011**, 152, 147-8	3.2	9
83	Effect of heart-rate recovery on long-term mortality among men and women. *International Journal of Cardiology*, **2010**, 144, 276-9	3.2	9
82	The association between serum cathepsin L and mortality in older adults. *Atherosclerosis*, **2016**, 254, 109-116	3.1	9
81	Comparison of Mortality and Nonfatal Cardiovascular Events in Adults With Atrial Fibrillation With Versus Without Levothyroxine Treatment. *American Journal of Cardiology*, **2017**, 120, 1974-1979	3	8
80	The prediction of colorectal cancer using anthropometric measures: A Swedish population-based cohort study with 22 years of follow-up. *United European Gastroenterology Journal*, **2019**, 7, 1250-1260	5.3	8
79	Low anthropometric measures and mortality--results from the Malm Diet and Cancer Study. *Annals of Medicine*, **2015**, 47, 325-31	1.5	8
78	Blood pressure measures and their predictive ability of cardiovascular mortality: a 26-year follow-up. *Blood Pressure Monitoring*, **2013**, 18, 72-7	1.3	8
77	Health care consumption and psychiatric diagnoses among adolescent girls 1 and 2 years after a first-time registered child sexual abuse experience: a cohort study in the Stockholm Region. *European Child and Adolescent Psychiatry*, **2021**, 30, 1803-1811	5.5	8
76	Warfarin treatment and risk of stroke among primary care patients with atrial fibrillation. *Scandinavian Cardiovascular Journal*, **2016**, 50, 311-316	2	8
75	Heart failure in immigrant groups: a cohort study of adults aged 45 years and over in Sweden. *Scandinavian Cardiovascular Journal*, **2018**, 52, 292-300	2	8
74	Diagnoses of sexual abuse and their common registered comorbidities in the total population of Stockholm. *Journal of Epidemiology and Community Health*, **2017**, 71, 592-598	5.1	7
73	Impact of physical activity on cardiovascular status in obesity. *European Journal of Clinical Investigation*, **2017**, 47, 167-175	4.6	7
72	Sex-specific risk of emergency department revisits and early readmission following myocardial infarction. *International Journal of Cardiology*, **2017**, 243, 54-58	3.2	7
71	Mortality in patients with atrial fibrillation and common co-morbidities - a cohort study in primary care. *Annals of Medicine*, **2018**, 50, 156-163	1.5	7
70	Association Between Relevant Co-Morbidities and Dementia With Atrial Fibrillation-A National Study. *Archives of Medical Research*, **2019**, 50, 29-35	6.6	6
69	Circulating endostatin as a risk factor for cardiovascular events in patients with stable coronary heart disease: A CLARICOR trial sub-study. *Atherosclerosis*, **2019**, 284, 202-208	3.1	6
68	Low fructosamine and mortality - A long term follow-up of 215,011 non-diabetic subjects in the Swedish AMORIS study. *Nutrition, Metabolism and Cardiovascular Diseases*, **2016**, 26, 1120-1128	4.5	6

67	Circulating endostatin and the incidence of heart failure. *Scandinavian Cardiovascular Journal*, **2018**, 52, 244-249	2	6
66	Prevalence of sarcopenia in community-dwelling older adults using the updated EWGSOP2 definition according to kidney function and albuminuria : The Screening for CKD among Older People across Europe (SCOPE) study. *BMC Geriatrics*, **2020**, 20, 327	4.1	6
65	Urinary KIM-1, but not urinary cystatin C, should be corrected for urinary creatinine. *Clinical Biochemistry*, **2016**, 49, 1164-1166	3.5	6
64	Patterns of multimorbidity and pharmacotherapy: a total population cross-sectional study. *Family Practice*, **2021**, 38, 132-140	1.9	6
63	Body composition measurements and risk of hematological malignancies: A population-based cohort study during 20 years of follow-up. *PLoS ONE*, **2018**, 13, e0202651	3.7	6
62	The association between gout and cardiovascular disease in patients with atrial fibrillation. *SN Comprehensive Clinical Medicine*, **2019**, 1, 304-310	2.7	5
61	The association between sociodemographic characteristics and dementia in patients with atrial fibrillation. *Aging Clinical and Experimental Research*, **2020**, 32, 2319-2327	4.8	5
60	Cardiovascular events in patients under age fifty with early findings of elevated lipid and glucose levels - The AMORIS study. *PLoS ONE*, **2018**, 13, e0201972	3.7	5
59	Effects of prescribed antithrombotics and other cardiovascular pharmacotherapies on all-cause mortality in patients with diabetes and atrial fibrillation - a cohort study from Sweden using propensity score analyses. *Diabetology and Metabolic Syndrome*, **2014**, 6, 2	5.6	5
58	Association between kidney function, nutritional status and anthropometric measures in older people : The Screening for CKD among Older People across Europe (SCOPE) study. *BMC Geriatrics*, **2020**, 20, 366	4.1	5
57	Predictive Capacity for Mortality and Severe Liver Disease of the Relative Fat Mass Algorithm. *Clinical Gastroenterology and Hepatology*, **2019**, 17, 2619-2620	6.9	5
56	Interleukin-8 is associated with increased total mortality in women but not in men-findings from a community-based cohort of elderly. *Annals of Medicine*, **2015**, 47, 28-33	1.5	4
55	Is medical urgency of elderly patients with traumatic brain injury underestimated by emergency department triage?. *Upsala Journal of Medical Sciences*, **2020**, 125, 58-63	2.8	4
54	Patient and provider perspectives on reducing risk of harm in primary health care: a qualitative questionnaire study in Sweden. *Scandinavian Journal of Primary Health Care*, **2020**, 38, 66-74	2.7	4
53	Patient-related factors associated with an increased risk of being a reported case of preventable harm in first-line health care: a case-control study. *BMC Family Practice*, **2020**, 21, 20	2.6	4
52	Massive open online courses (MOOCs) for long-distance education in geriatric medicine across Europe : A pilot project launched by the consortium of the project "Screening for Chronic Kidney Disease among Older People": SCOPE project. *European Geriatric Medicine*, **2019**, 10, 989-994	3	4
51	Systemic diseases and their association with open-angle glaucoma in the population of Stockholm. *International Ophthalmology*, **2021**, 1	2.2	4
50	Clinical Implications of Estimating Glomerular Filtration Rate with Three Different Equations Among Older People. Preliminary Results of the Project "Screening for Chronic Kidney Disease among Older People across Europe (SCOPE)". *Journal of Clinical Medicine*, **2020**, 9,	5.1	4

49	Chronic kidney disease in the context of multimorbidity patterns: the role of physical performance : The screening for CKD among older people across Europe (SCOPE) study. *BMC Geriatrics*, **2020**, 20, 350	4.1	4
48	Warfarin treatment and risk of myocardial infarction - A cohort study of patients with atrial fibrillation treated in primary health care. *International Journal of Cardiology*, **2016**, 221, 789-93	3.2	4
47	Effect of Levothyroxine Treatment on Incident Dementia in Adults with Atrial Fibrillation and Hypothyroidism. *Clinical Drug Investigation*, **2019**, 39, 187-195	3.2	4
46	Levothyroxine treatment is associated with an increased relative risk of overall and organ specific incident cancers - a cohort study of the Swedish population. *Cancer Epidemiology*, **2020**, 66, 101707	2.8	4
45	Gout in immigrant groups: a cohort study in Sweden. *Clinical Rheumatology*, **2017**, 36, 1091-1102	3.9	3
44	The use of anthropometric measures in the prediction of incident gout: results from a Swedish community-based cohort study. *Scandinavian Journal of Rheumatology*, **2019**, 48, 294-299	1.9	3
43	Associations between urinary kidney injury biomarkers and cardiovascular mortality risk in elderly men with diabetes. *Upsala Journal of Medical Sciences*, **2016**, 121, 174-8	2.8	3
42	Associations between relevant cardiovascular pharmacotherapies and incident heart failure in patients with atrial fibrillation: a cohort study in primary care. *Journal of Hypertension*, **2018**, 36, 1929-1935	1.9	3
41	Effect of cardiovascular drugs on mortality in atrial fibrillation and chronic heart failure. *Scandinavian Cardiovascular Journal*, **2014**, 48, 291-8	2	3
40	Lack of long-term effects from a pedagogical intervention upon blood pressure control--a randomized primary care study. *Scandinavian Cardiovascular Journal*, **2013**, 47, 289-96	2	3
39	Health care consumption among adolescent girls prior to diagnoses of sexual abuse, a case-control study in the Stockholm Region. *European Child and Adolescent Psychiatry*, **2020**, 29, 1363-1369	5.5	3
38	Impaired kidney function is associated with lower quality of life among community-dwelling older adults : The screening for CKD among older people across Europe (SCOPE) study. *BMC Geriatrics*, **2020**, 20, 340	4.1	3
37	The association between plasma proteomics and incident cardiovascular disease identifies MMP-12 as a promising cardiovascular risk marker in patients with chronic kidney disease. *Atherosclerosis*, **2020**, 307, 11-15	3.1	3
36	The impact of diabetes, education and income on mortality and cardiovascular events in hypertensive patients: A cohort study from the Swedish Primary Care Cardiovascular Database (SPCCD). *PLoS ONE*, **2020**, 15, e0237107	3.7	3
35	The association between circulating endostatin levels and incident myocardial infarction. *Scandinavian Cardiovascular Journal*, **2018**, 52, 315-319	2	3
34	Endothelial dysfunction is associated with impaired lung function in two independent community cohorts. *Respiratory Medicine*, **2018**, 143, 123-128	4.6	3
33	An evidence-based toolbox for the design and implementation of selective-prevention primary-care initiatives targeting cardio-metabolic disease. *Preventive Medicine Reports*, **2019**, 16, 100979	2.6	2
32	Parkinson's Disease Among Immigrant Groups and Swedish-Born Individuals: A Cohort Study of All Adults 50 Years of Age and Older in Sweden. *Journal of Parkinsonls Disease*, **2020**, 10, 1133-1141	5.3	2

31	An audit & feedback intervention for improved anticoagulant use in patients with atrial fibrillation in primary care. *International Journal of Cardiology*, **2020**, 310, 67-72	3.2	2
30	Distal forearm fractures in immigrant groups: A national Swedish study. *Bone*, **2020**, 138, 115508	4.7	2
29	Urolithiasis in immigrant groups: a nationwide cohort study in Sweden. *Scandinavian Journal of Urology*, **2019**, 53, 69-76	1.6	2
28	Personality measured as Murray's psychological needs and all-cause mortality: 41years of follow-up of a population-based sample. *Personality and Individual Differences*, **2014**, 68, 32-36	3.3	2
27	Endostatin predicts mortality in patients with acute dyspnea - A cohort study of patients seeking care in emergency departments. *Clinical Biochemistry*, **2020**, 75, 35-39	3.5	2
26	Is kidney function associated with cognition and mood in late life? : The Screening for CKD among Older People across Europe (SCOPE) study. *BMC Geriatrics*, **2020**, 20, 297	4.1	2
25	Multiple sclerosis among first- and second-generation immigrant groups in Sweden. *Acta Neurologica Scandinavica*, **2020**, 142, 339-349	3.8	2
24	Levothyroxine treatment and incident dementia in adults with atrial fibrillation. *Aging Clinical and Experimental Research*, **2020**, 32, 433-439	4.8	2
23	What should selective cardiometabolic prevention programmes in European primary care look like? A consensus-based design by the SPIMEU group. *European Journal of General Practice*, **2019**, 25, 101-108	2.8	1
22	End-Stage Kidney Diseases in Immigrant Groups: A Nationwide Cohort Study in Sweden. *American Journal of Nephrology*, **2019**, 49, 186-192	4.6	1
21	Validation and initial results of surveys exploring perspectives on risks and solutions for diagnostic and medication errors in primary care in Sweden. *Scandinavian Journal of Primary Health Care*, **2020**, 38, 381-390	2.7	1
20	Epilepsy in immigrants and Swedish-born: A cohort study of all adults 18 years of age and older in Sweden. *Seizure: the Journal of the British Epilepsy Association*, **2020**, 76, 116-122	3.2	1
19	The association between circulating endostatin and a disturbed circadian blood pressure pattern in patients with type 2 diabetes. *Blood Pressure*, **2018**, 27, 215-221	1.7	1
18	Response to: Modifiable lifestyle risks, cardiovascular disease, and all-cause mortality. *International Journal of Cardiology*, **2014**, 173, 560	3.2	1
17	High levels of leptin are associated with poor self-rated health in men and women with type 2 diabetes treated with metformin. *Nutrition, Metabolism and Cardiovascular Diseases*, **2013**, 23, e11-2	4.5	1
16	Cardiovascular medication in relation to renal function after coronary artery bypass surgery. *International Journal of Cardiology*, **2013**, 168, 4033-8	3.2	1
15	Association between antithrombotic treatment and hemorrhagic stroke in patients with atrial fibrillation-a cohort study in primary care. *European Journal of Clinical Pharmacology*, **2017**, 73, 215-221	2.8	1
14	Country of birth and mortality risk in hypertension with and without diabetes: the Swedish primary care cardiovascular database. *Journal of Hypertension*, **2021**, 39, 1155-1162	1.9	1

13	Kidney function and other factors and their association with falls : The screening for CKD among older people across Europe (SCOPE) study. *BMC Geriatrics*, **2020**, 20, 320	4.1	1
12	Use of cardiovascular drugs and risk of incident heart failure in patients with atrial fibrillation. *Journal of Clinical Hypertension*, **2020**, 22, 1396-1405	2.3	1
11	Hearing impairment among children in Sweden with foreign-born parents and natives: A national Swedish study. *Acta Paediatrica, International Journal of Paediatrics*, **2021**, 110, 2817-2824	3.1	1
10	Delayed healthcare access among victims of sexual abuse, understood through internal and external gatekeeping mechanisms. *Nordic Journal of Psychiatry*, **2021**, 75, 370-377	2.3	1
9	Osteoporotic fractures in second-generation immigrants and Swedish natives. *Osteoporosis International*, **2021**, 32, 1343-1350	5.3	1
8	Visual impairment among children in Sweden with foreign-born parents and natives: a national Swedish study. *Acta Paediatrica, International Journal of Paediatrics*, **2021**,	3.1	1
7	Unwillingness to participate in health checks for cardiometabolic diseases: A survey among primary health care patients in five European countries. *Health Science Reports*, **2021**, 4, e256	2.2	0
6	Effects of cardiovascular pharmacotherapies on incident dementia in patients with atrial fibrillation: A cohort study of all patients above 45 years diagnosed with AF in hospitals in Sweden. *International Journal of Cardiology*, **2019**, 297, 55-60	3.2	0
5	Atrial fibrillation in immigrants under the age of 45 y in Sweden. *International Health*, **2019**, 11, 193-202	2.4	0
4	Huntington's disease among immigrant groups and Swedish-born individuals: a cohort study of all adults 18 years of age and older in Sweden. *Neurological Sciences*, **2021**, 42, 3851-3856	3.5	0
3	Reply. *Journal of Hypertension*, **2016**, 34, 1441-2	1.9	
2	Urolithiasis in second-generation immigrant children younger than 18 years of age in Sweden. *Acta Paediatrica, International Journal of Paediatrics*, **2021**, 110, 340-346	3.1	
1	Amyotrophic lateral sclerosis (ALS) among immigrant groups and Swedish-born individuals: a cohort study of all adults 18 years of age and older in Sweden. *Journal of Neurology*, **2021**, 1	5.5	