
Jean-Christophe Bories

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5328962/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Bi38-3 is a novel CD38/CD3 bispecific T-cell engager with low toxicity for the treatment of multiple myeloma. Haematologica, 2021, 106, 1193-1197. | 3.5 | 23 |
| 2 | The long non-coding RNA CRNDE regulates growth of multiple myeloma cells via an effect on IL6 signalling. Leukemia, 2021, 35, 1710-1721. | 7.2 | 26 |
| 3 | Hematopoietic progenitors polarize in contact with bone marrow stromal cells in response to SDF1. Journal of Cell Biology, 2021, 220, . | 5.2 | 8 |
| 4 | Glomerular common gamma chain confers B- and T-cell–independent protection against glomerulonephritis. Kidney International, 2017, 91, 1146-1158. | 5.2 | 15 |
| 5 | Histone methyltransferase MMSET promotes AID-mediated DNA breaks at the donor switch region during class switch recombination. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, E10560-E10567. | 7.1 | 13 |
| 6 | Age-Associated Decrease of the Histone Methyltransferase SUV39H1 in HSC Perturbs Heterochromatin and B Lymphoid Differentiation. Stem Cell Reports, 2016, 6, 970-984. | 4.8 | 88 |
| 7 | TNFA, ANXA11 and BTNL2 Polymorphisms in CVID Patients with Granulomatous Disease. Journal of Clinical Immunology, 2016, 36, 110-112. | 3.8 | 7 |
| 8 | Dynamic recruitment of Ets1 to both nucleosome-occupied and -depleted enhancer regions mediates a transcriptional program switch during early T-cell differentiation. Nucleic Acids Research, 2016, 44, 3567-3585. | 14.5 | 39 |
| 9 | Global Genomic Analysis of Newly Diagnosed t(4 ;14) Multiple Myeloma Reveals a Specific Mutational Spectrum and Identifies PKD2 As a Potential Therapeutic Target. Blood, 2016, 128, 4462-4462. | 1.4 | 0 |
| 10 | Mice Haploinsufficient for Ets1 and Fli1 Display Middle Ear Abnormalities and Model Aspects of Jacobsen Syndrome. American Journal of Pathology, 2015, 185, 1867-1876. | 3.8 | 15 |
| 11 | The human immunodeficiency virus-1 protease inhibitor nelfinavir impairs proteasome activity and inhibits the proliferation of multiple myeloma cells in vitro and in vivo. Haematologica, 2012, 97, 1101-1109. | 3.5 | 42 |
| 12 | The Ets-1 transcription factor is required for Stat1-mediated T-bet expression and IgG2a class switching in mouse B cells. Blood, 2012, 119, 4174-4181. | 1.4 | 39 |
| 13 | Defects in the CD19 complex predispose to glomerulonephritis, as well as IgG1 subclass deficiency. Journal of Allergy and Clinical Immunology, 2011, 127, 538-541.e5. | 2.9 | 31 |
| 14 | The Ets-1 transcription factor controls the development and function of natural regulatory T cells. Journal of Experimental Medicine, 2010, 207, 2113-2125. | 8.5 | 98 |
| 15 | p16INK4A tumor suppressor gene expression and CD3ïµ deficiency but not pre-TCR deficiency inhibit TAL1-linked T-lineage leukemogenesis. Blood, 2007, 110, 2610-2619. | 1.4 | 14 |