

Katrina Albert

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5329226/publications.pdf>

Version: 2024-02-01

12
papers

302
citations

933447

10
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

444
citing authors

#	ARTICLE	IF	CITATIONS
1	Utilising Induced Pluripotent Stem Cells in Neurodegenerative Disease Research: Focus on Glia. <i>International Journal of Molecular Sciences</i> , 2021, 22, 4334.	4.1	14
2	Cerebral dopamine neurotrophic factor reduces α -synuclein aggregation and propagation and alleviates behavioral alterations in vivo. <i>Molecular Therapy</i> , 2021, 29, 2821-2840.	8.2	26
3	$\langle \text{scp} \rangle \text{GDNF} \langle / \text{scp} \rangle \langle \text{scp} \rangle \text{RET} \langle / \text{scp} \rangle$ Signaling Pathway Activation Eliminates Lewy Body Pathology in Midbrain Dopamine Neurons. <i>Movement Disorders</i> , 2020, 35, 2279-2289.	3.9	27
4	Back and to the Future: From Neurotoxin-Induced to Human Parkinson's Disease Models. <i>Current Protocols in Neuroscience</i> , 2020, 91, e88.	2.6	36
5	Neuroprotective and reparative effects of endoplasmic reticulum luminal proteins α -mesencephalic astrocyte-derived neurotrophic factor and cerebral dopamine neurotrophic factor. <i>Croatian Medical Journal</i> , 2019, 60, 99-109.	0.7	17
6	Cerebral Dopamine Neurotrophic Factor Diffuses Around the Brainstem and Does Not Undergo Anterograde Transport After Injection to the Substantia Nigra. <i>Frontiers in Neuroscience</i> , 2019, 13, 590.	2.8	7
7	Post-stroke Intranasal (+)-Naloxone Delivery Reduces Microglial Activation and Improves Behavioral Recovery from Ischemic Injury. <i>ENeuro</i> , 2018, 5, ENEURO.0395-17.2018.	1.9	35
8	Downregulation of tyrosine hydroxylase phenotype after AAV injection above substantia nigra: Caution in experimental models of Parkinson's disease. <i>Journal of Neuroscience Research</i> , 2018, 97, 346-361.	2.9	24
9	Nigral injection of a proteasomal inhibitor, lactacystin, induces widespread glial cell activation and shows various phenotypes of Parkinson's disease in young and adult mouse. <i>Experimental Brain Research</i> , 2017, 235, 2189-2202.	1.5	22
10	AAV Vector-Mediated Gene Delivery to Substantia Nigra Dopamine Neurons: Implications for Gene Therapy and Disease Models. <i>Genes</i> , 2017, 8, 63.	2.4	43
11	Characterization of a new low-dose 6-hydroxydopamine model of Parkinson's disease in rat. <i>Journal of Neuroscience Research</i> , 2016, 94, 318-328.	2.9	39
12	Comparison of the MK-801-induced appetitive extinction deficit with pressing for reward and associated pERK1/2 staining in prefrontal cortex and nucleus accumbens. <i>Behavioural Brain Research</i> , 2012, 228, 194-202.	2.2	10