

Dutra-Filho, Cs

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5329420/dutra-filho-cs-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

173
papers

4,451
citations

39
h-index

53
g-index

173
ext. papers

4,763
ext. citations

3.8
avg, IF

4.63
L-index

#	Paper	IF	Citations
173	Preliminary results of PBA-loaded nanoparticles development and the effect on oxidative stress and neuroinflammation in rats submitted to a chemically induced chronic model of MSUD. <i>Metabolic Brain Disease</i> , 2021 , 36, 1015-1027	3.9	1
172	Effects of Fish and Grape Seed Oils as Core of Haloperidol-Loaded Nanocapsules on Oral Dyskinesia in Rats. <i>Neurochemical Research</i> , 2018 , 43, 477-487	4.6	2
171	Neonatal hyperglycemia induces cell death in the rat brain. <i>Metabolic Brain Disease</i> , 2018 , 33, 333-342	3.9	3
170	Chronic Exposure to D-Alanine Generates Oxidative Stress and Alters Energy Metabolism in Cerebral Cortex and Cerebellum of Wistar Rats. <i>Molecular Neurobiology</i> , 2018 , 55, 5101-5110	6.2	7
169	Acute biochemical and physiological responses to swimming training series performed at intensities based on the 400-m front crawl speed. <i>Sport Sciences for Health</i> , 2018 , 14, 633-638	1.3	1
168	Evaluation of Oxidative Stress Parameters and Energy Metabolism in Cerebral Cortex of Rats Subjected to Sarcosine Administration. <i>Molecular Neurobiology</i> , 2017 , 54, 4496-4506	6.2	3
167	Voluntary Exercise Prevents Oxidative Stress in the Brain of Phenylketonuria Mice. <i>JIMD Reports</i> , 2016 , 27, 69-77	1.9	11
166	Brain zinc chelation by diethyldithiocarbamate increased the behavioral and mitochondrial damages in zebrafish subjected to hypoxia. <i>Scientific Reports</i> , 2016 , 6, 20279	4.9	11
165	Phenylalanine induces oxidative stress and decreases the viability of rat astrocytes: possible relevance for the pathophysiology of neurodegeneration in phenylketonuria. <i>Metabolic Brain Disease</i> , 2016 , 31, 529-37	3.9	20
164	Chemically induced acute model of sarcosinemia in wistar rats. <i>Metabolic Brain Disease</i> , 2016 , 31, 363-8	3.9	2
163	L-carnitine Prevents Oxidative Stress in the Brains of Rats Subjected to a Chemically Induced Chronic Model of MSUD. <i>Molecular Neurobiology</i> , 2016 , 53, 6007-6017	6.2	28
162	Urinary biomarkers of oxidative damage in Maple syrup urine disease: the L-carnitine role. <i>International Journal of Developmental Neuroscience</i> , 2015 , 42, 10-4	2.7	22
161	Neonatal hyperglycemia induces oxidative stress in the rat brain: the role of pentose phosphate pathway enzymes and NADPH oxidase. <i>Molecular and Cellular Biochemistry</i> , 2015 , 403, 159-67	4.2	16
160	L-Carnitine supplementation decreases DNA damage in treated MSUD patients. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2015 , 775, 43-7	3.3	23
159	Creatine and pyruvate prevent the alterations caused by tyrosine on parameters of oxidative stress and enzyme activities of phosphoryltransfer network in cerebral cortex of Wistar rats. <i>Molecular Neurobiology</i> , 2015 , 51, 1184-94	6.2	15
158	Acute exercise in treated phenylketonuria patients: Physical activity and biochemical response. <i>Molecular Genetics and Metabolism Reports</i> , 2015 , 5, 55-59	1.8	5
157	Investigation of inflammatory profile in MSUD patients: benefit of L-carnitine supplementation. <i>Metabolic Brain Disease</i> , 2015 , 30, 1167-74	3.9	21

156	Pipecolic acid induces oxidative stress in vitro in cerebral cortex of young rats and the protective role of lipoic acid. <i>Metabolic Brain Disease</i> , 2014 , 29, 175-83	3.9	20
155	Glutathione metabolism enzymes in brain and liver of hyperphenylalaninemic rats and the effect of lipoic acid treatment. <i>Metabolic Brain Disease</i> , 2014 , 29, 609-15	3.9	15
154	Prevention of DNA damage by L-carnitine induced by metabolites accumulated in maple syrup urine disease in human peripheral leukocytes in vitro. <i>Gene</i> , 2014 , 548, 294-8	3.8	16
153	Diabetic encephalopathy-related depression: experimental evidence that insulin and clonazepam restore antioxidant status in rat brain. <i>Cell Biochemistry and Function</i> , 2014 , 32, 711-9	4.2	11
152	Enzymatic scavengers in the epididymal fluid: comparison between pony and miniature breed stallions. <i>Animal Reproduction Science</i> , 2014 , 151, 164-8	2.1	5
151	Antioxidant treatment strategies for hyperphenylalaninemia. <i>Metabolic Brain Disease</i> , 2013 , 28, 541-50	3.9	12
150	Neurochemical evidence that the metabolites accumulating in 3-methylcrotonyl-CoA carboxylase deficiency induce oxidative damage in cerebral cortex of young rats. <i>Cellular and Molecular Neurobiology</i> , 2013 , 33, 137-46	4.6	12
149	Protein and lipid damage in maple syrup urine disease patients: l-carnitine effect. <i>International Journal of Developmental Neuroscience</i> , 2013 , 31, 21-4	2.7	36
148	Role of catalase and superoxide dismutase activities on oxidative stress in the brain of a phenylketonuria animal model and the effect of lipoic acid. <i>Cellular and Molecular Neurobiology</i> , 2013 , 33, 253-60	4.6	19
147	Experimental hyperprolinemia induces mild oxidative stress, metabolic changes, and tissue adaptation in rat liver. <i>Journal of Cellular Biochemistry</i> , 2012 , 113, 174-83	4.7	14
146	Pyruvate and creatine prevent oxidative stress and behavioral alterations caused by phenylalanine administration into hippocampus of rats. <i>Metabolic Brain Disease</i> , 2012 , 27, 79-89	3.9	20
145	Phenylpyruvic acid decreases glucose-6-phosphate dehydrogenase activity in rat brain. <i>Cellular and Molecular Neurobiology</i> , 2012 , 32, 1113-8	4.6	16
144	Effect of histidine administration to female rats during pregnancy and lactation on enzymes activity of phosphoryltransfer network in cerebral cortex and hippocampus of the offspring. <i>Metabolic Brain Disease</i> , 2012 , 27, 595-603	3.9	8
143	Exercício aeróbico agudo restaura a concentraçã de triptofano em cãebro de ratos com hiperfenilalaninemia. <i>Revista Brasileira De Medicina Do Esporte</i> , 2012 , 18, 338-340	0.5	2
142	Tyrosine impairs enzymes of energy metabolism in cerebral cortex of rats. <i>Molecular and Cellular Biochemistry</i> , 2012 , 364, 253-61	4.2	22
141	Administration of histidine to female rats induces changes in oxidative status in cortex and hippocampus of the offspring. <i>Neurochemical Research</i> , 2012 , 37, 1031-6	4.6	19
140	Dehydroepiandrosterone improves hepatic antioxidant reserve and stimulates Akt signaling in young and old rats. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2011 , 127, 331-6	5.1	14
139	Chronic hyperhomocysteinemia induces oxidative damage in the rat lung. <i>Molecular and Cellular Biochemistry</i> , 2011 , 358, 153-60	4.2	19

138	In vivo neuroprotective effect of L-carnitine against oxidative stress in maple syrup urine disease. <i>Metabolic Brain Disease</i> , 2011 , 26, 21-8	3.9	51
137	Tyrosine inhibits creatine kinase activity in cerebral cortex of young rats. <i>Metabolic Brain Disease</i> , 2011 , 26, 221-7	3.9	17
136	Regular exercise prevents oxidative stress in the brain of hyperphenylalaninemic rats. <i>Metabolic Brain Disease</i> , 2011 , 26, 291-7	3.9	21
135	Lipoic acid prevents oxidative stress in vitro and in vivo by an acute hyperphenylalaninemia chemically-induced in rat brain. <i>Journal of the Neurological Sciences</i> , 2010 , 292, 89-95	3.2	38
134	d-Serine administration provokes lipid oxidation and decreases the antioxidant defenses in rat striatum. <i>International Journal of Developmental Neuroscience</i> , 2010 , 28, 297-301	2.7	8
133	Experimental evidence that phenylalanine provokes oxidative stress in hippocampus and cerebral cortex of developing rats. <i>Cellular and Molecular Neurobiology</i> , 2010 , 30, 317-26	4.6	48
132	Neuroprotective role of lipoic acid against acute toxicity of N-acetylaspartic acid. <i>Molecular and Cellular Biochemistry</i> , 2010 , 344, 231-9	4.2	12
131	Acute administration of 5-oxoproline induces oxidative damage to lipids and proteins and impairs antioxidant defenses in cerebral cortex and cerebellum of young rats. <i>Metabolic Brain Disease</i> , 2010 , 25, 145-54	3.9	20
130	N-acetylaspartic acid impairs enzymatic antioxidant defenses and enhances hydrogen peroxide concentration in rat brain. <i>Metabolic Brain Disease</i> , 2010 , 25, 251-9	3.9	4
129	Evidence that 2-methylacetoacetate induces oxidative stress in rat brain. <i>Metabolic Brain Disease</i> , 2010 , 25, 261-7	3.9	7
128	Redox imbalance influence in the myocardial Akt activation in aged rats treated with DHEA. <i>Experimental Gerontology</i> , 2010 , 45, 957-63	4.5	21
127	Experimental evidence that ornithine and homocitrulline disrupt energy metabolism in brain of young rats. <i>Brain Research</i> , 2009 , 1291, 102-12	3.7	15
126	Effects of 1,4-butanediol administration on oxidative stress in rat brain: study of the neurotoxicity of gamma-hydroxybutyric acid in vivo. <i>Metabolic Brain Disease</i> , 2009 , 24, 271-82	3.9	28
125	Intracerebroventricular administration of N-acetylaspartic acid impairs antioxidant defenses and promotes protein oxidation in cerebral cortex of rats. <i>Metabolic Brain Disease</i> , 2009 , 24, 283-98	3.9	18
124	Tyrosine administration decreases glutathione and stimulates lipid and protein oxidation in rat cerebral cortex. <i>Metabolic Brain Disease</i> , 2009 , 24, 415-25	3.9	26
123	Glycine provokes lipid oxidative damage and reduces the antioxidant defenses in brain cortex of young rats. <i>Cellular and Molecular Neurobiology</i> , 2009 , 29, 253-61	4.6	18
122	Amino acids levels and lipid peroxidation in maple syrup urine disease patients. <i>Clinical Biochemistry</i> , 2009 , 42, 462-6	3.5	21
121	Hypermethioninemia provokes oxidative damage and histological changes in liver of rats. <i>Biochimie</i> , 2009 , 91, 961-8	4.6	21

120	Homocysteine induces oxidative stress, inflammatory infiltration, fibrosis and reduces glycogen/glycoprotein content in liver of rats. <i>International Journal of Developmental Neuroscience</i> , 2009 , 27, 337-44	2.7	56
119	Evidence that the major metabolites accumulating in hyperornithinemia-hyperammonemia-homocitrullinuria syndrome induce oxidative stress in brain of young rats. <i>International Journal of Developmental Neuroscience</i> , 2009 , 27, 635-41	2.7	9
118	Medium-chain fatty acids accumulating in MCAD deficiency elicit lipid and protein oxidative damage and decrease non-enzymatic antioxidant defenses in rat brain. <i>Neurochemistry International</i> , 2009 , 54, 519-25	4.4	27
117	Evidence that 3-hydroxyisobutyric acid inhibits key enzymes of energy metabolism in cerebral cortex of young rats. <i>International Journal of Developmental Neuroscience</i> , 2008 , 26, 293-9	2.7	9
116	Tyrosine promotes oxidative stress in cerebral cortex of young rats. <i>International Journal of Developmental Neuroscience</i> , 2008 , 26, 551-9	2.7	30
115	Induction of oxidative stress by the metabolites accumulating in isovaleric acidemia in brain cortex of young rats. <i>Free Radical Research</i> , 2008 , 42, 707-15	4	18
114	Sulfite increases lipoperoxidation and decreases the activity of catalase in brain of rats. <i>Metabolic Brain Disease</i> , 2008 , 23, 123-32	3.9	20
113	Oxidative stress in plasma from maple syrup urine disease patients during treatment. <i>Metabolic Brain Disease</i> , 2008 , 23, 71-80	3.9	36
112	Effects of cysteamine on oxidative status in cerebral cortex of rats. <i>Metabolic Brain Disease</i> , 2008 , 23, 81-93	3.9	14
111	Tryptophan administration induces oxidative stress in brain cortex of rats. <i>Metabolic Brain Disease</i> , 2008 , 23, 221-33	3.9	20
110	Influence of ketone bodies on oxidative stress parameters in brain of developing rats in vitro. <i>Metabolic Brain Disease</i> , 2008 , 23, 411-25	3.9	7
109	Inhibition of brain energy metabolism by the branched-chain amino acids accumulating in maple syrup urine disease. <i>Neurochemical Research</i> , 2008 , 33, 114-24	4.6	46
108	Antioxidant effect of cysteamine in brain cortex of young rats. <i>Neurochemical Research</i> , 2008 , 33, 737-44	4.6	34
107	Guanidinoacetate decreases antioxidant defenses and total protein sulfhydryl content in striatum of rats. <i>Neurochemical Research</i> , 2008 , 33, 1804-10	4.6	40
106	Evidence that 3-hydroxy-3-methylglutaric acid promotes lipid and protein oxidative damage and reduces the nonenzymatic antioxidant defenses in rat cerebral cortex. <i>Journal of Neuroscience Research</i> , 2008 , 86, 683-93	4.4	27
105	Oxidative stress induction by cis-4-decenoic acid: relevance for MCAD deficiency. <i>Free Radical Research</i> , 2007 , 41, 1261-72	4	17
104	5-Oxoproline reduces non-enzymatic antioxidant defenses in vitro in rat brain. <i>Metabolic Brain Disease</i> , 2007 , 22, 51-65	3.9	24
103	Energy metabolism is compromised in skeletal muscle of rats chronically-treated with glutaric acid. <i>Metabolic Brain Disease</i> , 2007 , 22, 111-23	3.9	12

102	Synaptic plasma membrane Na(+), K (+)-ATPase activity is significantly reduced by the alpha-keto acids accumulating in maple syrup urine disease in rat cerebral cortex. <i>Metabolic Brain Disease</i> , 2007 , 22, 77-88	3.9	6
101	Kynurenines impair energy metabolism in rat cerebral cortex. <i>Cellular and Molecular Neurobiology</i> , 2007 , 27, 147-60	4.6	26
100	Induction of oxidative stress by chronic and acute glutaric acid administration to rats. <i>Cellular and Molecular Neurobiology</i> , 2007 , 27, 423-38	4.6	46
99	Age and brain structural related effects of glutaric and 3-hydroxyglutaric acids on glutamate binding to plasma membranes during rat brain development. <i>Cellular and Molecular Neurobiology</i> , 2007 , 27, 805-18	4.6	21
98	Promotion of oxidative stress in kidney of rats loaded with cystine dimethyl ester. <i>Pediatric Nephrology</i> , 2007 , 22, 1121-8	3.2	14
97	In vitro evidence for an antioxidant role of 3-hydroxykynurenine and 3-hydroxyanthranilic acid in the brain. <i>Neurochemistry International</i> , 2007 , 50, 83-94	4.4	68
96	Gamma-hydroxybutyric acid induces oxidative stress in cerebral cortex of young rats. <i>Neurochemistry International</i> , 2007 , 50, 564-70	4.4	40
95	N-acetylaspartic acid promotes oxidative stress in cerebral cortex of rats. <i>International Journal of Developmental Neuroscience</i> , 2007 , 25, 317-24	2.7	16
94	Erythrocyte glutathione peroxidase activity and plasma selenium concentration are reduced in maple syrup urine disease patients during treatment. <i>International Journal of Developmental Neuroscience</i> , 2007 , 25, 335-8	2.7	21
93	Evidence for a synergistic action of glutaric and 3-hydroxyglutaric acids disturbing rat brain energy metabolism. <i>International Journal of Developmental Neuroscience</i> , 2007 , 25, 391-8	2.7	33
92	Inhibition of creatine kinase activity from rat cerebral cortex by 3-hydroxykynurenine. <i>Brain Research</i> , 2006 , 1124, 188-96	3.7	5
91	Inhibition of creatine kinase activity by cystine in the kidney of young rats. <i>Pediatric Research</i> , 2006 , 60, 190-5	3.2	4
90	Na+, K+ ATPase activity is markedly reduced by cis-4-decenoic acid in synaptic plasma membranes from cerebral cortex of rats. <i>Experimental Neurology</i> , 2006 , 197, 143-9	5.7	13
89	Promotion of oxidative stress by L-tryptophan in cerebral cortex of rats. <i>Neurochemistry International</i> , 2006 , 49, 87-93	4.4	27
88	Differential inhibitory effects of methylmalonic acid on respiratory chain complex activities in rat tissues. <i>International Journal of Developmental Neuroscience</i> , 2006 , 24, 45-52	2.7	42
87	Evidence that quinolinic acid severely impairs energy metabolism through activation of NMDA receptors in striatum from developing rats. <i>Journal of Neurochemistry</i> , 2006 , 99, 1531-42	6	49
86	A chemically-induced acute model of maple syrup urine disease in rats for neurochemical studies. <i>Journal of Neuroscience Methods</i> , 2006 , 155, 224-30	3	27
85	Inhibition of the electron transport chain and creatine kinase activity by ethylmalonic acid in human skeletal muscle. <i>Metabolic Brain Disease</i> , 2006 , 21, 11-9	3.9	19

84	Citrulline and ammonia accumulating in citrullinemia reduces antioxidant capacity of rat brain in vitro. <i>Metabolic Brain Disease</i> , 2006 , 21, 63-74	3.9	8
83	Evidence that oxidative stress is increased in plasma from patients with maple syrup urine disease. <i>Metabolic Brain Disease</i> , 2006 , 21, 279-86	3.9	71
82	Investigation of oxidative stress parameters in treated phenylketonuric patients. <i>Metabolic Brain Disease</i> , 2006 , 21, 287-96	3.9	53
81	Evaluation of the mechanisms involved in leucine-induced oxidative damage in cerebral cortex of young rats. <i>Free Radical Research</i> , 2005 , 39, 71-9	4	46
80	Cysteamine prevents and reverses the inhibition of creatine kinase activity caused by cystine in rat brain cortex. <i>Neurochemistry International</i> , 2005 , 46, 391-7	4.4	15
79	Protective effect of antioxidants on brain oxidative damage caused by proline administration. <i>Neuroscience Research</i> , 2005 , 52, 69-74	2.9	16
78	The effects of the interactions between amino acids on pyruvate kinase activity from the brain cortex of young rats. <i>International Journal of Developmental Neuroscience</i> , 2005 , 23, 509-14	2.7	16
77	Glutaric acid moderately compromises energy metabolism in rat brain. <i>International Journal of Developmental Neuroscience</i> , 2005 , 23, 687-93	2.7	24
76	Quinolinic acid reduces the antioxidant defenses in cerebral cortex of young rats. <i>International Journal of Developmental Neuroscience</i> , 2005 , 23, 695-701	2.7	40
75	Oxidative stress in patients with phenylketonuria. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2005 , 1740, 68-73	6.9	78
74	Mitochondrial energy metabolism is markedly impaired by D-2-hydroxyglutaric acid in rat tissues. <i>Molecular Genetics and Metabolism</i> , 2005 , 86, 188-99	3.7	79
73	Benzophenones from <i>Hypericum carinatum</i> . <i>Journal of Natural Products</i> , 2005 , 68, 784-6	4.9	41
72	alpha-keto acids accumulating in maple syrup urine disease stimulate lipid peroxidation and reduce antioxidant defences in cerebral cortex from young rats. <i>Metabolic Brain Disease</i> , 2005 , 20, 155-67	3.9	56
71	Inhibition of energy metabolism by 2-methylacetoacetate and 2-methyl-3-hydroxybutyrate in cerebral cortex of developing rats. <i>Journal of Inherited Metabolic Disease</i> , 2005 , 28, 501-15	5.4	16
70	Promotion of oxidative stress by 3-hydroxyglutaric acid in rat striatum. <i>Journal of Inherited Metabolic Disease</i> , 2005 , 28, 57-67	5.4	47
69	Inhibition of pyruvate kinase activity by cystine in brain cortex of rats. <i>Brain Research</i> , 2004 , 1012, 93-100, 7	3.7	17
68	Inhibition of energy metabolism in cerebral cortex of young rats by the medium-chain fatty acids accumulating in MCAD deficiency. <i>Brain Research</i> , 2004 , 1030, 141-51	3.7	30
67	The role of oxidative damage in the neuropathology of organic acidurias: insights from animal studies. <i>Journal of Inherited Metabolic Disease</i> , 2004 , 27, 427-48	5.4	132

66	Monosialoganglioside increases catalase activity in cerebral cortex of rats. <i>Free Radical Research</i> , 2004 , 38, 495-500	4	16
65	Tryptophan reduces creatine kinase activity in the brain cortex of rats. <i>International Journal of Developmental Neuroscience</i> , 2004 , 22, 95-101	2.7	11
64	Effects of histidine and imidazolelactic acid on various parameters of the oxidative stress in cerebral cortex of young rats. <i>International Journal of Developmental Neuroscience</i> , 2004 , 22, 67-72	2.7	9
63	Inhibition of creatine kinase activity from rat cerebral cortex by D-2-hydroxyglutaric acid in vitro. <i>Neurochemistry International</i> , 2004 , 44, 45-52	4.4	39
62	Alanine prevents the inhibition of pyruvate kinase activity caused by tryptophan in cerebral cortex of rats. <i>Metabolic Brain Disease</i> , 2003 , 18, 129-37	3.9	20
61	In vitro effect of homocysteine on some parameters of oxidative stress in rat hippocampus. <i>Metabolic Brain Disease</i> , 2003 , 18, 147-54	3.9	78
60	Effect of proline on creatine kinase activity in rat brain. <i>Metabolic Brain Disease</i> , 2003 , 18, 169-77	3.9	1
59	Proline reduces creatine kinase activity in the brain cortex of rats. <i>Neurochemical Research</i> , 2003 , 28, 1175-80	4.6	19
58	Evidence that antioxidants prevent the inhibition of Na ⁺ ,K ⁽⁺⁾ -ATPase activity induced by octanoic acid in rat cerebral cortex in vitro. <i>Neurochemical Research</i> , 2003 , 28, 1255-63	4.6	16
57	Inhibition of mitochondrial creatine kinase activity by D-2-hydroxyglutaric acid in cerebellum of young rats. <i>Neurochemical Research</i> , 2003 , 28, 1329-37	4.6	4
56	Effects of L-2-hydroxyglutaric acid on various parameters of the glutamatergic system in cerebral cortex of rats. <i>Metabolic Brain Disease</i> , 2003 , 18, 233-43	3.9	13
55	Effect of leucine administration on creatine kinase activity in rat brain. <i>Metabolic Brain Disease</i> , 2003 , 18, 17-25	3.9	21
54	Alanine prevents the in vitro inhibition of glycolysis caused by phenylalanine in brain cortex of rats. <i>Metabolic Brain Disease</i> , 2003 , 18, 87-94	3.9	12
53	Creatine kinase activity from rat brain is inhibited by branched-chain amino acids in vitro. <i>Neurochemical Research</i> , 2003 , 28, 675-9	4.6	33
52	In vivo and in vitro effects of proline on some parameters of oxidative stress in rat brain. <i>Brain Research</i> , 2003 , 991, 180-6	3.7	32
51	Glutaric acid induces oxidative stress in brain of young rats. <i>Brain Research</i> , 2003 , 964, 153-8	3.7	72
50	Characterization of the inhibition of pyruvate kinase caused by phenylalanine and phenylpyruvate in rat brain cortex. <i>Brain Research</i> , 2003 , 968, 199-205	3.7	31
49	Ascorbic acid prevents water maze behavioral deficits caused by early postnatal methylmalonic acid administration in the rat. <i>Brain Research</i> , 2003 , 976, 234-42	3.7	25

48	Induction of oxidative stress by L-2-hydroxyglutaric acid in rat brain. <i>Journal of Neuroscience Research</i> , 2003 , 74, 103-10	4.4	50
47	D-2-hydroxyglutaric acid induces oxidative stress in cerebral cortex of young rats. <i>European Journal of Neuroscience</i> , 2003 , 17, 2017-22	3.5	85
46	Evaluation of the effect of chronic administration of drugs on rat behavior in the water maze task. <i>Brain Research Protocols</i> , 2003 , 12, 109-15		8
45	Inhibition of brain energy metabolism by the alpha-keto acids accumulating in maple syrup urine disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2003 , 1639, 232-8	6.9	67
44	Ammonia potentiates methylmalonic acid-induced convulsions and TBARS production. <i>Experimental Neurology</i> , 2003 , 182, 455-60	5.7	23
43	Hyperphenylalaninemia reduces creatine kinase activity in the cerebral cortex of rats. <i>International Journal of Developmental Neuroscience</i> , 2003 , 21, 111-6	2.7	23
42	Proline induces oxidative stress in cerebral cortex of rats. <i>International Journal of Developmental Neuroscience</i> , 2003 , 21, 105-10	2.7	47
41	Kinetic studies on the inhibition of creatine kinase activity by branched-chain alpha-amino acids in the brain cortex of rats. <i>International Journal of Developmental Neuroscience</i> , 2003 , 21, 145-51	2.7	10
40	Induction of oxidative stress in rat brain by the metabolites accumulating in maple syrup urine disease. <i>International Journal of Developmental Neuroscience</i> , 2003 , 21, 327-32	2.7	68
39	GM1 ganglioside attenuates convulsions and thiobarbituric acid reactive substances production induced by the intrastriatal injection of methylmalonic acid. <i>International Journal of Biochemistry and Cell Biology</i> , 2003 , 35, 465-73	5.6	46
38	Ascorbic acid prevents cognitive deficits caused by chronic administration of propionic acid to rats in the water maze. <i>Pharmacology Biochemistry and Behavior</i> , 2002 , 73, 623-9	3.9	54
37	Stimulation of lipid peroxidation in vitro in rat brain by the metabolites accumulating in maple syrup urine disease. <i>Metabolic Brain Disease</i> , 2002 , 17, 47-54	3.9	56
36	Inhibition of the mitochondrial respiratory chain by phenylalanine in rat cerebral cortex. <i>Neurochemical Research</i> , 2002 , 27, 353-7	4.6	33
35	Inhibition of the mitochondrial respiratory chain by alanine in rat cerebral cortex. <i>Metabolic Brain Disease</i> , 2002 , 17, 123-30	3.9	4
34	Alanine prevents the reduction of pyruvate kinase activity in brain cortex of rats subjected to chemically induced hyperphenylalaninemia. <i>Neurochemical Research</i> , 2002 , 27, 947-52	4.6	14
33	Arginine administration reduces catalase activity in midbrain of rats. <i>NeuroReport</i> , 2002 , 13, 1301-4	1.7	12
32	Inhibition of cytochrome c oxidase activity in rat cerebral cortex and human skeletal muscle by D-2-hydroxyglutaric acid in vitro. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2002 , 1586, 81-91	6.9	69
31	Experimental hyperphenylalaninemia provokes oxidative stress in rat brain. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2002 , 1586, 344-52	6.9	49

30	Inhibition of the mitochondrial respiratory chain complex activities in rat cerebral cortex by methylmalonic acid. <i>Neurochemistry International</i> , 2002 , 40, 593-601	4.4	93
29	L-pyroglutamic acid inhibits energy production and lipid synthesis in cerebral cortex of young rats in vitro. <i>Neurochemical Research</i> , 2001 , 26, 1277-83	4.6	17
28	Nitric oxide synthase inhibition by L-NAME prevents the decrease of Na ⁺ ,K ⁺ -ATPase activity in midbrain of rats subjected to arginine administration. <i>Neurochemical Research</i> , 2001 , 26, 515-20	4.6	38
27	Effects of methylmalonic and propionic acids on glutamate uptake by synaptosomes and synaptic vesicles and on glutamate release by synaptosomes from cerebral cortex of rats. <i>Brain Research</i> , 2001 , 920, 194-201	3.7	21
26	In vitro stimulation of oxidative stress in cerebral cortex of rats by the guanidino compounds accumulating in hyperargininemia. <i>Brain Research</i> , 2001 , 923, 50-7	3.7	26
25	Reduced Na ⁽⁺⁾ , K ⁽⁺⁾ -ATPase activity in erythrocyte membranes from patients with phenylketonuria. <i>Pediatric Research</i> , 2001 , 50, 56-60	3.2	9
24	Inhibition of glutamate uptake into synaptic vesicles from rat brain by 3-nitropropionic acid in vitro. <i>Experimental Neurology</i> , 2001 , 172, 250-4	5.7	17
23	Reduction of large neutral amino acid levels in plasma and brain of hyperleucinemic rats. <i>Neurochemistry International</i> , 2001 , 38, 529-37	4.4	58
22	Inhibition of in vitro CO ₂ production and lipid synthesis by 2-hydroxybutyric acid in rat brain. <i>Brazilian Journal of Medical and Biological Research</i> , 2001 , 34, 627-31	2.8	8
21	Chronic postnatal administration of methylmalonic acid provokes a decrease of myelin content and ganglioside N-acetylneuraminic acid concentration in cerebrum of young rats. <i>Brazilian Journal of Medical and Biological Research</i> , 2001 , 34, 227-31	2.8	11
20	Propionic and L-methylmalonic acids induce oxidative stress in brain of young rats. <i>NeuroReport</i> , 2000 , 11, 541-4	1.7	72
19	Quinolinic acid inhibits glutamate uptake into synaptic vesicles from rat brain. <i>NeuroReport</i> , 2000 , 11, 249-53	1.7	81
18	Effect of phenylalanine and p-chlorophenylalanine on Na ⁺ , K ⁺ -ATPase activity in the synaptic plasma membrane from the cerebral cortex of rats. <i>Metabolic Brain Disease</i> , 2000 , 15, 105-14	3.9	6
17	Platelet Na ⁺ , K ⁺ -ATPase activity as a possible peripheral marker for the neurotoxic effects of phenylalanine in phenylketonuria. <i>Metabolic Brain Disease</i> , 2000 , 15, 115-21	3.9	3
16	Inhibition of energy production in vitro by glutaric acid in cerebral cortex of young rats. <i>Metabolic Brain Disease</i> , 2000 , 15, 123-31	3.9	28
15	Effect of collection, transport, processing and storage of blood specimens on the activity of lysosomal enzymes in plasma and leukocytes. <i>Brazilian Journal of Medical and Biological Research</i> , 2000 , 33, 1003-13	2.8	7
14	Inhibition of glutamate uptake into synaptic vesicles of rat brain by the metabolites accumulating in maple syrup urine disease. <i>Journal of the Neurological Sciences</i> , 2000 , 181, 44-9	3.2	57
13	Inhibition of rat brain lipid synthesis in vitro by 4-hydroxybutyric acid. <i>Metabolic Brain Disease</i> , 1999 , 14, 157-64	3.9	10

12	Proline administration decreases Na ⁺ ,K ⁺ -ATPase activity in the synaptic plasma membrane from cerebral cortex of rats. <i>Metabolic Brain Disease</i> , 1999 , 14, 265-72	3.9	16
11	2-Hydroxybutyrate and 4-hydroxybutyrate inhibit CO ₂ formation from labeled substrates by rat cerebral cortex. <i>Biochemical Society Transactions</i> , 1995 , 23, 228S	5.1	12
10	Inhibition of citrate oxidation in vitro by 2-hydroxybutyrate and 4-hydroxybutyrate in cerebral cortex of young rats. <i>Biochemical Society Transactions</i> , 1995 , 23, 229S	5.1	3
9	Possible high frequency of tetrahydrobiopterin deficiency in south Brazil. <i>Journal of Inherited Metabolic Disease</i> , 1994 , 17, 223-9	5.4	6
8	Inhibition of succinate dehydrogenase and beta-hydroxybutyrate dehydrogenase activities by methylmalonate in brain and liver of developing rats. <i>Journal of Inherited Metabolic Disease</i> , 1993 , 16, 147-53	5.4	53
7	Effects of methylmalonate and propionate on uptake of glucose and ketone bodies in vitro by brain of developing rats. <i>Biochemical Medicine and Metabolic Biology</i> , 1991 , 45, 56-64		24
6	Seven-year experience of a reference laboratory for detection of inborn errors of metabolism in Brazil. <i>Journal of Inherited Metabolic Disease</i> , 1991 , 14, 400-2	5.4	4
5	Effect of phenylalanine, p-chlorophenylalanine and alpha-methylphenylalanine on glucose uptake in vitro by the brain of young rats. <i>Biochemical Society Transactions</i> , 1990 , 18, 419	5.1	9
4	Influence of methylmalonate on the uptake of ketone bodies in vitro by the brain of young rats. <i>Biochemical Society Transactions</i> , 1990 , 18, 421-2	5.1	1
3	Reduced locomotor activity of rats made histidinemic by injection of histidine. <i>Journal of Nutrition</i> , 1989 , 119, 1223-7	4.1	10
2	Inborn errors of metabolism. Sensitivity of screening tests in high risk patients. <i>Clinical Pediatrics</i> , 1989 , 28, 494-7	1.2	2
1	An improved specific laboratory test for homocystinuria. <i>Clinica Chimica Acta</i> , 1982 , 125, 367-9	6.2	6