Timothy Bagguley

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5339174/publications.pdf

Version: 2024-02-01

		1306789	1058022	
18	252	7	14	
papers	citations	h-index	g-index	
20	20	20	F20	
20	20	20	530	
all docs	docs citations	times ranked	citing authors	

#	Article	IF	CITATIONS
1	Health-related quality of life in lower-risk MDS patients compared with age- and sex-matched reference populations: a European LeukemiaNet study. Leukemia, 2018, 32, 1380-1392.	3.3	66
2	Cohort Profile: The Haematological Malignancy Research Network (HMRN): a UK population-based patient cohort. International Journal of Epidemiology, 2018, 47, 700-700g.	0.9	47
3	Multiple myeloma: routes to diagnosis, clinical characteristics and survival – findings from a <scp>UK</scp> populationâ€based study. British Journal of Haematology, 2017, 177, 67-71.	1.2	39
4	Labile plasma iron levels predict survival in patients with lower-risk myelodysplastic syndromes. Haematologica, 2018, 103, 69-79.	1.7	35
5	Toxic iron species in lower-risk myelodysplastic syndrome patients: course of disease and effects on outcome. Leukemia, 2021, 35, 1745-1750.	3.3	15
6	Frontline management of nodular lymphocyte predominant Hodgkin lymphoma – a retrospective UK multicentre study. British Journal of Haematology, 2019, 186, e214-e217.	1.2	9
7	PCN351 - HEALTH-STATE UTILITY VALUES IN DIFFUSE LARGE B-CELL LYMPHOMA. Value in Health, 2018, 21, S74.	0.1	8
8	Hodgkin lymphoma detection and survival: findings from the Haematological Malignancy Research Network. BJGP Open, 2019, 3, bjgpopen19X101668.	0.9	8
9	Cohort Profile Update: The Haematological Malignancy Research Network (HMRN) UK population-based cohorts. International Journal of Epidemiology, 2022, 51, e87-e94.	0.9	7
10	Health impact of monoclonal gammopathy of undetermined significance (MGUS) and monoclonal B-cell lymphocytosis (MBL): findings from a UK population-based cohort. BMJ Open, 2021, 11, e041296.	0.8	5
11	Skeletal-Related Events In Myeloma: A Population-Based Study. Blood, 2013, 122, 3158-3158.	0.6	5
12	Labile Plasma Iron (LPI) Is a Clinical Indicator of Overt Iron Overload in Patients with Lower-Risk Myelodysplastic Syndromes (MDS) from the European Leukemianet MDS Registry. Blood, 2015, 126, 2865-2865.	0.6	3
13	Elevated Labile Plasma Iron Levels (LPI) and Increased Oxidative Stress Are Associated with Red Blood Cell Transfusions in Patients with Lower-Risk Myelodysplastic Syndromes (MDS) Subtitle: from the European Leukemianet MDS Registry. Blood, 2016, 128, 4327-4327.	0.6	1
14	212 TRANSFUSIONS AND PRESENCE OF RINGSIDEROBLASTS INFLUENCE HEPCIDIN AND NTBI LEVELS IN PATIENTS WITH LOWER-RISK MYELODYSPLASTIC SYNDROMES (MDS) - A REPORT FROM THE EUROPEAN LEUKEMIANET MDS REGISTRY. Leukemia Research, 2015, 39, S106-S107.	0.4	0
15	Hepcidin and GDF15 Levels during the First 2 Years Follow-up in Patients with Low and Int-1 Risk Myelodysplastic Syndromes (MDS) from the European Leukemianet MDS Registry. Blood, 2014, 124, 3267-3267.	0.6	O
16	Novel Targeted Therapies and Their Impact on Survival from Multiple Myeloma (MM) and Chronic Lymphocytic Leukaemia (CLL) in the Real World. Blood, 2016, 128, 3546-3546.	0.6	0
17	Elevated Labile Plasma Iron (LPI) Levels in Patients with Lower-Risk Myelodysplastic Syndromes (MDS) Are Associated with Decreased Quality of Life and Reduced Survival. Blood, 2018, 132, 4392-4392.	0.6	0
18	Matching Adjusted Indirect Comparison to Assess the Relative Efficacy of Idelalisib in Double-Refractory Follicular Lymphoma. Blood, 2018, 132, 5842-5842.	0.6	0