

Stefanie Hessel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5340425/publications.pdf>

Version: 2024-02-01

35
papers

561
citations

567281

15
h-index

677142

22
g-index

35
all docs

35
docs citations

35
times ranked

619
citing authors

#	ARTICLE	IF	CITATIONS
1	Organic Cation Transporter I and Na ⁺ /taurocholate Co ⁺ Transporting Polypeptide are Involved in Retrorsine ⁺ and Senecionine ⁺ Induced Hepatotoxicity in HepaRG cells. <i>Molecular Nutrition and Food Research</i> , 2022, 66, e2100800.	3.3	4
2	Identification of microRNAs Implicated in Modulating Senecionine-Induced Liver Toxicity in HepaRG Cells. <i>Foods</i> , 2022, 11, 532.	4.3	2
3	The chemical structure impairs the intensity of genotoxic effects promoted by 1,2-unsaturated pyrrolizidine alkaloids in vitro. <i>Food and Chemical Toxicology</i> , 2022, 164, 113049.	3.6	5
4	Mixture effects of food-relevant polycyclic aromatic hydrocarbons on the activation of nuclear receptors and gene expression, benzo[a]pyrene metabolite profile and DNA damage in HepaRG cells. <i>Food and Chemical Toxicology</i> , 2021, 147, 111884.	3.6	11
5	Pyrrolizidine Alkaloids Disturb Bile Acid Homeostasis in the Human Hepatoma Cell Line HepaRG. <i>Foods</i> , 2021, 10, 161.	4.3	6
6	Active Transport of Hepatotoxic Pyrrolizidine Alkaloids in HepaRG Cells. <i>International Journal of Molecular Sciences</i> , 2021, 22, 3821.	4.1	8
7	The Food Contaminants Pyrrolizidine Alkaloids Disturb Bile Acid Homeostasis Structure-Dependently in the Human Hepatoma Cell Line HepaRG. <i>Foods</i> , 2021, 10, 1114.	4.3	4
8	Pyrrolizidine alkaloid-induced transcriptomic changes in rat lungs in a 28-day subacute feeding study. <i>Archives of Toxicology</i> , 2021, 95, 2785-2796.	4.2	5
9	Pyrrolizidine Alkaloids Induce Cell Death in Human HepaRG Cells in a Structure-Dependent Manner. <i>International Journal of Molecular Sciences</i> , 2021, 22, 202.	4.1	15
10	Polycyclic Aromatic Hydrocarbons Activate the Aryl Hydrocarbon Receptor and the Constitutive Androstane Receptor to Regulate Xenobiotic Metabolism in Human Liver Cells. <i>International Journal of Molecular Sciences</i> , 2021, 22, 372.	4.1	26
11	The pyrrolizidine alkaloid senecionine induces CYP-dependent destruction of sinusoidal endothelial cells and cholestasis in mice. <i>Archives of Toxicology</i> , 2020, 94, 219-229.	4.2	33
12	Hepatotoxic pyrrolizidine alkaloids induce DNA damage response in rat liver in a 28-day feeding study. <i>Archives of Toxicology</i> , 2020, 94, 1739-1751.	4.2	25
13	Comparison of long-term versus short-term effects of okadaic acid on the apoptotic status of human HepaRG cells. <i>Chemico-Biological Interactions</i> , 2020, 317, 108937.	4.0	9
14	Okadaic acid activates Wnt/ β -catenin-signaling in human HepaRG cells. <i>Archives of Toxicology</i> , 2019, 93, 1927-1939.	4.2	10
15	Human CYP3A4-mediated toxification of the pyrrolizidine alkaloid lasiocarpine. <i>Food and Chemical Toxicology</i> , 2019, 130, 79-88.	3.6	35
16	Sensitization of Human Liver Cells Toward Fas ⁺ Mediated Apoptosis by the Metabolically Activated Pyrrolizidine Alkaloid Lasiocarpine. <i>Molecular Nutrition and Food Research</i> , 2019, 63, e1801206.	3.3	12
17	Metabolism of the lipophilic phycotoxin 13-Desmethylspirolide C using human and rat in vitro liver models. <i>Toxicology Letters</i> , 2019, 307, 17-25.	0.8	0
18	Combined effects of okadaic acid and pectenotoxin-2, 13-desmethylspirolide C or yessotoxin in human intestinal Caco-2 ⁺ cells. <i>Chemosphere</i> , 2019, 228, 139-148.	8.2	12

#	ARTICLE	IF	CITATIONS
19	The marine biotoxin okadaic acid affects intestinal tight junction proteins in human intestinal cells. <i>Toxicology in Vitro</i> , 2019, 58, 150-160.	2.4	19
20	In vitro characterization of hepatic toxicity of Alternaria toxins. <i>Mycotoxin Research</i> , 2019, 35, 157-168.	2.3	16
21	Pyrrrolizidine alkaloid-induced alterations of prostanoid synthesis in human endothelial cells. <i>Chemico-Biological Interactions</i> , 2019, 298, 104-111.	4.0	6
22	Structure-dependent induction of apoptosis by hepatotoxic pyrrolizidine alkaloids in the human hepatoma cell line HepaRG: Single versus repeated exposure. <i>Food and Chemical Toxicology</i> , 2018, 114, 215-226.	3.6	28
23	PXR: Structure-specific activation by hepatotoxic pyrrolizidine alkaloids. <i>Chemico-Biological Interactions</i> , 2018, 288, 38-48.	4.0	15
24	Metabolism of okadaic acid by NADPH-dependent enzymes present in human or rat liver S9 fractions results in different toxic effects. <i>Toxicology in Vitro</i> , 2017, 42, 161-170.	2.4	15
25	Metabolism of the Marine Phycotoxin PTX-2 and Its Effects on Hepatic Xenobiotic Metabolism: Activation of Nuclear Receptors and Modulation of the Phase I Cytochrome P450. <i>Toxins</i> , 2017, 9, 212.	3.4	8
26	Differences in metabolism of the marine biotoxin okadaic acid by human and rat cytochrome P450 monooxygenases. <i>Archives of Toxicology</i> , 2016, 90, 2025-2036.	4.2	18
27	Disturbance of gene expression in primary human hepatocytes by hepatotoxic pyrrolizidine alkaloids: A whole genome transcriptome analysis. <i>Toxicology in Vitro</i> , 2015, 29, 1669-1682.	2.4	31
28	Utility of an appropriate reporter assay: Heliotrine interferes with GAL4/upstream activation sequence-driven reporter gene systems. <i>Analytical Biochemistry</i> , 2015, 487, 45-48.	2.4	10
29	Dose-dependent induction of signaling pathways by the flavonoid quercetin in human primary hepatocytes: A transcriptomic study. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 1117-1129.	3.3	10
30	Structure-activity relationship in the passage of different pyrrolizidine alkaloids through the gastrointestinal barrier: ABCB1 excretes heliotrine and echimidine. <i>Molecular Nutrition and Food Research</i> , 2014, 58, 995-1004.	3.3	24
31	Active elimination of the marine biotoxin okadaic acid by P-glycoprotein through an in vitro gastrointestinal barrier. <i>Toxicology Letters</i> , 2014, 225, 311-317.	0.8	23
32	Polycyclic aromatic hydrocarbons stimulate human CYP3A4 promoter activity via PXR. <i>Toxicology Letters</i> , 2013, 222, 180-188.	0.8	79
33	Multidrug resistance-associated proteins are involved in the transport of the glutathione conjugates of the ultimate carcinogen of benzo[a]pyrene in human Caco-2 cells. <i>Archives of Toxicology</i> , 2013, 87, 269-280.	4.2	16
34	Analysis of GSH Conjugates of Bay- and Fjord-Region Dihydrodiol Epoxides of Benzo[a]pyrene and Dibenz[a,l]pyrene and their Transport in Enterocyte-like Caco-2 Cells. <i>Polycyclic Aromatic Compounds</i> , 2012, 32, 221-237.	2.6	5
35	All-trans retinoic acid enhances the transport of phase II metabolites of benzo[a]pyrene by inducing the Breast Cancer Resistance Protein expression in Caco-2 cells. <i>Toxicology Letters</i> , 2010, 197, 151-155.	0.8	16