

Yu Hung Chang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/534117/publications.pdf>

Version: 2024-02-01

22
papers

631
citations

759233

12
h-index

677142

22
g-index

22
all docs

22
docs citations

22
times ranked

1355
citing authors

#	ARTICLE	IF	CITATIONS
1	Pioglitazone and statins lower incidence of Parkinson disease in patients with diabetes mellitus. <i>European Journal of Neurology</i> , 2021, 28, 430-437.	3.3	19
2	<p>Common and Unique Factors and the Bidirectional Relationship Between Chronic Kidney Disease and Nonalcoholic Fatty Liver in Type 2 Diabetes Patients</p>. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020, Volume 13, 1203-1214.	2.4	12
3	Elevated serum cytokeratin-18 concentration in patients with type 2 diabetes mellitus and non-alcoholic fatty liver disease. <i>Annals of Clinical Biochemistry</i> , 2019, 56, 141-147.	1.6	12
4	Renal function preservation with pioglitazone or with basal insulin as an add-on therapy for patients with type 2 diabetes mellitus. <i>Acta Diabetologica</i> , 2017, 54, 561-568.	2.5	10
5	The association of obstructive sleep apnea and renal outcomesâ€”a systematic review and meta-analysis. <i>BMC Nephrology</i> , 2017, 18, 313.	1.8	30
6	Serum uric acid level as an indicator for CKD regression and progression in patients with type 2 diabetes mellitusâ€”a 4.6â€”year cohort study. <i>Diabetes/Metabolism Research and Reviews</i> , 2016, 32, 557-564.	4.0	51
7	Benefits of Rosuvastatin in Cardiovascular Protection Remain Unclear After HOPE-3. <i>Review of Diabetic Studies</i> , 2016, 13, 212-214.	1.3	2
8	Changes in high-density-lipoprotein cholesterol upon statin treatment in type 2 diabetic patients: A meta-analysis. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014, 24, 1067-1073.	2.6	2
9	High-density lipoprotein cholesterol and the risk of nephropathy in type 2 diabetic patients. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2013, 23, 751-757.	2.6	44
10	Developing and Psychometric Testing of a Short-Form Problem Areas in Diabetes Scale in Chinese Patients. <i>The Journal of Nursing Research: JNR</i> , 2013, 21, 212-217.	1.7	19
11	Reassessing the Benefits of Statins in the Prevention of Cardiovascular Disease in Diabetic Patients - A Systematic Review and Meta-Analysis. <i>Review of Diabetic Studies</i> , 2013, 10, 157-170.	1.3	14
12	Paradoxical Negative HDL Cholesterol Response to Atorvastatin and Simvastatin Treatment in Chinese Type 2 Diabetic Patients. <i>Review of Diabetic Studies</i> , 2013, 10, 213-222.	1.3	6
13	Serum osteoprotegerin and tumor necrosis factor related apoptosis inducing-ligand (TRAIL) are elevated in type 2 diabetic patients with albuminuria and serum osteoprotegerin is independently associated with the severity of diabetic nephropathy. <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 1064-1069.	3.4	42
14	Visfatin in overweight/obesity, type 2 diabetes mellitus, insulin resistance, metabolic syndrome and cardiovascular diseases: a metaâ€”analysis and systemic review. <i>Diabetes/Metabolism Research and Reviews</i> , 2011, 27, 515-527.	4.0	276
15	One-Year Efficacy and Safety of the Telehealth System in Poorly Controlled Type 2 Diabetic Patients Receiving Insulin Therapy. <i>Telemedicine Journal and E-Health</i> , 2011, 17, 683-687.	2.8	17
16	Albuminuria Is the Stronger Risk Factor for Peripheral Arterial Disease than eGFR Decline in a Type 2 Diabetic Taiwanese Population. <i>Kidney and Blood Pressure Research</i> , 2010, 33, 352-359.	2.0	9
17	Fulminant Type 1 Diabetes Mellitus Leading to Fetal Loss in a Pregnant Chinese Woman. <i>Kaohsiung Journal of Medical Sciences</i> , 2010, 26, 316-320.	1.9	4
18	Diabetic Muscle Infarction in Association with Terlipressin Therapy: A Case Report. <i>Kaohsiung Journal of Medical Sciences</i> , 2009, 25, 25-28.	1.9	3

#	ARTICLE	IF	CITATIONS
19	Elevated serum retinol-binding protein 4 concentrations are associated with renal dysfunction and uric acid in type 2 diabetic patients. <i>Diabetes/Metabolism Research and Reviews</i> , 2008, 24, 629-634.	4.0	22
20	High-sensitivity C-reactive protein and silent myocardial ischemia in Chinese with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2008, 57, 1533-1538.	3.4	11
21	Thiazolidinedione addition reduces the serum retinol-binding protein 4 in type 2 diabetic patients treated with metformin and sulfonylurea. <i>Translational Research</i> , 2008, 151, 309-314.	5.0	19
22	Insulinoma-associated Transient Hypothalamus-Pituitary-Adrenal Axis Impairment and Amelioration by Steroid Therapy and Surgical Intervention: A Case Report. <i>Kaohsiung Journal of Medical Sciences</i> , 2007, 23, 526-530.	1.9	7