

Jose Luis Rosa

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5344618/jose-luis-rosa-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

90
papers

7,270
citations

34
h-index

85
g-index

93
ext. papers

8,294
ext. citations

5.2
avg, IF

5
L-index

#	Paper	IF	Citations
90	NRF2 function in osteocytes is required for bone homeostasis and drives osteocytic gene expression. <i>Redox Biology</i> , 2021 , 40, 101845	11.3	10
89	HERC Ubiquitin Ligases in Cancer. <i>Cancers</i> , 2020 , 12,	6.6	8
88	The ubiquitin ligase HERC1 regulates cell migration via RAF-dependent regulation of MKK3/p38 signaling. <i>Scientific Reports</i> , 2020 , 10, 824	4.9	9
87	Regulation of the MDM2-p53 pathway by the ubiquitin ligase HERC2. <i>Molecular Oncology</i> , 2020 , 14, 69-86	6.9	14
86	HERC1 Ubiquitin Ligase Is Required for Hippocampal Learning and Memory. <i>Frontiers in Neuroanatomy</i> , 2020 , 14, 592797	3.6	2
85	The HERC1 ubiquitin ligase regulates presynaptic membrane dynamics of central synapses. <i>Scientific Reports</i> , 2020 , 10, 12057	4.9	5
84	A new homozygous HERC1 gain-of-function variant in MDFPMR syndrome leads to mTORC1 hyperactivation and reduced autophagy during cell catabolism. <i>Molecular Genetics and Metabolism</i> , 2020 , 131, 126-134	3.7	2
83	HERCing: Structural and Functional Relevance of the Large HERC Ubiquitin Ligases. <i>Frontiers in Physiology</i> , 2019 , 10, 1014	4.6	13
82	Interplay between BMPs and Reactive Oxygen Species in Cell Signaling and Pathology. <i>Biomolecules</i> , 2019 , 9,	5.9	19
81	Glucose Restriction Promotes Osteocyte Specification by Activating a PGC-1 β -Dependent Transcriptional Program. <i>iScience</i> , 2019 , 15, 79-94	6.1	12
80	Large HERCs Function as Tumor Suppressors. <i>Frontiers in Oncology</i> , 2019 , 9, 524	5.3	4
79	Inhibition of phosphatidylinositol 3-kinase (PI3K) prevents heterotopic ossification. <i>EMBO Molecular Medicine</i> , 2019 , 11, e10567	12	12
78	ACVR1 Function in Health and Disease. <i>Cells</i> , 2019 , 8,	7.9	16
77	Tris-Acetate Polyacrylamide Gradient Gels for the Simultaneous Electrophoretic Analysis of Proteins of Very High and Low Molecular Mass. <i>Methods in Molecular Biology</i> , 2019 , 1855, 269-277	1.4	0
76	Leucine reduces the proliferation of MC3T3-E1 cells through DNA damage and cell senescence. <i>Toxicology in Vitro</i> , 2018 , 48, 1-10	3.6	4
75	HERC1 Ubiquitin Ligase Is Required for Normal Axonal Myelination in the Peripheral Nervous System. <i>Molecular Neurobiology</i> , 2018 , 55, 8856-8868	6.2	10
74	The E3 ubiquitin ligase HERC1 controls the ERK signaling pathway targeting C-RAF for degradation. <i>Oncotarget</i> , 2018 , 9, 31531-31548	3.3	21

73	Mesenchymal stem cells improves survival in LPS-induced acute lung injury acting through inhibition of NETs formation. <i>Journal of Cellular Physiology</i> , 2017 , 232, 3552-3564	7	53
72	Liraglutide improves liver microvascular dysfunction in cirrhosis: Evidence from translational studies. <i>Scientific Reports</i> , 2017 , 7, 3255	4.9	41
71	Guaranã Highly Caffeinated Food, Presents in vitro Antitumor Activity in Colorectal and Breast Cancer Cell Lines by Inhibiting AKT/mTOR/S6K and MAPKs Pathways. <i>Nutrition and Cancer</i> , 2017 , 69, 800-810	2.8	16
70	Mesenchymal stem cells decrease lung inflammation during sepsis, acting through inhibition of the MAPK pathway. <i>Stem Cell Research and Therapy</i> , 2017 , 8, 289	8.3	29
69	TGF- β targets Smad, p38 MAPK, and PI3K/Akt signaling pathways to induce PFKFB3 gene expression and glycolysis in glioblastoma cells. <i>FEBS Journal</i> , 2017 , 284, 3437-3454	5.7	79
68	p53 inhibits SP7/Osterix activity in the transcriptional program of osteoblast differentiation. <i>Cell Death and Differentiation</i> , 2017 , 24, 2022-2031	12.7	34
67	Extracellular calcium promotes bone formation from bone marrow mesenchymal stem cells by amplifying the effects of BMP-2 on SMAD signalling. <i>PLoS ONE</i> , 2017 , 12, e0178158	3.7	43
66	NEURL4 regulates the transcriptional activity of tumor suppressor protein p53 by modulating its oligomerization. <i>Oncotarget</i> , 2017 , 8, 61824-61836	3.3	9
65	A nonsense variant in HERC1 is associated with intellectual disability, megalencephaly, thick corpus callosum and cerebellar atrophy. <i>European Journal of Human Genetics</i> , 2016 , 24, 455-8	5.3	32
64	Class I PI-3-Kinase Signaling Is Critical for Bone Formation Through Regulation of SMAD1 Activity in Osteoblasts. <i>Journal of Bone and Mineral Research</i> , 2016 , 31, 1617-30	6.3	11
63	Tris-acetate polyacrylamide gradient gel electrophoresis for the analysis of protein oligomerization. <i>Analytical and Bioanalytical Chemistry</i> , 2016 , 408, 1715-9	4.4	5
62	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016 , 12, 1-222	10.2	3838
61	Functional and pathological relevance of HERC family proteins: a decade later. <i>Cellular and Molecular Life Sciences</i> , 2016 , 73, 1955-68	10.3	52
60	Mesenchymal Stem Cells Within Gelatin/CaSO ₄ Scaffolds Treated Ex Vivo with Low Doses of BMP-2 and Wnt3a Increase Bone Regeneration. <i>Tissue Engineering - Part A</i> , 2016 , 22, 41-52	3.9	13
59	The HERC2 ubiquitin ligase is essential for embryonic development and regulates motor coordination. <i>Oncotarget</i> , 2016 , 7, 56083-56106	3.3	13
58	HERC 1 Ubiquitin Ligase Mutation Affects Neocortical, CA3 Hippocampal and Spinal Cord Projection Neurons: An Ultrastructural Study. <i>Frontiers in Neuroanatomy</i> , 2016 , 10, 42	3.6	18
57	Analysis of Protein Oligomerization by Electrophoresis. <i>Methods in Molecular Biology</i> , 2016 , 1449, 341-8	1.4	2
56	Therapeutic ultrasound stimulates MC3T3-E1 cell proliferation through the activation of NF- κ B, p38 and mTOR. <i>Lasers in Surgery and Medicine</i> , 2015 , 47, 765-72	3.6	10

55	Capsaicin modulates proliferation, migration, and activation of hepatic stellate cells. <i>Cell Biochemistry and Biophysics</i> , 2014 , 68, 387-96	3.2	12
54	The p38MAPK function in osteoprecursors is required for bone formation and bone homeostasis in adult mice. <i>PLoS ONE</i> , 2014 , 9, e102032	3.7	21
53	Functional EF-hands in neuronal calcium sensor GCAP2 determine its phosphorylation state and subcellular distribution in vivo, and are essential for photoreceptor cell integrity. <i>PLoS Genetics</i> , 2014 , 10, e1004480	6	12
52	Mitogen-activated protein kinase (MAPK)-regulated interactions between Osterix and Runx2 are critical for the transcriptional osteogenic program. <i>Journal of Biological Chemistry</i> , 2014 , 289, 27105-27117	5.4	76
51	The E3 ubiquitin protein ligase HERC2 modulates the activity of tumor protein p53 by regulating its oligomerization. <i>Journal of Biological Chemistry</i> , 2014 , 289, 14782-95	5.4	39
50	PFKFB3 activation in cancer cells by the p38/MK2 pathway in response to stress stimuli. <i>Biochemical Journal</i> , 2013 , 452, 531-43	3.8	51
49	Osterix induces Col1a1 gene expression through binding to Sp1 sites in the bone enhancer and proximal promoter regions. <i>Bone</i> , 2013 , 52, 548-56	4.7	54
48	Mutation of HERC2 causes developmental delay with Angelman-like features. <i>Journal of Medical Genetics</i> , 2013 , 50, 65-73	5.8	50
47	MicroRNA-322 (miR-322) and its target protein Tob2 modulate Osterix (Osx) mRNA stability. <i>Journal of Biological Chemistry</i> , 2013 , 288, 14264-14275	5.4	63
46	Akt-dependent activation of the heart 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase (PFKFB2) isoenzyme by amino acids. <i>Journal of Biological Chemistry</i> , 2013 , 288, 10640-51	5.4	50
45	Contribution of S6K1/MAPK signaling pathways in the response to oxidative stress: activation of RSK and MSK by hydrogen peroxide. <i>PLoS ONE</i> , 2013 , 8, e75523	3.7	14
44	Antiproliferative effect of catechin in GRX cells. <i>Biochemistry and Cell Biology</i> , 2012 , 90, 575-84	3.6	9
43	Progestins activate 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 3 (PFKFB3) in breast cancer cells. <i>Biochemical Journal</i> , 2012 , 442, 345-56	3.8	30
42	Tris-acetate polyacrylamide gradient gels for the simultaneous electrophoretic analysis of proteins of very high and low molecular mass. <i>Methods in Molecular Biology</i> , 2012 , 869, 205-13	1.4	15
41	Fructose-1,6-bisphosphate protects against Zymosan-induced acute lung injury in mice. <i>Inflammation</i> , 2012 , 35, 1198-203	5.1	7
40	Fructose-1,6-bisphosphate reduces the mortality in <i>Candida albicans</i> bloodstream infection and prevents the septic-induced platelet decrease. <i>Inflammation</i> , 2012 , 35, 1256-61	5.1	8
39	Sertoli-secreted FGF-2 induces PFKFB4 isozyme expression in mouse spermatogenic cells by activation of the MEK/ERK/CREB pathway. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012 , 303, E695-707	6	14
38	Treatment with N-methyl-D-aspartate receptor antagonist (MK-801) protects against oxidative stress in lipopolysaccharide-induced acute lung injury in the rat. <i>International Immunopharmacology</i> , 2011 , 11, 706-11	5.8	25

37	The transcriptional activation of the cyclooxygenase-2 gene in zymosan-activated macrophages is dependent on NF-kappa B, C/EBP, AP-1, and CRE sites. <i>Inflammation</i> , 2011 , 34, 653-8	5.1	16
36	Conserved regulatory motifs in osteogenic gene promoters integrate cooperative effects of canonical Wnt and BMP pathways. <i>Journal of Bone and Mineral Research</i> , 2011 , 26, 718-29	6.3	46
35	Noncanonical BMP signaling regulates cyclooxygenase-2 transcription. <i>Molecular Endocrinology</i> , 2011 , 25, 1006-17		22
34	Amino acids activate mammalian target of rapamycin complex 2 (mTORC2) via PI3K/Akt signaling. <i>Journal of Biological Chemistry</i> , 2011 , 286, 6128-42	5.4	132
33	The p38/MK2/Hsp25 pathway is required for BMP-2-induced cell migration. <i>PLoS ONE</i> , 2011 , 6, e16477	3.7	31
32	p38 regulates expression of osteoblast-specific genes by phosphorylation of osterix. <i>Journal of Biological Chemistry</i> , 2010 , 285, 31985-94	5.4	101
31	The combination of sirolimus and rosiglitazone produces a renoprotective effect on diabetic kidney disease in rats. <i>Life Sciences</i> , 2010 , 87, 147-53	6.8	21
30	Simultaneous electrophoretic analysis of proteins of very high and low molecular mass using Tris-acetate polyacrylamide gels. <i>Electrophoresis</i> , 2010 , 31, 1318-21	3.6	38
29	Progressive Purkinje cell degeneration in tambaleante mutant mice is a consequence of a missense mutation in HERC1 E3 ubiquitin ligase. <i>PLoS Genetics</i> , 2009 , 5, e1000784	6	44
28	The RCC1 superfamily: from genes, to function, to disease. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2008 , 1783, 1467-79	4.9	82
27	ERK and p38 pathways regulate amino acid signalling. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2008 , 1783, 2241-54	4.9	38
26	BMP-2 induces Osterix expression through up-regulation of Dlx5 and its phosphorylation by p38. <i>Journal of Biological Chemistry</i> , 2008 , 283, 3816-26	5.4	172
25	BMP2 induction of actin cytoskeleton reorganization and cell migration requires PI3-kinase and Cdc42 activity. <i>Journal of Cell Science</i> , 2008 , 121, 3960-70	5.3	89
24	Simultaneous electrophoretic analysis of proteins of very high and low molecular weights using low-percentage acrylamide gel and a gradient SDS-PAGE gel. <i>Electrophoresis</i> , 2006 , 27, 3935-8	3.6	16
23	TSC1 stabilizes TSC2 by inhibiting the interaction between TSC2 and the HERC1 ubiquitin ligase. <i>Journal of Biological Chemistry</i> , 2006 , 281, 8313-6	5.4	169
22	Requirement of phosphatidylinositol-4,5-bisphosphate for HERC1-mediated guanine nucleotide release from ARF proteins. <i>FEBS Letters</i> , 2005 , 579, 343-8	3.8	11
21	The HERC proteins: functional and evolutionary insights. <i>Cellular and Molecular Life Sciences</i> , 2005 , 62, 1826-38	10.3	58
20	6-Phosphofructo-2-kinase (pfkfb3) gene promoter contains hypoxia-inducible factor-1 binding sites necessary for transactivation in response to hypoxia. <i>Journal of Biological Chemistry</i> , 2004 , 279, 53562-70	5.4	171

19	BMP-2 decreases Mash1 stability by increasing Id1 expression. <i>EMBO Journal</i> , 2004 , 23, 3527-37	13	87
18	The giant protein HERC1 is recruited to aluminum fluoride-induced actin-rich surface protrusions in HeLa cells. <i>FEBS Letters</i> , 2004 , 559, 77-83	3.8	8
17	Interaction between HERC1 and M2-type pyruvate kinase. <i>FEBS Letters</i> , 2003 , 539, 78-84	3.8	31
16	Regulation of ubiquitous 6-phosphofructo-2-kinase by the ubiquitin-proteasome proteolytic pathway during myogenic C2C12 cell differentiation. <i>FEBS Letters</i> , 2003 , 550, 23-9	3.8	30
15	Direct binding of Smad1 and Smad4 to two distinct motifs mediates bone morphogenetic protein-specific transcriptional activation of Id1 gene. <i>Journal of Biological Chemistry</i> , 2002 , 277, 3176-85	5.4	239
14	Inhibition of PI3K/p70 S6K and p38 MAPK cascades increases osteoblastic differentiation induced by BMP-2. <i>FEBS Letters</i> , 2002 , 510, 99-104	3.8	103
13	The human ubiquitous 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene (PFKFB3): promoter characterization and genomic structure. <i>Gene</i> , 2001 , 264, 131-8	3.8	32
12	HERC3 binding to and regulation by ubiquitin. <i>FEBS Letters</i> , 2001 , 488, 74-80	3.8	23
11	Interaction and functional cooperation of NF-kappa B with Smads. Transcriptional regulation of the junB promoter. <i>Journal of Biological Chemistry</i> , 2000 , 275, 28937-46	5.4	90
10	Effect of protein kinase A activity on the association of ADP-ribosylation factor 1 to golgi membranes. <i>Journal of Biological Chemistry</i> , 2000 , 275, 19050-9	5.4	29
9	A zinc-finger transcription factor induced by TGF-beta promotes apoptotic cell death in epithelial Mv1Lu cells. <i>FEBS Letters</i> , 1999 , 457, 478-82	3.8	84
8	JunB is involved in the inhibition of myogenic differentiation by bone morphogenetic protein-2. <i>Journal of Biological Chemistry</i> , 1998 , 273, 537-43	5.4	89
7	Expression of the F-type 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase mRNA during liver regeneration. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1997 , 1334, 256-60	4	3
6	A giant protein that stimulates guanine nucleotide exchange on ARF1 and Rab proteins forms a cytosolic ternary complex with clathrin and Hsp70. <i>Oncogene</i> , 1997 , 15, 1-6	9.2	54
5	Hepatocyte growth factor and transforming growth factor beta regulate 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene expression in rat hepatocyte primary cultures. <i>Biochemical Journal</i> , 1996 , 314 (Pt 1), 235-40	3.8	5
4	Calmodulin expression during rat liver regeneration. <i>Hepatology</i> , 1994 , 20, 1002-8	11.2	12
3	c-met mRNA overexpression in human hepatocellular carcinoma 1994 , 19, 88		5
2	Effect of galactosamine on hepatic carbohydrate metabolism: protective role of fructose 1,6-bisphosphate. <i>Hepatology</i> , 1992 , 15, 1147-53	11.2	65

- 1 Fructose 2,6-bisphosphate in hypoglycemic rat brain. *Journal of Neurochemistry*, **1991**, 57, 200-3 6 14